

*Short Communication***Methylated DNA Collected by Tampons—A New Tool to Detect Endometrial Cancer**

Heidi Fiegl,¹ Conny Gatringer,¹ Andreas Widschwendter,¹ Alois Schneitter,¹ Angela Ramoni,¹ Daniela Sarlay,¹ Inge Gaugg,¹ Georg Goebel,² Hannes M. Müller,¹ Elisabeth Mueller-Holzner,¹ Christian Marth,¹ and Martin Widschwendter¹

¹Department of Obstetrics and Gynecology, Innsbruck University Hospital and ²Department of Biostatistics and Documentation, University of Innsbruck, Innsbruck, Austria

Abstract

This proof of principle study aimed to define a new and simple strategy for detection of endometrial cancer using epigenetic markers. We investigated DNA isolated from vaginal secretion collected from tampon for aberrant methylation of five genes (*CDH13*, *HSPA2*, *MLH1*, *RASSF1A*, and *SOCS2*) using MethyLight in 15 patients with endometrial cancer and 109 patients without endometrial cancer. All endometrial cancer

patients revealed three or more methylated genes, whereas 91% (99 of 109) of the patients without endometrial cancer had no or fewer than three genes methylated in their vaginal secretion. The methods developed in this study provide the basis for a prospective clinical trial to screen asymptomatic women who are at high risk for endometrial cancer. (Cancer Epidemiol Biomarkers Prev 2004;13(5):882–8)

Introduction

Endometrial cancer is one of the most common genital cancers in women worldwide. The highest incidence rates are observed in western Europe and North America (1). The well-known risk factors for endometrial cancer include obesity, type 2 diabetes mellitus, and hypertension. Additionally, anovulation and long-term use of unopposed estrogens for hormone replacement therapy increase the risk for endometrial cancer. Genetic causes of endometrial cancer are uncommon, although there is an association with hereditary nonpolyposis colon cancer syndrome, in which the individual risk rises to a cumulative incidence of 40% by age 70 years (2). In 2001, the American Cancer Society concluded that there was insufficient evidence to recommend screening for endometrial cancer for women at average risk or increased risk due to history of unopposed estrogen therapy, nulliparity, infertility or failure to ovulate, obesity, diabetes, or hypertension (3). Studies examining endometrial carcinoma screening methods for asymptomatic postmenopausal women have used ultrasound-determined endometrial thickness as an indication of risk. Transvaginal ultrasonography compared with en-

dometrial biopsy for the detection of endometrial disease had a positive predictive value of only 9% for detecting any abnormality, with 90% sensitivity and 48% specificity (4). There is a need for a sensitive and specific screening test for high-risk women. It has been shown that genetic abnormalities can be used to detect endometrial cancer (5). Changes in the status of DNA methylation are among the most common molecular alterations in human neoplasias (6). It has been increasingly recognized over the past 4–5 years that the CpG islands of a large number of genes, which are unmethylated in normal tissue, are methylated to varying degrees in multiple types of human cancer (6, 7). Aberrant methylation of CpG islands within the promoter regions of several genes such as *E-cadherin*, *adenomatous polyposis coli* (*APC*), *MLH1*, *p16*, *estrogen receptor*, *progesterone receptor*, and *PTEN* (*MMAC1*) has been identified in endometrial cancer tissue (8–14). Up to now, no investigations have been undertaken to assess the methylation status of DNA obtained from cervicovaginal secretion from endometrial cancer patients.

Our proof of principle study was performed to determine whether it is possible to detect endometrial cancer by analyzing methylated DNA in cervicovaginal secretion.

Materials and Methods

Patients and Samples. One hundred twenty-four patients were recruited for this study: 15 patients had endometrial cancer, while the no endometrial cancer group contained 5 patients with invasive cervical cancer, 35 patients with cervical intraepithelial neoplasia

Received 9/24/03; revised 11/18/03; accepted 12/24/03.

Grant support: "Fonds zur Förderung der Wissenschaftlichen Forschung," P15995-B05 and P16159-B05.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Note: H. Fiegl, C. Gatringer, and A. Widschwendter contributed equally to this work.

Requests for reprints: Martin Widschwendter, Department of Obstetrics and Gynecology, Innsbruck University Hospital, Anichstrasse 35, A-6020 Innsbruck, Austria. Phone: 43-512-504-4155; Fax: 43-512-504-3112. E-mail: martin.widschwendter@uibk.ac.at

(CIN I, 3 cases; CIN II, 19 cases; CIN III, 13 cases), and 69 patients with benign disease of the uterus. Histology of the endometrial cancers was adenocarcinoma in 12 cases, adenosquamous carcinoma in 2 cases, and malignant mixed mullerian tumor in 1 case. Most of the carcinomas (10 cases) were well differentiated (tumor grade 1). Only four carcinomas had tumor grade 2. All FIGO stages were represented in the endometrial cancer group (FIGO I, 8 patients; FIGO II, 2 patients; FIGO III, 3 patients; FIGO IV, 1 patient; unknown FIGO stage, 1 patient). Sample collection was done between January 1, 2003 and May 31, 2003 at the Department of Obstetrics and Gynecology, Innsbruck University Hospital, Austria. All patients who were scheduled to undergo surgery of the uterus on the next day including a histological diagnosis were invited to attend the study. Samples and clinical data were collected after informed consent was obtained. To ensure standardized sample collection, a tampon was inserted in the patient by a physician after speculum examination and retained intravaginal for 30 min. Preparation of the samples is shown in Fig. 1.

DNA Isolation and Methylation Analysis. Genomic DNA from samples was isolated using the High Pure Viral Nucleic Acid Kit (Roche Diagnostics, Mannheim, Germany) according to the manufacturer's protocol with some modifications for multiple loading of the DNA extraction columns to gain a sufficient amount of DNA. Sodium bisulfite-treated genomic DNA was analyzed by means of MethyLight, a fluorescence-based, real-time PCR assay, as described previously (15, 16). Briefly, two sets of primers and probes, designed specifically for bisulfite-converted DNA, were used: a methylated set for the gene of interest and a reference set, β -actin (*ACTB*), to normalize for input DNA. Specificity of the reactions for methylated DNA was confirmed separately using *SssI* (New England Biolabs, Frankfurt, Germany)-treated human WBC DNA (heavily methylated). The percentage of fully methylated reference (PMR) molecules at a specific locus was calculated by dividing the *GENE:ACTB* ratio of a sample by the *GENE:ACTB* ratio of *SssI*-treated WBC DNA and multiplying by 100. PMR indicates this measurement. A gene was deemed methylated if the PMR value was >0 . Primer and probes specific for methylated DNA and used for MethyLight reactions are listed in Table 1.

Statistical Analysis. Associations between categorical variables were tested using the χ^2 test. Differences in median of age were examined with the Mann-Whitney *U* test or, between more than two groups, with the Kruskal-Wallis test. Due to a significant age difference between endometrial cancer patients and the no endometrial cancer group, an age-matched group of the no endometrial cancer group with a matching ratio 1:2 was randomly selected. The computation of the matching group was done using MATLAB R12 (<http://www.mathworks.com>). For determination of diagnostic accuracy, a nonparametric receiver operating curve analysis with linear interpolation was performed. $P > 0.05$ was considered statistically significant. All statistical calculations were performed using SPSS version 11.0 for Windows.




Figure 1. Flowchart of procedures for tampon insertion and sample preparation.

Results

Aberrant methylation of 38 genes in DNA obtained from vaginal secretion from the first five patients with endometrial cancer and the first four patients with benign disease was analyzed to determine appropriate genes for further study. The most appropriate genes for our further

Table 1. Primer and probes specific for methylated DNA and used for MethyLight reactions

HUGO gene nomenclature	Forward primer sequence	Reverse primer sequence	Probe oligo sequence
<i>ACTB</i>	TGGTGATGGAGGAGGTTTAG TAAGT	AACCAATAAAACCTACTCCTC CCTTAA	6FAM-ACCACCACCCAACACACAATA ACAAACACA-BHQ-1
<i>APC</i>	GAACCAAAACGCTCCCAT	TTATATGTGGTACGTGCGTT TATAT	6FAM-CCCGTCGAAAACCCGCCGA TTA-BHQ-1
<i>ARHI</i>	GCGTAAGCGGAATTTATGT TTGT	CCGCGATTTATATTCGACTT	6FAM-CGCACAAAAACGAAATACGAA AACGCAAA-BHQ-1
<i>BLT1</i>	GCGTTGGTTTTATCGGAAGG	AAACCGTAATTCGGCTCG	6FAM-GACTCCGCCAACTTCGCCAA AA-BHQ-1
<i>BRCA1</i>	GAGAGGTTGTTGTTAGCGG TAGTT	CGCGCAATCGCAATTTTAAAT	6FAM-CCGCGCTTTTCGGTACCA CGA-BHQ-1
<i>CALCA</i>	GTTTTGGAAGTATGAGGGTG ACG	TTCCC GCCGTATAAATCG	6FAM-ATTCCGCCAATACACAACA ACCAATAACG-BHQ-1
<i>CDH1</i>	AATTTTAGGTTAGAGGGTT ATCGCGT	TCCCCAAAACGAACTAAC GAC	6FAM-CGCCCACCCGACCTCGCAT- BHQ-1
<i>CDH13</i>	AATTCGTTGTTTTGTGCGT	CTACCCGTACCGAACGATCC	6FAM-AACGCAAAACGCGCCCGACA- BHQ-1
<i>CDKN2A</i>	TGGAGTTTTCGGTTGATTGGTT	AACAACGCCCGCACCTCT	6FAM-ACCCGACCCCGAACCGCG-BHQ-1
<i>CYP11B1</i>	GTGCGTTTGACGGGAGTT	AACGCGACCTAACAAAAC GAA	6FAM-CGCCGCACACCAAACCCGC TT-BHQ-1
<i>DAPK1</i>	TCGTCGTCGTTTCGGTTAGTT	TCCCTCCGAAACGCTATCG	6FAM-CGACCATAAACGCCAACG CCG-BHQ-1
<i>ESR1</i>	GGCGTTCGTTTTGGGATTG	GCCGACACGCGAACTCTAA	6FAM-CGATAAAACCGAACGACCC GACGA-BHQ-1
<i>ESR2</i>	TTTGAAATTTGAGGGCGAA GAGTAG	ACCCGTCGCAACTCGAATAA	6FAM-CCGACCCAACGCTCGCCG-BHQ-1
<i>FGF18</i>	ATCTCCTCCTCCGCGTCTCT	TCGCGGTAGAAAACGTTT	6FAM-CGACCGTACGCATCGCCGC-BHQ-1
<i>GSTM3</i>	GCGGAACGCCCTAACT	AACGTCGGTATTAGTCGCGTTT	6FAM-CCCGTCTCCTCCCTTACCTCC- BHQ-1
<i>GSTP1</i>	GTCGGCGTCGTGATTTAGT ATTG	AAACTACGACGACGAAAC TCCAA	6FAM-AAACCTCGCGACCTCCGAACCTTA TAAAA-BHQ-1
<i>HIC1</i>	GTTAGGCGGTTAGGGCGTC	CCGAACGCCTCCATCGTAT	6FAM-CAACATCGTCTACCCAACACACTC TCCTACG-BHQ-1
<i>HLA-G</i>	CACCCCATATACGCGCTAA	GGTCGTTACGTTTCGGGTAG TTTA	6FAM-CGCGCTCACACGCTCAAAAACCT- BHQ-1
<i>HSD17B4</i>	TATCGTTGAGGTTGACGGG	TCCAACCTTCGCATACTACC	6FAM-CCCGCGCCGATAACCAATACCA- BHQ-1
<i>HSPA2</i>	CACGAACACTACCAACAAC TCAACT	GGGAGCGGATTGGGTTTG	6FAM-CCGCGCCCAATTCGGATTCT- BHQ-1
<i>IGFBP2</i>	CTCGCGCCGACAAATAAATAC	CGGGAAGAGTAGGGAATTT TTAGAGT	6FAM-ACGCCCGCTCGCCACCT-BHQ-1
<i>MGMT</i>	GCGTTTCGACGTTCTGAGGT	CACCTTCCGAAAACGAAACG	6FAM-CGCAAACGATACGCACCCGGA- BHQ-1
<i>MLH1</i>	AGGAAGAGCGGATAGCGATT	TCTTCGTCCTCCCTAAAACG	6FAM-CCCGCTACCTAAAAAATATACG CTTACGCG-BHQ-1
<i>MLLT7</i>	CCTCACGATACCTCCCTCAA	TTAGGGATTAGCGTTTTGG GATT	6FAM-AAACACATTCCTACCAATCTTC AAAAATCGCG-BHQ-1
<i>MT3</i>	CGATAAACGAACTTCTCCA AACAA	GCGCGGTGCGTAGGG	6FAM-AAACGCGCGACTTAACTAATA ACAACAATAACGA-BHQ-1
<i>MYOD1</i>	GAGCGCGGTAGTTAGCG	TCCGACACGCCCTTCC	6FAM-CTCCAACACCCGACTACTATAT CCGCGAAA-BHQ-1
<i>PGR</i>	TTATAATTCGAGGCGGTTAG TGTTT	TCGAACTTCTACTAATCC GTAACGTA	6FAM-ATCATCTCCGAAAATCTCAAT CCCAATAACG-BHQ-1
<i>PPP1R13B</i>	CCTCACCCACCGACATCATC	TCGGAGCGGTGGGTATAGTTC	6FAM-AAAAATCCGCGACGCCCTCGA- BHQ-1
<i>PTGS2</i>	CGGAAGCGTTCGGGTAAAG	AATTCACCCGCCCAAAC	6FAM-TTCCGCCAAATATCTTTTCTTC TTCGCA-BHQ-1
<i>RASSF1A</i>	ATTGAGTTGCGGGAGTTGGT	ACACGCTCCAACCGAATACG	6FAM-CCCTTCCCAACCGCCCA-BHQ-1
<i>SOCS1</i>	GCGTCGAGTTCGTGGGTATTT	CCGAAACCATCTTCACGCTAA	6FAM-ACAATTCGGTAAACGACTATC GCGCA-BHQ-1
<i>SOCS2</i>	TCCCTTCCCCGCCATT	TTGTTTTGTGCGGGTATT	6FAM-CCGAAAAACTCAAAACCCGCAA AATCATBHQ-1
<i>SYK</i>	GGGCGGATATTGGGAG	GCGACTTCTCTATTTTAA CAAC	6FAM-CCTTAAACGCGCCGAAACAAACG- BHQ-1
<i>TERT</i>	GGATTGCGGGTATAGACGTT	CGAAATCCGCGCGAAA	6FAM-CCCAATCCCTCCGCCACGTAAAA- BHQ-1
<i>TFF1</i>	TAAGGTTACGGTGGTTATT TCGTGA	ACCTTAATCCAAATCCTACT CATATCTAAAA	6FAM-CCCTCCGCCAAAATAAATACTAT ACTACTACAAAA-BHQ-1
<i>TIMP3</i>	GCGTCGGAGGTTAAGGTTGTT	CTCTCCAAAATTACCGTACGCG	6FAM-AACTCGTCCGCCGCCGAA- BHQ-1

(Continued on following page)

Table 1. Primer and probes specific for methylated DNA and used for MethyLight reactions (Cont'd)

HUGO gene nomenclature	Forward primer sequence	Reverse primer sequence	Probe oligo sequence
<i>TITF1</i>	CGAAATAAACCGAATCCTCC TTAA	TGTTTGTGTTTITAGCGTTTACGT	6FAM-CTCGCGTTTATTTTAACCCG ACCCA-BHQ-1
<i>TP53BP2</i>	ACCCCTAACGCGACTTTATC	GTTTCGATTCGGGATTAGTTGGT	6FAM-CGCTCGTAACGATCGAAACT CCCTCCT-BHQ-1
<i>TWIST</i>	GTAGCGCGGCGAACGT	AAACGCAACGAATCATAACCAAC	6FAM-CCAACGCACCCAATCGCTAAA CGA-BHQ-1

analyses were determined to be those that revealed the greatest difference in PMR values between patients with benign disease of the uterus and endometrial cancer patients (Fig. 2). Five genes (*i.e.*, *RASSF1A*, *hMLH1*, *CDH13*, *HSPA2*, and *SOCS2*) were selected for further

analysis. DNA methylation in three or more of these five genes was observed in cervicovaginal secretion of all five patients with endometrial cancer, whereas all four patients without endometrial cancer showed no or fewer than three genes to be methylated. We therefore





Figure 2. PMR values of 38 genes in patients without endometrial cancer ($n = 4$) and with endometrial cancer ($n = 5$). Arrows, genes with the greatest difference in PMR values.



*cervical intraepithelial neoplasia grade III (CIN III); † invasive cervical cancer

Figure 3. A. Methylation status of the five investigated genes (full box, methylated; empty box, unmethylated), status (endometrial cancer or no endometrial cancer), and age. B. Receiver operating curve for detection of endometrial cancer by methylation analysis of DNA obtained from vaginal secretion for patients aged between 50 and 75 years (excluding patients with CIN III or cervical cancer).

determined the cutoff value between no endometrial cancer and endometrial cancer as methylation positive in three or more of the five investigated genes.

The overwhelming majority of the patients without endometrial cancer (99 of 109) revealed no or fewer than three genes methylated, whereas all of the 15 endometrial cancer patients had three or more genes methylated in their vaginal secretion ($P < 0.001$, χ^2 test; Fig. 3A). Histological examination of the 10 patients in the no endometrial cancer group with three or more genes methylated revealed invasive cervical cancer (four cases), CIN III (one case), endometrium polyp (four cases), and fibroids (one case). Samples were collected after primary surgery (curettage, punch biopsy of the cervix, or hysteroscopic operation) and before secondary surgery (hysterectomy) in 16 of 124 patients: 9 of 16 patients had endometrial cancer, 3 of 16 patients had CIN III, and 4 of 16 patients had benign disease of the endometrium. All nine endometrial cancer patients had a diagnosis of residual cancer at the time of secondary surgery. All nine endometrial cancer patients had three or more genes methylated, the three CIN III patients revealed no methylated genes, and one of the patients with benign disease showed one gene to be methylated. Within the group of patients from whom the vaginal secretion was collected prior to any surgery, one patient presented due to sonographically detected serometra with complete stenosis of the cervicouterine canal. Even this patient showed methylation of three of the five tested genes. DNA methylation of the five genes identified seems to increase with age although statistically not significant (data not shown). Using all 15 endometrial cancer cases and 109 controls, the area under the receiver operating curve was 0.973 (data not shown). After exclusion of the nine pilot cases (five endometrial cancer patients and four patients without endometrial cancer), which were initially used to determine appropriate genes for further analyses, the area under the receiver operating curve was 0.963. To rule out the possibility that abnormal DNA methylation is merely a surrogate for age rather than a cancer-specific marker, we randomly age matched two nonendometrial cancer controls for each endometrial cancer case. Investigation of DNA methylation in the cervicovaginal secretion of these 45 patients was still able to discriminate between endometrial cancer and patients without endometrial cancer ($P < 0.001$, χ^2 test) with a sensitivity of 100% and a specificity of 80%.

When analyzing all patients aged between 50 and 75 years and excluding patients with CIN III or cervical cancer, sensitivity was 100% and specificity rose to 97.2% (Fig. 3B). In this group, only 1 of 35 samples was false positive. The same sensitivity and specificity values were obtained after exclusion of the pilot cases from these analyses.

Discussion

Previous studies have described the importance of DNA methylation in human cancers. Recently, aberrant methylation of various genes has been identified in endometrial cancer tissue (8–14). Methylated DNA has been investigated as a possible screening marker for neoplastic disease in several body fluids (17). However,

up to now, no investigations have been undertaken to assess the methylation status of DNA obtained from cervicovaginal secretion from endometrial cancer patients.

In our study, all endometrial cancer patients revealed three or more of the five investigated genes methylated, whereas 99 of 109 patients without endometrial cancer had no or fewer than three genes methylated. Four of 10 patients in the no endometrial cancer group with three or more genes methylated had invasive cervical cancer. These cases indicate that some cervical cancer patients can also be identified with this assay.

In some cases (16 of 124), samples were collected after primary surgery and before secondary surgery. All endometrial cancer patients in this group had three or more genes methylated. These results demonstrate that aberrant methylation analysis can detect endometrial cancer even after primary surgery.

As endometrial cancer is more prevalent in older women and abnormal DNA methylation in nonmalignant tissues seems to increase with age (18), we especially addressed this problem within this project. Comparison of DNA methylation in the cervicovaginal secretion of endometrial cancer patients and age-matched nonendometrial cancer controls revealed still highly significant differences between these two groups.

Endometrial cancer occurs in almost all cases after menopause. Therefore, we analyzed all patients aged between 50 and 75 years and excluded patients with CIN III or invasive cervical cancer. These patients represent the group that will benefit from an endometrial cancer screening assay. In these group, the sensitivity and specificity to detect patients with endometrial cancer was 100% and 97.2%, respectively.

The methods developed in this study, involving tampons to collect cervicovaginal secretion, DNA isolation, and a high-throughput method to analyze DNA methylation, provide the basis for a prospective clinical trial to screen asymptomatic women who are at high risk for endometrial cancer.

Acknowledgments

We thank the staff of the Department of Obstetrics and Gynecology, Innsbruck University Hospital, for patient management.

References

1. Parkin DM, Pisani P, Ferlay J. Estimates of the worldwide incidence of 25 major cancers in 1990. *Int J Cancer* 1999;80:827-41.
2. Vasen HF, Wijnen JT, Menko FH, et al. Cancer risk in families with hereditary nonpolyposis colorectal cancer diagnosed by mutation analysis. *Gastroenterology* 1996;110:1020-7.
3. Smith RA, von Eschenbach AC, Wender R, et al. American Cancer Society guidelines for the early detection of cancer: update of early detection guidelines for prostate, colorectal, and endometrial cancers. Also: Update 2001—testing for early lung cancer detection. *CA Cancer J Clin* 2001;51:38-75.
4. Langer RD, Pierce JJ, O'Hanlan KA, et al. Transvaginal ultrasonography compared with endometrial biopsy for the detection of endometrial disease. Postmenopausal Estrogen/Progestin Interventions Trial. *N Engl J Med* 1997;337:1792-8.
5. Al Jehani RM, Jeyarajah AR, Hagen B, et al. Model for the molecular genetic diagnosis of endometrial cancer using K-ras mutation analysis. *J Natl Cancer Inst* 1998;90:540-2.
6. Jones PA. DNA methylation errors and cancer. *Cancer Res* 1996;56:2463-7.
7. Jones PA, Laird PW. Cancer epigenetics comes of age. *Nat Genet* 1999;21:163-7.
8. Saito T, Nishimura M, Yamasaki H, Kudo R. Hypermethylation in

- promoter region of E-cadherin gene is associated with tumor dedifferentiation and myometrial invasion in endometrial carcinoma. *Cancer* 2003;97:1002-9.
9. Zysman M, Saka A, Millar A, Knight J, Chapman W, Bapat B. Methylation of adenomatous polyposis coli in endometrial cancer occurs more frequently in tumors with microsatellite instability phenotype. *Cancer Res* 2002;62:3663-6.
 10. Esteller M, Levine R, Baylin SB, Ellenson LH, Herman JG. MLH1 promoter hypermethylation is associated with the microsatellite instability phenotype in sporadic endometrial carcinomas. *Oncogene* 1998;17:2413-7.
 11. Tsuda H, Yamamoto K, Inoue T, Uchiyama I, Umesaki N. The role of p16-cyclin d/CDK-pRb pathway in the tumorigenesis of endometrioid-type endometrial carcinoma. *Br J Cancer* 2000;82:675-82.
 12. Sasaki M, Kotcherguina L, Dharia A, Fujimoto S, Dahiya R. Cytosine-phosphoguanine methylation of estrogen receptors in endometrial cancer. *Cancer Res* 2001;61:3262-6.
 13. Sasaki M, Dharia A, Oh BR, Tanaka Y, Fujimoto S, Dahiya R. Progesterone receptor B gene inactivation and CpG hypermethylation in human uterine endometrial cancer. *Cancer Res* 2001;61:97-102.
 14. Salvesen HB, MacDonald N, Ryan A, et al. PTEN methylation is associated with advanced stage and microsatellite instability in endometrial carcinoma. *Int J Cancer* 2001;91:22-6.
 15. Eads CA, Danenberg KD, Kawakami K, et al. MethyLight: a high-throughput assay to measure DNA methylation. *Nucleic Acids Res* 2000;28:E32.
 16. Eads CA, Lord RV, Wickramasinghe K, et al. Epigenetic patterns in the progression of esophageal adenocarcinoma. *Cancer Res* 2001;61:3410-8.
 17. Muller HM, Widschwendter M. Methylated DNA as a possible screening marker for neoplastic disease in several body fluids. *Exp Rev Mol Diagn* 2003;3:443-58.
 18. Toyota M, Issa JP. CpG island methylator phenotypes in aging and cancer. *Semin Cancer Biol* 1999;9:349-57.

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Methylated DNA Collected by Tampons—A New Tool to Detect Endometrial Cancer

Heidi Fiegl, Conny Gatringer, Andreas Widschwendter, et al.

Cancer Epidemiol Biomarkers Prev 2004;13:882-888.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/13/5/882>

Cited articles This article cites 17 articles, 5 of which you can access for free at:
<http://cebp.aacrjournals.org/content/13/5/882.full#ref-list-1>

Citing articles This article has been cited by 7 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/13/5/882.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and
Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications
Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/13/5/882>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's
(CCC)
Rightslink site.