

Cervical Cancer Control Research in Vietnamese American Communities

Victoria M. Taylor,^{1,2} Tung T. Nguyen,⁴ J. Carey Jackson,³ and Stephen J. McPhee⁴

¹Cancer Prevention Program, Division of Public Health Sciences, Fred Hutchinson Cancer Research Center; ²Department of Health Services, School of Public Health, and ³Division of General Internal Medicine, Department of Medicine, University of Washington, Seattle, Washington and ⁴Division of General Internal Medicine, Department of Medicine, University of California at San Francisco, San Francisco, California

Abstract

Census data show that the U.S. Vietnamese population now exceeds 1,250,000. Cervical cancer among Vietnamese American women has been identified as an important health disparity. Available data indicate the cervical cancer disparity may be due to low Papanicolaou (Pap) testing rates rather than variations in human papillomavirus infection rates and/or types. The cervical cancer incidence rates among Vietnamese and non-Latina White women in California during 2000 to 2002 were 14.0 and 7.3 per 100,000, respectively. Only 70% of Vietnamese women who participated in the 2003 California Health Interview Survey reported a recent Pap smear compared with 84% of non-Latina White women. Higher levels of cervical cancer screening participation among Viet-

namese women are strongly associated with current/previous marriage, having a usual source of care/doctor, and previous physician recommendation. Vietnamese language media campaigns and lay health worker intervention programs have been effective in increasing Pap smear use in Vietnamese American communities. Cervical cancer control programs for Vietnamese women should address knowledge deficits, enable women who are without a usual source of care to find a primary care doctor, and improve patient-provider communication by encouraging health-care providers to recommend Pap testing as well as by empowering women to ask for testing. (Cancer Epidemiol Biomarkers Prev 2008;17(11):2924–30)

Introduction

The majority of Vietnamese Americans came to the United States as refugees or immigrants over the last three decades (1). According to the 2004 American Community Survey, 11% of Asian Americans are of Vietnamese descent and the Vietnamese population now exceeds 1,250,000 (2). Vietnamese are the second fastest-growing Asian American group after South Asian Indians (1). California, Texas, and Washington State have the largest Vietnamese communities (3). Compared with the general U.S. population and all Asian Americans, Vietnamese Americans are economically disadvantaged and linguistically isolated (Table 1). Over one-third (38%) have less than a high school education and nearly two-thirds (62%) have limited English proficiency (4).

The President's Advisory Commission on Asian Americans recently identified cervical cancer among

Vietnamese women as one of the most important health disparities experienced by Asian American populations (5). Human papillomavirus (HPV) infection is a universal risk factor for cervical cancer (6, 7). Although little is known about HPV infection among Vietnamese American women, available data indicate that the cervical cancer disparity may be due to low Papanicolaou (Pap) testing rates rather than variations in HPV infection rates and/or types (8, 9).

American Cancer Society guidelines specify that women should be screened for cervical cancer every 1 to 3 years depending on their risk factors for disease and previous screening history (10). Further, national cervical cancer screening goals for the year 2010 specify that at least 97% of women will have been screened on at least one occasion, and 90% will have received a Pap smear within the previous 3 years (11). In this review article, we present cervical cancer incidence data for Vietnamese women; provide information about levels of Pap testing, factors associated with Pap smear receipt, and cervical cancer control intervention programs; and summarize limitations of the available data.

Cervical Cancer Incidence

Cervical cancer incidence data for Vietnamese populations are summarized in Table 2 (6, 12–15). Twelve years ago, the Surveillance, Epidemiology and End Results

Received 4/29/08; revised 7/1/08; accepted 7/15/08.

Grant support: National Cancer Institute (The Asian American Network for Cancer Awareness, Research, and Training Community Network Program) grant U01-CA-114640, Centers for Disease Control and Prevention (The Vietnamese REACH for Health Initiative) cooperative agreement U50-CCU-922156, National Cancer Institute (Cervical Cancer Control in Vietnamese Women) grant R01-CA-115564, and Centers for Disease Control and Prevention (The Alliance for Reducing Cancer Northwest) cooperative agreement U48-DP-000050.

Requests for reprints: Victoria M. Taylor, Cancer Prevention Program, Division of Public Health Sciences, Fred Hutchinson Cancer Research Center (M3-B232), 1100 Fairview Avenue North, Seattle, WA 98109. Phone: 206-667-5114; Fax: 206-667-5977. E-mail: vtaylor@fhcrc.org

Copyright © 2008 American Association for Cancer Research.
doi:10.1158/1055-9965.EPI-08-0386

Table 1. Characteristics of selected U.S. populations, 2000

Characteristics	Vietnamese (%)	Asian (%)	U.S. (%)
Less than high school education*	38	20	20
Below poverty level	16	13	12
Foreign-born	76	69	11
Speak language other than English at home	93	79	18
Do not speak English very well	62	40	8

*People ages ≥ 25 years.

program published a monograph addressing racial/ethnic patterns of cancer in the United States. This monograph showed that Vietnamese women had higher rates of invasive cervical cancer than any other racial/ethnic group. For the 1988 to 1992 period, the age-adjusted invasive cervical cancer incidence rate among Vietnamese women was 43.0 per 100,000 compared with 7.5 per 100,000 among non-Latina White women, 13.2 per 100,000 among Black women, and 16.2 per 100,000 among Latina women (14).

More recent cancer registry data indicate that cervical cancer incidence rates among Vietnamese Americans have gone down steadily during the last decade (15, 16). However, the incidence rate among Vietnamese women in California was still nearly twice the incidence rate among non-Latina White women during 2000 to 2002 (14.0 versus 7.3 per 100,000; ref. 6). In summary, available data show that Vietnamese American women experience a cervical cancer disparity relative to other racial/ethnic groups, but this disparity has decreased over time.

Pap Testing Levels

McPhee and Nguyen previously summarized findings from population-based surveys of Pap testing use among Vietnamese women, conducted before 2000 (17). These California and Massachusetts surveys found that only about one-half (between 43% and 53%) of Vietnamese women ages ≥ 18 years had ever received a Pap smear (18-21). Findings from more recent population-based surveys of Vietnamese women are given in Table 3 (22-27). Surveys conducted in Texas during 2000, 2004, and 2007; California during 2000; and Washington during 2002 consistently found that approximately three-quarters (between 74% and 78%) of Vietnamese women had ever been screened for cervical cancer (22, 25, 27).⁵ In contrast, a Washington survey of Vietnamese women, conducted during 2006 and 2007, found that 93% had received at least one Pap smear and 84% were adherent to interval screening guidelines.⁵

California Health Interview Survey data from 2003 provide the most recent direct comparisons of cervical cancer screening rates among Vietnamese and other racial/ethnic groups. This survey found that 70% of Vietnamese women ages ≥ 18 years reported Pap testing in the previous 3 years compared with 84% of White, 87% of Black, and 85% of Latina women. Pap testing levels among Filipino, Japanese, South Asian, Chinese, and Korean women were 86%, 75%, 73%, 68%, and 67%,

respectively (26). Overall, recent survey data indicate that Vietnamese women have lower levels of adherence to Pap testing guidelines than most other racial/ethnic groups. However, recent survey data also indicate that adherence levels among Vietnamese women vary by geographic area of the United States.

Factors Associated with Pap Testing Use

Older Vietnamese women have generally been shown to have lower levels of Pap testing participation than the female Vietnamese population in general (20-22, 25, 27). For example, Nguyen et al. reported that women ages ≥ 65 years were significantly less likely to have ever been screened for cervical cancer than younger women ($P < 0.01$; ref. 22). Marital status has consistently been shown to be associated with Pap smear receipt among Vietnamese Americans. Specifically, never married women have lower levels of screening participation than currently/previously married women (18-22, 25, 27, 28). Multiple studies have examined the associations between education, income, and Pap testing use among Vietnamese women with inconclusive findings (18-22, 25, 27).

To assess the effect of acculturation, researchers have considered the relationships between length of U.S. residence (among immigrants), English language proficiency, and cervical cancer screening. A majority (but not all) of these studies found that higher Pap testing levels are associated with longer U.S. residence (18-23, 25, 27). The relationship between English language proficiency and Pap smear receipt remains unclear (18-23, 25, 27, 29). Table 4 shows the associations between demographic characteristics and recent Pap testing use among Vietnamese women ages 20 to 69 years who were surveyed in metropolitan Seattle during late 2006 and early 2007.⁵

Some have speculated that traditional health beliefs may be barriers to preventive care for Southeast Asian and other immigrant groups (30, 31). However, several studies have failed to document an association between traditional Vietnamese beliefs and Pap testing (30, 32). For example, although Do et al. found that 71% of Vietnamese women believed that proper observance of the "sitting month" (a set of traditional postpartum practices that include the avoidance of wind and water) protects women from cervical cancer, this belief was not associated with recent Pap smear receipt (32).

Nearly all the studies that examined the role of having a usual source of care and/or usual doctor in Pap testing participation among Vietnamese women have documented associations (20, 22, 25, 27, 33). As would be expected, Kagawa-Singer et al. found that Vietnamese women who reported at least one doctor's visit during

⁵ Unpublished data.

Cervical Cancer Control in Vietnamese American Communities

the prior year were more likely to also report a recent Pap smear than those who did not (65% versus 43%; $P < 0.01$; ref. 28). Positive correlations have been found between having a female physician, having a non-Vietnamese physician, and cervical cancer screening (20-22, 25, 27). Additionally, one study found that women who received care at a community/county hospital clinic or a multi-specialty clinic were more likely to report a recent Pap smear than those who received care at a private physician's office (25). Results from surveys that looked at health insurance coverage in relation to Vietnamese women's cervical cancer screening behavior are inconsistent (19-22, 25, 27).

Two recent studies have used the Pathways Model (which originated in the PRECEDE-PROCEED planning framework) and multivariable methods to systematically examine relationships health-care system access and attitudes and the Pap testing practices of Vietnamese women (22, 27). In the first analysis (using 2000 data), having a female doctor (odds ratio, 1.9; 95% confidence interval, 1.2-2.9), having a respectful doctor (odds ratio, 2.0; 95% confidence interval, 1.1-3.6), having a physician recommend testing (odds ratio, 8.0; 95% confidence interval, 5.7-11.9), and having requested the test (odds ratio, 8.7; 95% confidence interval, 5.8-13.0) were associated with receipt of at least one Pap smear (22). The second analysis (using 2004 data) showed that, in addition to factors identified by the earlier analysis, the following factors were associated with previous Pap testing: having health insurance, having a usual place for health care, and having a Vietnamese male physician (negative association; ref. 27).

Another study used the theoretical perspective of the Health Behavior Framework to examine individual factors associated with recent Pap testing and found strong correlations ($P < 0.001$) between the following variables and recent Pap smear receipt: believing regular Pap tests decrease the risk of cancer and Pap testing is necessary for asymptomatic, sexually inactive, and postmenopausal women; reporting concern about pain/discomfort as a barrier to Pap testing; family members and friends had suggested Pap testing; and doctors had recommended Pap testing and had asked doctors for Pap testing. In a logistic regression model, believing Pap smears are necessary for asymptomatic women, doctors had recommended Pap testing, and had asked doctors for Pap testing were significantly associated with adherence to interval screening guidelines (33).

To summarize, levels of Pap testing use among Vietnamese American women have consistently been shown to be associated with some demographic and acculturation variables but not others. Health-care and physician factors are important determinants of cervical cancer screening participation. There is some evidence that beliefs about Pap testing are associated with adherence to cervical cancer screening guidelines.

Intervention Studies

Because many Vietnamese Americans are relatively new immigrants with limited English speaking proficiency and access to culturally appropriate health care, community-based interventions may be the best way to educate women and encourage cervical cancer screening participation (17). Further, the Cochrane Collaboration recently concluded that lay health worker outreach is a promising approach to improving health outcomes among racial/ethnic minority populations (34). As shown in Table 5, the Vietnamese Community Health Promotion Project in San Francisco has evaluated the effectiveness of lay health worker interventions as well as Vietnamese language media campaigns in increasing Pap testing use among Vietnamese women (27, 35-40).

In one study, indigenous lay health workers conducted a series of small group educational sessions with Vietnamese women in San Francisco, whereas Sacramento served as a control community. Pre- and post-intervention surveys showed that the proportion of women reporting at least one Pap smear increased significantly in the experimental area (46% pre-intervention versus 66% post-intervention; $P < 0.001$) but did not increase in the control area (40% pre-intervention versus 42% post-intervention; $P > 0.05$; ref. 36).

Another research project evaluated a Vietnamese language media campaign that included use of television, newspaper, and billboard advertising as well as the distribution of audiovisual and print educational materials. Post-intervention, no differences in recent Pap testing use existed between women in two northern California experimental counties (Alameda and Santa Clara) and two southern California control counties (Los Angeles and Orange). However, women in the experimental area were significantly more likely to be planning future Pap testing than women in the control area (37).

Table 2. Age-adjusted invasive cervical cancer incidence rates per 100,000 Women

Author, publication year	Group	Geographic area	Period	Rate in Southeast Asian/Vietnamese women	Rate in non-Latina White women
Taylor, 1996 (12)	Southeast Asian*	Surveillance, Epidemiology and End Results regions	1976-1984	39.4	8.2
Perkins, 1995 (13)	Southeast Asian*	California	1988-1992	35.2	7.5
Miller, 1996 (14)	Vietnamese	Surveillance, Epidemiology and End Results regions	1988-1992	43.0	7.5
Gomez, 2005 (15)	Vietnamese	Greater San Francisco Bay	1990-1993	49.0	—
			1994-1997	26.3	—
			1998-2000	15.2	—
McCracken, 2007 (6)	Vietnamese	California	2000-2002	14.0	7.3

* Cambodian, Hmong, Laotian, and Vietnamese.

† Los Angeles, San Francisco/Oakland, San Jose/Monterey, and Seattle/Puget Sound.

Table 3. Cervical cancer screening rates among Vietnamese Women ages ≥ 18 years

Author, publication year	Survey year(s)	Geographic area	Survey method	Cooperation rate* (%)	Sample size	Ever screened (%)	Screened last year [†] (%)	Screened last 3 y [†] (%)
Nguyen, 2002 (22)	2000	Harris County, Texas	Telephone	54	768	74 [‡]	66 [‡]	—
Nguyen, 2002 (22)	2000	Santa Clara County, California	Telephone	63	798	78 [‡]	60 [‡]	—
Kandula, 2006 (23) [§]	2001	California	Telephone	64	425	—	—	62 [‡]
Centers for Disease Control, 2004 (24)	2001-2002	Los Angeles, Orange, and Santa Clara Counties, California	Telephone	72	1,667	—	—	66 [‡]
Taylor, 2004 (25) [¶]	2002 [§]	King County, Washington	In-person	84	544	74	45	68
Holtby, 2006 (26) [§]	2003	California	Telephone	—	—	—	—	70
Nguyen, 2006 (27)	2004	Harris County, Texas	Telephone	50	1,005	71	53	—
Unpublished data ^{**}	2006-2007	King County, Washington	In-person	72	1,332	93	56	84
Unpublished data ^{††}	2007	Harris County, Texas	Telephone	74	765	76	49	66

*Response among reachable and eligible women (completed/completed and refused).

[†] National guidelines for interval Pap testing changed from every year to every 3 years in 2002.

[‡] Includes women without uteri.

[§] Study used California Health Interview Survey data.

^{||} Cooperation rate for all racial/ethnic groups.

[¶] Women ages 18 to 64 years.

^{**} Women ages 20 to 69 years.

^{††} Women ages ≥ 40 years.

The recently completed Vietnamese REACH for Health Initiative included a quasi-experimental study to evaluate a community-based cervical cancer control intervention as well as a randomized controlled trial of a Pap testing lay health worker intervention (27, 40). The community-based intervention was multifaceted and included capacity-building activities but primarily targeted women through a Vietnamese language media campaign (27, 38, 41). The intervention was implemented in Santa Clara County, California, whereas Harris County, Texas served as a control community. Intervention effect was measured through cross-sectional, pre-

intervention (2000) and post-intervention (2004) surveys in the experimental and control areas. The proportion of women reporting at least one Pap test increased in the experimental community (78-84%; $P \leq 0.001$) but not in the control community (74-71%; $P > 0.05$; ref. 27).

In the trial component of the Vietnamese REACH for Health Initiative, 1,005 women in Santa Clara County were randomized to lay health worker group education plus media-based education (combined intervention) or media-based education alone (media-only intervention). Women provided information about their Pap testing history 4 months after randomization. The combined

Table 4. Demographic characteristics associated with recent Pap testing among Vietnamese Women-King County, Washington, 2006 to 2007

Variable	n (%)	Screened last 3 y (%)	Odds ratio (95% confidence interval)*
Age (y)			
60-69	210 (16)	75 [†]	0.4 (0.2-0.9)
50-59	368 (28)	85	0.9 (0.5-1.8)
40-49	289 (22)	87	1.1 (0.6-2.1)
30-39	352 (27)	88	1.4 (0.7-2.5)
20-29	108 (8)	74	Reference
Marital status			
Currently married	1,077 (81)	86 [†]	3.6 (2.3-5.8)
Previously married	124 (9)	81	3.3 (1.7-6.4)
Never married	125 (9)	63	Reference
Years of education			
>12	372 (28)	85	1.2 (0.8-2.0)
12	321 (24)	84	1.1 (0.7-1.6)
<12	634 (48)	82	Reference
Household income (dollars)			
$\geq 30,000$	506 (38)	90 [†]	2.0 (1.3-3.2)
<30,000	603 (45)	82	1.4 (0.9-2.1)
Unknown	219 (16)	75	Reference
Years in U.S.			
≥ 20 [†]	219 (16)	84	1.6 (0.9-2.8)
10-19	813 (61)	85	1.6 (1.1-2.4)
<10	297 (22)	80	Reference
English proficiency			
Speaks very well/fluent	175 (13)	82	0.6 (0.3-1.3)
Speaks quite well (so-so)	660 (50)	85	0.9 (0.6-1.3)
Does not speak well/at all	492 (37)	82	Reference

*Adjusted for all other demographic variables.

[†] $P < 0.001$.

[‡] Includes eight U.S.-born women.

Table 5. Cervical cancer control intervention studies

Author, publication year	Study design	Assignment unit	Intervention component(s)	Evaluation method	Main finding(s)
Bird, 1998 (36)	Quasi-experimental	Community	Group educational sessions delivered by lay health workers Print educational materials Health fairs	Pre- and post-intervention Cross-sectional surveys	Effective in increasing Pap testing use
Jenkins, 1999 (37)	Quasi-experimental	Community	Vietnamese language media campaign Audiovisual and print educational materials	Pre- and post-intervention Cross-sectional surveys	No effect on Pap testing behavior Increased Pap testing intentions
Nguyen, 2006 (27)	Quasi-experimental	Community	Vietnamese language media campaign Print educational materials Community capacity-building activities*	Pre- and post-intervention Cross-sectional surveys	Effective in increasing Pap testing use
Mock, 2007 (40)	Randomized controlled trial [†]	Individual	Group educational sessions delivered by lay health workers	Survey four months after randomization	Effective in increasing Pap testing use

NOTE: All the studies included women ages ≥ 18 years.

*Community capacity-building activities included restoration of a government-funded low-cost screening program, a Pap testing clinic specifically for Vietnamese women, a Pap testing reminder system, and continuing medical education for Vietnamese physicians.

[†]Women in a community with an ongoing Pap testing Vietnamese language media campaign were randomized to a combined intervention group (lay health worker plus media education) or a media education alone group.

intervention was more effective than the media-only intervention in increasing the rate of previous Pap testing receipt (66-82% versus 70-76%; $P < 0.001$). Among those who had never been screened, significantly more women in the combined intervention group (46%) than in the media-only group (27%) obtained Pap tests ($P < 0.001$; ref. 40).

We were only able to identify four studies that evaluated cervical cancer control intervention programs for Vietnamese women in the United States, and all but one of these studies used a quasi-experimental design. However, our review of the limited intervention literature suggests that Vietnamese language media campaigns and lay health worker intervention programs have the potential to be effective in increasing Pap smear use in Vietnamese American communities.

Limitations of Available Data

Swallen et al. examined the reliability of Vietnamese racial classification in population-based cancer registry data. Specifically, persons with cancer diagnosed in Northern California during 1989 to 1992 and whom the registry considered Vietnamese were interviewed and asked to specify their race/ethnicity. The study findings suggested that 20% of cancer cases classified as Vietnamese were probably not Vietnamese. It is possible that the observed changes in cervical cancer incidence rates among Vietnamese women over time are, at least partly, a result of reductions in racial classification errors (42).

Evidence exists that the quality of survey data may differ by race/ethnicity. For example, the Pathways Project found that test-retest reliabilities for the question "have you ever had a mammogram" were significantly lower among Chinese, Vietnamese, Latina, and Black women than among White women (43). As another example, the Pathfinders Project examined medical records to validate Pap smear self-reports among survey respondents in multiethnic Alameda County, California. The proportions of Pap smear self-reports that could be validated among White, Black, Latina, Chinese, and Filipina women were 85%, 66%, 66%, 68%, and 67%, respectively (44). Therefore, data from surveys of Vietnamese women may overestimate cervical cancer screening use. Also, studies evaluating intervention programs should, ideally, verify women's Pap testing self-reports with provider reports.

Some of the previous surveys of Vietnamese women have had relatively low response rates, and all have been cross-sectional in nature (Table 3). Survey responders and nonresponders may have different levels of Pap smear participation. Indeed, the apparent increase in Pap testing levels between 2002 and 2006 to 2007 in Washington may be partly a reflection of the lower response rate in 2006 to 2007. It is also unclear whether higher levels of knowledge about cervical cancer and Pap testing lead to higher rates of cervical cancer screening participation or whether interactions with health-care providers during Pap smear appointments result in increased knowledge.

As reported in this article, researchers from the Vietnamese Community Health Promotion Project have successfully used quasi-experimental approaches to evaluate several cervical cancer control programs for

Vietnamese women. Specifically, they have documented increases in Pap testing rates among women in experimental communities with no change among women in control communities (together with an absence of any other promotional activities in the experimental and control communities; refs. 27, 37). However, other investigators have reported difficulties evaluating Pap testing interventions, using a two-community design, because of unanticipated promotional activities in their experimental and/or control communities (45, 46). Further, Murray et al. recently specified that group-randomized trials should be considered the gold standard for studies designed to evaluate cancer control interventions that operate at a group level, manipulate the social or physical environment, or cannot be delivered to individuals. They also specified that community intervention trials should be adequately powered with a sufficient number of randomized communities (47).

Discussion

An analysis of 2001 California Health Interview Survey data indicated that disparities in Pap smear use among Black, White, and Latina women no longer exist in California (48). Recent surveys suggest that disparities in Pap smear use between Vietnamese and other racial/ethnic groups have decreased over time in Washington but not in Texas. Over the last 5 years, the Vietnamese community in King County, Washington has been the focus of targeted cervical cancer control efforts by the National Breast and Cervical Cancer Early Detection Program as well as a community clinic system serving Asian Americans with limited English proficiency. The relatively high Pap testing levels among Vietnamese women who responded to the 2006 to 2007 King County survey may reflect the success of these efforts.

Our review suggests that (as in other disadvantaged and immigrant populations) the cervical cancer disparity among Vietnamese women is likely a marker for health-care access inequities (31, 49, 50). Efforts to increase screening participation in Vietnamese communities should enable women who are without a usual source of care to find a primary care doctor. Additionally, intervention programs should improve patient-provider communication by encouraging health-care providers (especially Vietnamese physicians serving women living in ethnic enclaves) to recommend Pap testing as well as by empowering women to ask for testing.

Although previous community intervention studies suggest that community-based approaches to cervical cancer control among Vietnamese American women hold promise, findings from these studies are inconclusive because the study samples only included two communities. Because most Vietnamese Americans live in a few geographic areas of the United States, there are a limited number of Vietnamese communities available for randomization into group-randomized trials (3). However, there is a compelling need for well-designed, adequately powered community intervention trials to decrease the cancer burden experienced by Vietnamese immigrants.

During 1998 to 2002, the cervical cancer incidence rate among Cambodian Americans in California and the Puget Sound area of Washington State was 15.0 per

100,000 women compared with 7.7 per 100,000 among non-Latina White women (51). Yang et al. examined cervical cancer among Hmong women in California. Their analysis showed that Hmong women experience high rates of invasive cervical cancer, and the incidence has not decreased over time. Specifically, the age-adjusted incidence rates during 1988 to 1991 and 1996 to 2001 were 37.5 and 33.7 per 100,000, respectively (52). Although these other Southeast Asian groups come from the same geographic area and have similar recent immigration histories to the Vietnamese, there are many cultural differences (1, 53). With ~600,000 individuals of Cambodian, Hmong, and Laotian descent in the United States, it is important that future efforts to increase Pap testing focus on these communities (3).

The U.S. Food and Drug Administration recently approved the quadrivalent HPV vaccine (HPV-16, HPV-18, HPV-6, and HPV-11) for children, adolescents, and young women ages 9 to 26 years (54). Future cervical cancer control intervention programs for Vietnamese and other racial/ethnic groups should promote use of the HPV vaccine for age-eligible individuals as well regular interval Pap testing for adult women who may have already been infected with HPV.

Conclusion

In conclusion, there is a paucity of information about effective cervical cancer control interventions for Vietnamese women in the United States. Survey data indicate that efforts to increase Pap testing receipt among Vietnamese women must address both cognitive and contextual influences. Future community intervention studies should use adequately powered, group-randomized designs to further evaluate Vietnamese language media campaigns and lay health worker programs as well as interventions targeting the health-care providers who serve Vietnamese women. When possible, such intervention programs should be evaluated using medical records verification of Pap testing self-reports. In the future, dissemination studies will be needed to examine effective methods of disseminating evidence-based cervical cancer control interventions to Vietnamese communities throughout the United States (55).

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

References

1. Chan S, Lee E. Families with Asian roots. In: Developing cross-cultural competence. In Lynch EW, editors. Hanson (MJ). Baltimore: Paul Brookes Publishing; 2004.
2. U.S. Census Bureau. The American community-Asians: 2004. Washington (DC): U.S. Department of Commerce; 2007.
3. Niedzweicki M, Tuong TC. Southeast Asian American statistical profile. Washington (DC): Southeast Asia Resource Action Center; 2004.

Cervical Cancer Control in Vietnamese American Communities

4. U.S. Census Bureau. We the people: Asians in the United States. Washington (DC): U.S. Department of Commerce; 2004.
5. President's Advisory Commission on Asian Americans and Pacific Islanders. Asian Americans and Pacific Islanders—addressing health disparities: opportunities for building a healthier America. Washington (DC): Department of Health and Human Services; 2003.
6. McCracken M, Olsen M, Chen MS, et al. Cancer incidence, mortality, and associated risk factors among Asian Americans of Chinese, Filipino, Vietnamese, Korean, and Japanese ethnicities. *CA Cancer J Clin* 2007;57:190–205.
7. Parkin DM. The global health burden of infection-associated cancers in the year 2000. *Int J Cancer* 2006;118:3030–44.
8. Pham TH, Nguyen TH, Herrero R, et al. Human papillomavirus infection among women in South and North Vietnam. *Int J Cancer* 2003;104:213–20.
9. Vo PD, Nguyen TT, Nguyen P, et al. Human papillomavirus and abnormal Pap test results in Vietnamese American women: a pilot case-control study. *J Lower Genital Tract Dis* 2004;8:217–23.
10. Saslow D, Runowicz CD, Solomon D, et al. American Cancer Society guideline for the early detection of cervical neoplasia and cancer. *CA Cancer J Clin* 2002;52:342–62.
11. U.S. Department of Health and Human Services. Healthy People 2010—understanding and improving health. Washington (DC): U.S. Government Printing Office; 2000.
12. Taylor VM, Jackson JC, Schwartz SM, Tu S-P, Thompson B. Cervical cancer among Asian American women: a neglected public health problem? *Asian Am Pac Isl J Health* 1996;4:327–42.
13. Perkins CI, Morris CR, Wright WE, Young JL. Cancer incidence and mortality in California by detailed race/ethnicity, 1988-1992. Sacramento: California Department of Health Services; 1995.
14. Miller BA, Kolonel LN, Bernstein L, et al. Racial/ethnic patterns of cancer in the United States: 1988-1992. Bethesda: National Cancer Institute; 1996.
15. Gomez SL, Le GM, Miller T, et al. Cancer incidence among Asians in the Greater Bay Area: 1990-2002. Fremont: Northern California Cancer Center; 2005.
16. Cockburn M, Deapen D. Cancer incidence and mortality in California—trends by ethnicity: 1988-2001. Los Angeles: Norris Comprehensive Cancer Center; 2004.
17. McPhee SJ, Nguyen TT. Cancer, cancer risk factors, and community-based cancer control trials in Vietnamese Americans. *Asian Am Pac Isl J Health* 2000;8:19–31.
18. Centers for Disease Control. Behavioral risk Factor survey of Vietnamese-California, 1991. *Morb Mortal Wkly Rep* 1992;41:69–72.
19. Yi JK. Factors associated with cervical cancer screening behavior among Vietnamese women. *J Community Health* 1994;19:189–200.
20. McPhee SJ, Bird JA, Davis T, Ha NT, Jenkins CNH, Le B. Barriers to breast and cervical cancer screening among Vietnamese American women. *Am J Prev Med* 1997;13:205–13.
21. McPhee SJ, Stewart S, Brock KC, Bird JA, Jenkins CNH, Pham GQ. Factors associated with breast and cervical cancer screening practices among Vietnamese American women. *Cancer Detect Prev* 1997;21:510–21.
22. Nguyen TT, McPhee SJ, Nguyen T, Lam T, Mock J. Predictors of cervical cancer awareness, intention, and receipt among Vietnamese American women. *Am J Prev Med* 2002;23:207–14.
23. Kandula NR, Wen M, Jacobs EA, Lauderdale DS. Low rates of colorectal, cervical, and breast cancer screening in Asian Americans compared with non-Hispanic Whites: cultural influences or access to care. *Cancer* 2006;107:184–92.
24. Centers for Disease Control. Health status of Cambodians and Vietnamese, United States, 2001-2002. *Morb Mortal Wkly Rep* 2004;53:760–5.
25. Taylor VM, Schwartz SM, Yasui Y, et al. Pap testing among Vietnamese women: the health care system and physician factors. *J Community Health* 2004;29:437–50.
26. Holtby S, Zahnd E, Lordi N, McCain C, Chia YJ, Kurata J. Health of California's adults, adolescents, and children. Sacramento: California Department of Health Services; 2006.
27. Nguyen TT, McPhee SJ, Gildengorin G, et al. Papanicolaou testing among Vietnamese Americans: results of a multifaceted intervention. *Am J Prev Med* 2006;31:1–9.
28. Kagawa-Singer M, Pourat N, Breen N, et al. Breast and cervical cancer screening rates of subgroups of Asian American women in California. *Med Care Res Rev* 2007. Epub ahead of print 2007 Sep 5.
29. Ponce N, Gatchell M, Brown R. Cancer screening rates among Asian ethnic groups. UCLA Center for Health Policy Research health policy fact sheet; 2003 Nov.
30. Jenkins CNH, Le T, McPhee SJ, Stewart S, Ha NT. Health care access and preventive care among Vietnamese immigrants. *Soc Sci Med* 1996;43:1049–56.
31. Schleider E. Immigrant women and cervical cancer prevention in the United States. Baltimore: John Hopkins Bloomberg School of Health; 2007.
32. Do HH, Taylor VM, Burke N, Yasui Y, Schwartz SM, Jackson JC. Knowledge about cervical cancer risk factors, traditional health beliefs, and Pap testing among Vietnamese American women. *J Immigrant Minority Health* 2007;9:109–14.
33. Taylor VM, Yasui Y, Burke N, et al. Pap testing adherence among Vietnamese American women. *Cancer Epidemiol Biomarkers Prev* 2004;13:613–9.
34. Lewan SA, Dick J, Pond P, et al. Lay health workers in primary and community care. <http://www.cochrane.org>. Accessed 2007.
35. McPhee SJ, Bird JA, Ha NT, Jenkins CNH, Fordham D, Le B. Pathways to early cancer detection for Vietnamese women. *Health Educ Q (Suppl)* 1996;23:66–75.
36. Bird JA, McPhee SJ, Ha NT, Le B, Davis T, Jenkins CNH. Opening pathways to cancer screening for Vietnamese-American women: lay health workers hold a key. *Prev Med* 1998;27:821–9.
37. Jenkins CNH, McPhee SJ, Bird JA, et al. Effect of a media-led education campaign on breast and cervical cancer screening among Vietnamese-American women. *Prev Med* 1999;28:395–406.
38. Nguyen TT, McPhee SJ, Bui-Tong N, et al. Community-based participatory research increases cervical cancer screening among Vietnamese Americans. *J Health Care Poor Underserved* 2006;17:31–54.
39. Lam TK, McPhee SJ, Mock J, et al. Encouraging Vietnamese American women to obtain Pap tests through lay health worker outreach and media education. *J Gen Intern Med* 2003;18:516–24.
40. Mock J, McPhee SJ, Nguyen T, et al. Effective lay health worker outreach and media-based education for promoting cervical cancer screening among Vietnamese American women. *Am J Public Health* 2007;97:1693–700.
41. Lai KQ, Nguyen TT, Mock J, McPhee SJ, Doan HT, Pham TH. Increasing Vietnamese American physicians' knowledge of cervical cancer and Pap testing: impact of continuing medical education programs. *Ethn Dis* 2004;14:124–8.
42. Swallen KC, Glaser SL, Stewart SL, Jenkins CN, McPhee SJ. Accuracy of racial classification of Vietnamese patients in a population-based cancer registry. *Ethn Dis* 1998;8:218–27.
43. Pasick RJ, Stewart SL, Bird JA, D'Onofrio CN. Quality of data in multi-ethnic health surveys. *Public Health Rep (Suppl)* 2001;116:223–43.
44. McPhee SJ, Nguyen TT, Shema SJ, et al. Validation of recall of breast and cervical cancer screening by women in an ethnically diverse population. *Prev Med* 2002;35:463–73.
45. Suarez L, Roche RA, Pulley LV, Weiss NS, Goldman D, Simpson DM. Why a peer intervention program for Mexican American women failed to modify the secular trend in cancer screening. *Am J Prev Med* 1997;13:411–7.
46. Wismer BA, Moskowitz JM, Min K, et al. Interim assessment of a community intervention to improve breast and cervical cancer screening among Korean American women. *J Public Health Manage Pract* 2001;7:61–70.
47. Murray DM, Pals SL, Blitstein JL, Alfano CM, Lehman J. Design and analysis of group-randomized trials in cancer: a review of current practices. *J Natl Cancer Inst* 2008;100:483–91.
48. De Alba I, Ngo-Metzger Q, Sweningson JM, Hubbell FA. Pap smear use in California: are we closing the racial/ethnic gap? *Prev Med* 2004;40:747–55.
49. Freeman HP, Wingrove PK. Excess cervical cancer mortality: a marker for low access to health care in poor communities. Rockville: National Cancer Institute; 2005.
50. Tanjasiri SP, Tran JH, Kagawa-Singer M, et al. Exploring access to cancer control services for Asian-American and Pacific islander communities in Southern California. *Ethn Dis* 2004;14:14–9.
51. Kem R, Chu KC. Cambodian cancer incidence rates in California and Washington, 1998-2002. *Cancer* 2007;110:1370–5.
52. Yang RC, Mills PK, Riordan DG. Cervical cancer among Hmong women in California, 1988 to 2000. *Am J Prev Med* 2004;27:132–8.
53. McGarvey EL, Clavet GJ, Johnson JB, Butler A, Cook KO, Pennini B. Cancer screening practices and attitudes: comparison of low-income women in three ethnic groups. *Ethn Health* 2003;8:71–82.
54. Markowitz LE, Dunne EF, Saraiya M, Lawson HW, Chesson H, Unger ER; Centers for Disease Control and Prevention (CDC); Advisory Committee on Immunization Practices (ACIP). Quadrivalent human papillomavirus vaccine: recommendations of the Advisory Committee on Immunization Practices (ACIP). *Morb Mortal Wkly Rep* 2007;56:1–24.
55. Kerner J, Rimer B, Emmons K. Dissemination research and research dissemination: how can we close the gap? *Health Psychol* 2005;24:443–6.

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Cervical Cancer Control Research in Vietnamese American Communities

Victoria M. Taylor, Tung T. Nguyen, J. Carey Jackson, et al.

Cancer Epidemiol Biomarkers Prev 2008;17:2924-2930.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/17/11/2924>

Cited articles This article cites 39 articles, 1 of which you can access for free at:
<http://cebp.aacrjournals.org/content/17/11/2924.full#ref-list-1>

Citing articles This article has been cited by 1 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/17/11/2924.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/17/11/2924>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.