

Metabolic Phenotype and Risk of Colorectal Cancer in Normal-Weight Postmenopausal Women

Xiaoyun Liang¹, Karen L. Margolis², Michael Hendryx³, Thomas E. Rohan⁴, Erik J. Groessl^{5,6}, Cynthia A. Thomson⁷, Candyce H. Kroenke⁸, Michael S. Simon⁹, Dorothy Lane¹⁰, Marcia Stefanick¹¹, and Juhua Luo¹²

Abstract

Background: The prevalence of metabolically unhealthy phenotype in normal-weight adults is 30%, and few studies have explored the association between metabolic phenotype and colorectal cancer incidence in normal-weight individuals. Our aim was to compare the risk of colorectal cancer in normal-weight postmenopausal women who were characterized by either the metabolically healthy phenotype or the metabolically unhealthy phenotype.

Methods: A large prospective cohort, the Women's Health Initiative, was used. The analytic sample included 5,068 postmenopausal women with BMI 18.5 to <25 kg/m². Metabolic phenotype was defined using the Adult Treatment Panel-III definition, excluding waist circumference; therefore, women with one or none of the four components (elevated triglycerides, low high-density lipoprotein cholesterol, elevated blood pressure,

and elevated fasting glucose) were classified as metabolically healthy. Multivariable Cox proportional hazards regression was used to estimate adjusted HRs for the association between metabolic phenotype and risk of colorectal cancer.

Results: Among normal-weight women, those who were metabolically unhealthy had higher risks of colorectal cancer (HR, 1.49; 95% CI, 1.02–2.18) compared with those who were metabolically healthy.

Conclusions: A metabolically unhealthy phenotype was associated with higher risk of colorectal cancer among normal-weight women.

Impact: Normal-weight women should still be evaluated for metabolic health and appropriate steps taken to reduce their risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev*; 26(2): 155–61. ©2017 AACR.

Introduction

According to estimates by the International Agency for Research on Cancer (Lyon, France), in 2012, colorectal cancer accounted for 9.2% of all incident cancers among women in the

world, which made it the second most common cancer in women (1). In 2016 in the United States, it is estimated that 63,670 women will be diagnosed with colorectal cancer and 23,170 will die from it (2).

The risk of developing colorectal cancer is associated with genetic, environmental, socioeconomic, and lifestyle factors. These include age, sex, race, and family history, as well as modifiable risk factors, including smoking, alcohol consumption, diet (intake of high percent calories from fat, low consumption of dietary fiber), physical inactivity, and obesity. Of particular interest for the current study, overweight and obesity are already well-established risk factors for colorectal cancer (3, 4). Protective factors include current or past regular use of NSAIDs and prior screening examinations (5, 6).

Obesity often cooccurs with metabolic disorders, but normal weight with a metabolically unhealthy phenotype has been recognized since the 1980s (7) and yet has been understudied. The definition of metabolic health varies among studies, but most researchers use the Adult Treatment Panel-III (ATP-III), which includes five components: elevated waist circumference, elevated triglyceride, low HDL-C (high-density lipoprotein cholesterol), elevated blood pressure, and elevated glucose (8). The metabolically unhealthy phenotype is defined by two or more of four of these components except for waist circumference, and the metabolic syndrome is identified by the presence of three or more of five components (9–11). A recent worldwide meta-analysis shows that 30.0%

¹School of Social Development and Public Policy, Beijing Normal University, Beijing, China. ²HealthPartners Institute, Minneapolis, Minnesota. ³Department of Applied Health Science, School of Public Health, Indiana University Bloomington, Bloomington, Indiana. ⁴Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, New York. ⁵Department of Family Medicine and Public Health, University of California San Diego, La Jolla, California. ⁶VA San Diego Healthcare System, San Diego, California. ⁷University of Arizona Cancer Center, Tucson, Arizona. ⁸Division of Research, Kaiser Permanente Northern California, Oakland, California. ⁹Department of Oncology, Karmanos Cancer Institute, Wayne State University, Detroit, Michigan. ¹⁰Department of Family, Population and Preventive Medicine, School of Medicine, Stony Brook University, Stony Brook, New York. ¹¹Department of Medicine, Stanford Prevention Research Center, Stanford University, Stanford, California. ¹²Department of Epidemiology and Biostatistics, School of Public Health, Indiana University Bloomington, Bloomington, Indiana.

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Xiaoyun Liang, Beijing Normal University, 19 Xijiekouwai Street, Beijing 100875, China. Phone: 8613-8118-79287; Fax: 8610-5880-0366; E-mail: liangxiaoyun@bnu.edu.cn

doi: 10.1158/1055-9965.EPI-16-0761

©2017 American Association for Cancer Research.

Liang et al.

[95% confidence interval (CI), 25.6–35.6%] of normal-weight adults were metabolically unhealthy (12). Using criteria of two or more metabolic abnormalities, Wildman and colleagues found that 23.5% of normal-weight U.S. adults were metabolically unhealthy (13).

Two meta-analyses of studies that evaluated the association of metabolic syndrome with colorectal cancer found a 30% to 40% higher risk in both men and women (3, 14). It is unclear whether the higher risk is independent of overweight and obesity, as waist circumference is a component of the metabolic syndrome and is highly correlated with body mass index (BMI). Few studies have assessed the association between metabolically defined unhealthy phenotype and risk of colorectal cancer, particularly in normal-weight people.

To build upon and contribute to the evidence relating metabolic phenotype to colorectal cancer risk, we used the rich Women's Health Initiative (WHI) dataset to compare the risk of colorectal cancer between normal-weight postmenopausal women who were characterized by either metabolically healthy phenotype or metabolically unhealthy phenotype.

Materials and Methods

Population

We used data from the WHI Observational Study (OS) and Clinical Trial (CT), a prospective cohort study enrolled through 40 clinical centers throughout the United States. A total of 161,808 postmenopausal women between 50 and 79 years of age were recruited from 1993 to 1998. Three CTs ($n = 68,132$) included hormone therapy, dietary modification, and calcium plus vitamin D supplementation. All of the OS and CT women completed screening and enrolment questionnaires by self-report, interview, physical examination, and fasting blood sample collection. Written informed consent and appropriate Institutional Review Board approval were obtained by each participating WHI site.

The WHI dataset includes metabolic biomarker data from several subsets of women. These data were collected prospectively on a 6% random sample of women in the CT ($N = 4,544$), and a 1% sample of women in the OS ($N = 1,062$) at baseline (15). Stored samples were used in WHI ancillary studies, including the SNP Health Association Resource cohort, and the European American Hormone Trial subcohort to provide additional




Figure 1.
Flow diagram of participants included in the analysis.

metabolic biomarker results that were applied in the current analysis. The SNP Health Association Resource cohort includes 12,007 OS and CT participants (8,405 African Americans and 3,602 Hispanics), and the European American Hormone Trial includes 10,306 participants. Blood samples collected at baseline and included in this analysis were analyzed for glucose, triglycerides, total cholesterol, and HDL-C. In addition, baseline height, weight, waist circumference, and systolic and diastolic blood pressure were measured by study staff using a standardized protocol at clinical visits.

In this study, women were excluded if they had any prior diagnosis of cancer other than nonmelanoma skin cancer before the date of the baseline questionnaire administration. Women with missing data on BMI ($n = 164$), triglycerides ($n = 8$), HDL cholesterol ($n = 1,599$), blood pressure ($n = 187$), or fasting glucose ($n = 39$) were excluded. As the exposure of interest for this analysis focused on normal-weight women with metabolic phenotype, those who were overweight, obese, or underweight (BMI ≥ 25 kg/m² or BMI < 18.5 kg/m²; $n = 16,612$) were excluded. Women with missing covariate data, that is, age, ethnicity, smoking, alcohol consumption, were also excluded. For those missing data on family history of colorectal cancer, we created an indicator variable and included it in the multivariable model. A flowchart showing derivation of the included study population is presented as Fig. 1.

Exposure measurements

We defined the metabolic phenotype using the ATP-III metabolic syndrome definition excluding the waist circumference ≥ 80 cm component due to its significant collinearity with BMI. Women were classified as metabolically healthy if they had less than two of the following four ATP-III components, and women with two or more of the four components were classified as metabolically unhealthy: (i) elevated triglycerides (≥ 150 mg/dL); (ii) low HDL-C (< 50 mg/dL or use of medication for reduced HDL-C); (iii) elevated blood pressure (systolic/diastolic blood pressure $\geq 130/85$ mm Hg or use of antihypertensive medication); and (iv) elevated fasting glucose (≥ 100 mg/dL or use of diabetes medication; refs. 8, 16).

Metabolic syndrome was defined as having 3 or more of 5 components (elevated waist circumference, elevated triglyceride, low HDL-C, elevated blood pressure, and elevated glucose). Seventeen additional women were excluded because of missing waist circumference.

HOMA-IR is a standard measure of insulin resistance and is defined by a formula that incorporates both insulin and glucose levels [(fasting insulin (mU/L) \times fasting glucose (mmol/L))/22.5; ref. 17]. We used the same method to define HOMA-IR as another WHI study (18), categorized HOMA-IR into quartiles (q1–q4), and used the 75th percentile value as the cut-off point to define metabolic phenotype: metabolically healthy (HOMA-IR–q1q2q3),

Table 1. Baseline characteristics of the WHI normal-weight participants by metabolic phenotype ($N = 5,068$)

	Metabolically healthy	Metabolically unhealthy	P
Total number of women	3,358	1,710	
Age at cohort entry (mean \pm SD, y)	63.8 \pm 7.7	66.7 \pm 6.9	<0.001
Race/ethnicity (%)			
Black or African American	23.8	20.5	0.014
Hispanic/Latino	15.9	15.4	
NonHispanic white	56.1	58.7	
Others	4.2	5.4	
Smoking status (%)			<0.001
Never smokers	52.5	52.8	
Former smokers	37.0	32.4	
Current smokers	10.5	14.8	
Alcohol intake (7+ drinks/wk, %)	41.1	33.9	<0.001
Total energy intake [(mean \pm SD), med serv/day]	1,515.0 \pm 678.4	1,472.3 \pm 670.7	0.033
Dietary fiber [(mean \pm SD), med serv/day]	15.3 \pm 7.2	14.9 \pm 7.3	0.031
Percent calories from fat [(mean \pm SD), med serv/day]	32.0 \pm 8.6	31.9 \pm 8.7	0.532
Physical activity (%)			<0.001
0–1.5 METs/wk	18.0	19.8	
>1.5–8 METs/wk	24.5	29.2	
>8–19 METs/wk	29.5	28.3	
>19 METs/wk	28.0	22.8	
Family history of colorectal cancer (%)			0.807
No	75.8	75.0	
Yes	14.1	14.7	
Missing	10.1	10.3	
NSAID use (%)	13.5	13.4	0.899
Systolic blood pressure (mean \pm SD, mm Hg)	122.2 \pm 17.1	134.6 \pm 18.2	<0.001
Diastolic blood pressure (mean \pm SD, mm Hg)	73.0 \pm 8.9	75.8 \pm 9.6	<0.001
Total cholesterol (mean \pm SD, mg/dL)	221.6 \pm 36.0	236.2 \pm 46.5	<0.001
HDL cholesterol (mean \pm SD, mg/dL)	66.0 \pm 13.9	52.4 \pm 14.2	<0.001
Triglycerides (mean \pm SD, mg/dL)	95.7 \pm 39.3	165.2 \pm 104.5	<0.001
Glucose (mean \pm SD, mg/dL)	88.9 \pm 12.2	102.9 \pm 32.1	<0.001
C-reactive protein (mean \pm SD, mg/mL)	2.3 \pm 4.3	3.7 \pm 11.0	<0.001
HOMA-IR (mean \pm SD, mmol/L ⁺ mU/L)	1.4 \pm 1.1	2.6 \pm 7.1	<0.001
Waist circumference (mean \pm SD, cm)	74.5 \pm 7.0	77.8 \pm 7.9	<0.001
Hypertension (%)	25.0	61.1	<0.001
Diabetes (%)	1.0	13.0	<0.001
Metabolic syndrome (%)	0	40.0	<0.001

Abbreviation: serv, servings.

Liang et al.

metabolically unhealthy (HOMA-IR-q4; ref. 19). A total of 182 additional women were excluded from the models because of missing insulin level.

Outcome measurements

The primary outcome of interest was incident colorectal cancer. In the WHI, incident colorectal cancer was documented and coded for primary site, anatomic subsite, diagnosis date, stage, tumor size, and grade. The diagnosis of colorectal cancer was ascertained through self-administered questionnaires, then confirmed by a centralized review of pathology reports from diagnostic aspirations, biopsies, surgeries, and discharge summaries, and subsequently adjudicated by trained, central adjudicators (20). For this study, the end of follow-up time was September 30, 2015.

Laboratory methods

Fasting blood samples collected at baseline were maintained at 4°C for up to one hour until plasma or serum was separated from cells. Plasma/serum aliquots were then frozen at -70°C and sent on dry ice to the central repository (Rockville, MD), where they were stored at -80°C. Glucose was measured using the hexokinase method on a Hitachi 747. Monthly interassay coefficients of variation were less than 2%. Total cholesterol and triglycerides were analyzed by enzymatic methods on a Hitachi 747 analyzer (Boehringer Mannheim Diagnostics). HDL-C was isolated using heparin manganese chloride. Coefficients of variation for total cholesterol, triglycerides, and HDL-C were all 2% or less. Serum insulin was measured in a step-wise sandwich ELISA procedure on an ES 300 (Boehringer Mannheim Diagnostics). An ongoing monthly quality assurance program was maintained with the Diabetes Diagnostic Laboratory at the University of Missouri (Columbia, MO; ref. 21).

Anthropometric measures and blood pressure

Participants were asked to remove their shoes for anthropomorphic measures. Height (cm) was measured using a wall-mounted stadiometer to the nearest 0.1 cm. Weight (kg) was measured using a balance beam scale to the nearest 0.1 kg. Waist circumference at the natural waist or narrowest part of the torso was measured to the nearest 0.1 cm. Blood pressure was measured twice in the right arm with the participant in the seated position and rested for 5 minutes using a conventional mercury sphygmomanometer and appropriately sized cuffs. Two blood pressure measurements were obtained at least 30 seconds apart, and the average of the two measurements was used in the analysis (21).

Statistical analysis

Characteristics of women with the metabolically healthy phenotype were compared with those of women with the

metabolically unhealthy phenotype. Means (\pm SDs) were used to summarize continuous characteristics at baseline, whereas proportions were used for categorical variables. *t* tests were used to compare continuous variables, and χ^2 tests were used to compare the distributions of categorical variables.

Survival time was measured as the date from enrolment to colorectal or colon cancer diagnosis, death, or end of follow-up, whichever came first. Multivariable Cox proportional hazards regression was used to estimate adjusted HRs for colorectal and colon cancer incidence. Each model was adjusted for age, and we also ran additional models that included age, race/ethnicity, smoking, alcohol consumption, physical activity, total energy intake, dietary fiber, percent calories from fat, family history of colorectal cancer, NSAIDs use, and treatment arm in each CT. Proportional hazards assumptions were tested on the basis of the Schoenfeld residuals, and no violation was observed.

We also evaluated the associations between each of the four components of metabolic phenotype (elevated triglycerides, low HDL-C, elevated blood pressure, and elevated fasting glucose) and cancer incidence.

Sensitivity analyses were carried out in all models, including (i) metabolic unhealthy phenotype was replaced with metabolic syndrome, (ii) using HOMA-IR to define metabolic phenotype, (iii) excluding women with diabetes at baseline enrollment, and (iv) limiting analysis to nonusers of NSAIDs.

Results

Of 5,068 normal-weight women in the analysis, women with metabolically healthy and metabolically unhealthy phenotypes accounted for 66.3% and 33.7%, respectively (Table 1). There were significant differences between the two groups with respect to age, race/ethnicity, smoking status, alcohol intake, total energy intake, and physical activity. Metabolically unhealthy women were older, were more likely to be nonHispanic white, were more likely to currently smoke, had lower alcohol consumption, total energy intake and dietary fiber, and reported less physical activity. There were 114 cases of colorectal cancer that occurred during mean follow-up time of 14.3 years, and average follow-up time from study entry to diagnosis of colorectal cancer was 5.3 years (0.2–18.0 years).

In both age-adjusted and fully adjusted models, compared with women with metabolically healthy normal weight, metabolically unhealthy normal weight was associated with a higher risk of colorectal cancer. The HR for colorectal cancer in the fully adjusted model was 1.49 (95% CI, 1.02–2.18; Table 2).

Among the individual components of metabolic syndrome, elevated fasting glucose was associated with a higher risk of colorectal and colon cancer (HR, 1.71; 95% CI, 1.12–2.58; HR,

Table 2. HRs and 95% CIs for the association between metabolic phenotype and colorectal/colon cancer among WHI normal-weight women

	Cases	Age-adjusted HR (95% CI) ^a	Fully adjusted HR (95% CI) ^b
Colorectal cancer			
Metabolically healthy	64	1	1
Metabolically unhealthy	50	1.51 (1.03–2.19)	1.49 (1.02–2.18)
Colon cancer			
Metabolically healthy	50	1	1
Metabolically unhealthy	38	1.48 (0.96–2.27)	1.51 (0.98–2.33)

^aHR and 95% CI adjusted for age.

^bHR and 95% CI adjusted for age, ethnicity, smoking, alcohol consumption, physical activity, total energy intake, dietary fiber, percent calories from fat, family history of colorectal cancer, NSAIDs use, and treatment arm in each CT.

Table 3. HRs and 95% CIs for the associations between individual components of metabolic phenotype and colorectal/colon cancer among normal-weight women

	Colorectal cancer		Colon cancer	
	Cases	HR (95% CI)	Cases	HR (95% CI)
Elevated triglycerides ^a				
No	76	1	55	1
Yes	38	0.99 (0.65–1.52)	33	1.32 (0.82–2.12)
Low HDL-C ^b				
No	82	1	65	1
Yes	32	1.33 (0.84–2.09)	23	1.13 (0.67–1.91)
Elevated blood pressure ^c				
No	51	1	40	1
Yes	63	1.09 (0.74–1.60)	48	1.04 (0.67–1.62)
Elevated fasting glucose ^d				
No	81	1	63	1
Yes	33	1.70 (1.12–2.58)	25	1.68 (1.04–2.71)

NOTE: HR and 95% CI adjusted for age, ethnicity, smoking, alcohol consumption, physical activity, total energy intake, dietary fiber, percent calories from fat, family history of colorectal cancer, NSAIDs use, and treatment arm in each CT.

^aElevated triglycerides: ≥ 150 mg/dL.

^bLow HDL-C: < 50 mg/dL or use of medication for reduced HDL-C.

^cElevated blood pressure: systolic/diastolic blood pressure $\geq 130/85$ mm Hg or use of antihypertensive medication.

^dElevated fasting glucose: ≥ 100 mg/dL or use of diabetes medication.

1.68; 95% CI, 1.04–2.71, respectively). Elevated triglycerides, low HDL-C, and elevated blood pressure were not associated with colorectal or colon cancer (Table 3).

A sensitivity analysis showed that when metabolic syndrome was used instead of metabolic phenotype, women with metabolic syndrome had a higher risk for both colorectal and colon cancer than those in the main results (HR, 2.14; 95% CI, 1.38–3.32; HR, 2.42; 95% CI, 1.48–3.95, respectively; Supplementary Table S1). When HOMA-IR was used to define metabolic phenotype, the results were similar to the main results. In addition, when women with diabetes ($n = 205$) were excluded from the analysis, the results were similar to the main results (Supplementary Table S2). Among nonusers of NSAIDs, metabolically unhealthy women had a higher risk for both colorectal and colon cancer than those in the main results (HR, 1.56; 95% CI, 1.04–2.35; HR, 1.61; 95% CI, 1.01–2.56, respectively; Supplementary Table S3).

Discussion

The results of this long-term prospective study of normal-weight postmenopausal women suggest that metabolically unhealthy women have a higher risk of colorectal cancer than metabolically healthy women.

Using WHI data, Gunter and colleagues found that, compared with metabolically healthy normal-weight women, metabolically unhealthy normal-weight women were at higher risk of breast cancer (18). We observed a similar positive association for the metabolically unhealthy normal-weight phenotype and colorectal cancer. With younger participants (around 57 years old on average), a recently published nested case-control study in Europe by Murphy and colleagues observed that, compared with metabolically healthy normal-weight individuals, a higher colorectal cancer risk was observed among metabolically unhealthy normal-weight participants (OR, 1.59; 95% CI, 1.10–2.28; ref. 22), findings consistent with our study (HR, 1.49; 95% CI, 1.02–2.18), despite the fact that they classified individuals as metabolically healthy or unhealthy based on the C-peptide level, which is not a generally accepted clinical definition. In addition, use of aspirin and other NSAIDs that were associated with reduced risk of colorectal cancer were not included in the Murphy and colleagues' study.

Colorectal cancer includes colon cancer, rectosigmoid cancer, and rectal cancer. In our study, the association between metabolic phenotype and risk of colon cancer was not significant. These results are similar with those of Murphy and colleagues who did not find a statistically significant association between metabolic unhealthy phenotype and risk of colon cancer among normal-weight individuals (OR, 1.49; 95% CI, 0.92–2.43; ref. 22).

Kabat and colleagues used WHI data to identify 81 incident cases of colorectal cancer among 4,862 eligible women with median follow-up of 12 years and found that among the individual components of metabolic syndrome, only elevated fasting glucose was associated with higher risk of colorectal and colon cancer (23), which is consistent with our findings. In other studies, elevated fasting glucose or diabetes has been repeatedly shown to be associated with the risk of colorectal and colon cancer (24, 25).

The reason why colorectal cancer risk is higher in normal-weight women with metabolic abnormalities is not entirely clear, but mechanistically, these results may suggest that the phenotype represents a proinflammatory state. In fact, anti-inflammatory medications, such as aspirin, are among the strongest preventive agents for primary prevention of colorectal cancer (4). The current study shows that the association between metabolic phenotype and colorectal cancer risk was strengthened among nonusers of NSAIDs, which indicates that there is some residual confounding after the basic adjustment for NSAID use. In addition, proinflammatory and anti-inflammatory cytokine concentrations have been associated with colorectal cancer risk (26, 27). Furthermore, compared with metabolically healthy normal-weight individuals, metabolically unhealthy normal-weight individuals may have less physical activity (28) and consume more saturated fat and less fiber (27); however, physical activity, saturated fat, and dietary fiber were included as covariates in this study, and it is unlikely that the differences in these lifestyle factors would account for the observed association between metabolic abnormality and risk of colorectal cancer.

These results indicate that among normal-weight women, a metabolically unhealthy phenotype is a relevant risk factor for colorectal cancer. Current guidelines recommend commencing colorectal cancer screening based primarily on age. Earlier identification of individuals at higher risk based on obesity or high-risk metabolic phenotype could result not only in appropriate

Liang et al.

preventive interventions, but also earlier screening thus increased likelihood of early-stage diagnosis and improved survival. Consideration of this possibility would require analyses of the potential benefits of alternative screening recommendations relative to costs.

The strengths of this study lie in the prospective long-term follow-up among several subcohorts of the WHI. In addition, multiple important confounding factors were included in the analysis, such as NSAID use and dietary information, which have been linked with the risk of colorectal cancer. However, two limitations should be taken into account. First, BMI and metabolic factors were measured at baseline, and we were not able to evaluate possible changes during follow-up. Therefore, there might be misclassification bias due to changes in these risk factors over time. As a consequence of aging, we would expect metabolically healthy women at baseline to be more likely to become metabolically unhealthy at follow-up, rather than the reverse. Thus, this type of misclassification bias would likely make our estimates conservative. Second, the study population consisted of four WHI subsets of women, which could be generalized to postmenopausal women, but not men or younger women.

In summary, a metabolically unhealthy phenotype was associated with higher risk of colorectal cancer among normal-weight postmenopausal women. These data suggest that normal-weight postmenopausal women should be evaluated for metabolic health and appropriate steps taken to reduce their risk of diseases, including colorectal cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Disclaimer

The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Authors' Contributions

Conception and design: X. Liang, M. Hendryx, J. Luo

Development of methodology: X. Liang, M. Stefanick, J. Luo

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): X. Liang, C.A. Thomson, D. Lane, M. Stefanick

References

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359–86.
2. American Cancer Society. *Cancer facts & figures 2016*. Atlanta, GA: American Cancer Society; 2016. Available from: <http://www.cancer.org/acs/groups/content/@research/documents/document/acspc-047079.pdf>
3. Esposito K, Chiodini P, Capuano A, Bellastella G, Maiorino MI, Rafaniello C, et al. Colorectal cancer association with metabolic syndrome and its components: a systematic review with meta-analysis. *Endocrine* 2013;44:634–47.
4. World Cancer Research Fund/American Institute for Cancer Research. Continuous update project report. Food, nutrition, physical activity, and the prevention of colorectal cancer. Washington, DC: American Institute for Cancer Research; 2011.
5. Aran V, Victorino AP, Thuler LC, Ferreira CG. Colorectal cancer: epidemiology, disease mechanisms and interventions to reduce onset and mortality. *Clin Colorectal Cancer* 2016;15:195–203.
6. Strum WB. Colorectal adenomas. *N Engl J Med* 2016;374:1065–75.
7. Conus F, Rabasa-Lhoret R, Peronnet F. Characteristics of metabolically obese normal-weight (MONW) subjects. *Appl Physiol Nutr Metab* 2007;32:4–12.
8. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002;106:3143–421.
9. Hinnouho GM, Czernichow S, Dugravot A, Batty GD, Kivimaki M, Singh-Manoux A. Metabolically healthy obesity and risk of mortality: does the definition of metabolic health matter? *Diabetes Care* 2013;36:2294–300.
10. Blüher M. The distinction of metabolically 'healthy' from 'unhealthy' obese individuals. *Curr Opin Lipidol* 2010;21:38–43.
11. Kramer C, Zinman B, Retnakaran R. Are metabolically healthy overweight and obesity benign conditions?: A systematic review and meta-analysis. *Ann Intern Med* 2013;159:758–69.
12. Wang B, Zhuang R, Luo X, Yin L, Pang C, Feng T, et al. Prevalence of metabolically healthy obese and metabolically obese but normal weight in adults worldwide: a meta-analysis. *Horm Metab Res* 2015;47:839–45.
13. Wildman RP, Muntner P, Reynolds K, McGinn AP, Rajpathak S, Wylie-Rosett J, et al. The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: prevalence

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): X. Liang, K.L. Margolis, M. Hendryx, E.J. Groessl, C.H. Kroenke, J. Luo

Writing, review, and/or revision of the manuscript: X. Liang, K.L. Margolis, M. Hendryx, T.E. Rohan, E.J. Groessl, C.A. Thomson, C.H. Kroenke, M.S. Simon, D. Lane, M. Stefanick, J. Luo

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): X. Liang

Study supervision: X. Liang, K.L. Margolis, C.A. Thomson, J. Luo

Acknowledgments

A short list of WHI investigators is as follows:

Program Office (National Heart, Lung, and Blood Institute, Bethesda, MD): Jacques Rossouw, Shari Ludlam, Joan McGowan, Leslie Ford, and Nancy Geller
Clinical Coordinating Center (Fred Hutchinson Cancer Research Center, Seattle, WA): Garnet Anderson, Ross Prentice, Andrea LaCroix, and Charles Kooperberg

Investigators and Academic Centers: (Brigham and Women's Hospital, Harvard Medical School, Boston, MA) JoAnn E. Manson; (MedStar Health Research Institute/Howard University, Washington, DC) Barbara V. Howard; (Stanford Prevention Research Center, Stanford, CA) Marcia L. Stefanick; (The Ohio State University, Columbus, OH) Rebecca Jackson; (University of Arizona, Tucson/Phoenix, AZ) Cynthia A. Thomson; (University at Buffalo, Buffalo, NY) Jean Wactawski-Wende; (University of Florida, Gainesville/Jacksonville, FL) Marian Limacher; (University of Iowa, Iowa City/Davenport, IA) Jennifer Robinson; (University of Pittsburgh, Pittsburgh, PA) Lewis Kuller; (Wake Forest University School of Medicine, Winston-Salem, NC) Sally Shumaker; (University of Nevada, Reno, NV) Robert Brunner; (University of Minnesota, Minneapolis, MN) Karen L. Margolis

Women's Health Initiative Memory Study: (Wake Forest University School of Medicine, Winston-Salem, NC) Mark Espeland

Grant Support

The WHI program is funded by the National Heart, Lung, and Blood Institute, NIH, and the U.S. Department of Health and Human Services, through contracts HHSN268201100046C, HHSN268201100001C, HHSN268201100002C, HHSN268201100003C, HHSN268201100004C, and HHSN271201100004C. Research reported in this publication was supported by the Youth Scholars Program of Beijing Normal University and by the NCI of the NIH under award number R15CA179463 (to J. Luo).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received September 25, 2016; revised November 10, 2016; accepted November 14, 2016; published OnlineFirst February 1, 2017.

- and correlates of 2 phenotypes among the US population (NHANES 1999–2004). *Arch Intern Med* 2008;168:1617–24.
14. Alfa-Wali M, Sharma A, Boniface S, Tekkis P, Hackshaw A, Antoniou A. Metabolic syndrome (MetS) and risk of colorectal cancer (CRC): a systematic review and meta-analysis. *Ann Oncol* 2012;23:22–3.
 15. Kabat GC, Kim M, Caan BJ, Chlebowski RT, Gunter MJ, Ho GY, et al. Repeated measures of serum glucose and insulin in relation to postmenopausal breast cancer. *Int J Cancer* 2009;125:2704–10.
 16. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome. An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005;112:2735–52.
 17. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412–9.
 18. Gunter MJ, Xie X, Xue X, Kabat GC, Rohan TE, Wassertheil-Smoller S, et al. Breast cancer risk in metabolically healthy but overweight postmenopausal women. *Cancer Res* 2015;75:270–4.
 19. Radikova Z, Koska J, Huckova M, Ksinantova L, Imrich R, Vigas M, et al. Insulin sensitivity indices: a proposal of cut-off points for simple identification of insulin-resistant subjects. *Exp Clin Endocrinol Diabetes* 2006;114:249–56.
 20. Curb JD, McTiernan A, Heckbert SR, Kooperberg C, Stanford J, Nevitt M, et al. Outcomes ascertainment and adjudication methods in the Women's Health Initiative. *Ann Epidemiol* 2003;13:S122–8.
 21. Anderson GL, Manson J, Wallace R, Lund B, Hall D, Davis S, et al. Implementation of the women's health initiative study design. *Ann Epidemiol* 2003;13:S5–S17.
 22. Murphy N, Cross AJ, Abubakar M, Jenab M, Aleksandrova K, Boutron-Ruault MC, et al. A nested case-control study of metabolically defined body size phenotypes and risk of colorectal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). *PLoS Med* 2016;13:e1001988.
 23. Kabat GC, Kim MY, Peters U, Stefanick M, Hou L, Wactawski-Wende J, et al. A longitudinal study of the metabolic syndrome and risk of colorectal cancer in postmenopausal women. *Eur J Cancer Prev* 2012;21:326–32.
 24. Ahmed RL, Schmitz KH, Anderson KE, Rosamond WD, Folsom AR. The metabolic syndrome and risk of incident colorectal cancer. *Cancer* 2006;107:28–36.
 25. Sturmer T, Buring JE, Lee IM, Gaziano JM, Glynn RJ. Metabolic abnormalities and risk for colorectal cancer in the physicians' health study. *Cancer Epidemiol Biomarkers Prev* 2006;15:2391–7.
 26. De Lorenzo A, Del Gobbo V, Premrov MC, Bigioni M, Galvano F, Di Renzo L. Normal-weight obese syndrome: early inflammation? *Am J Clin Nutr* 2007;85:40–5.
 27. Hyun YJ, Koh SJ, Chae JS, Kim JY, Kim OY, Lim HH, et al. Atherogenicity of LDL and unfavorable adipokine profile in metabolically obese, normal-weight woman. *Obesity* 2008;16:784–9.
 28. Dvorak RV, DeNino WF, Ades PA, Poehlman ET. Phenotypic characteristics associated with insulin resistance in metabolically obese but normal-weight young women. *Diabetes* 1999;48:2210–4.

Cancer Epidemiology, Biomarkers & Prevention

Metabolic Phenotype and Risk of Colorectal Cancer in Normal-Weight Postmenopausal Women

Xiaoyun Liang, Karen L. Margolis, Michael Hendryx, et al.

Cancer Epidemiol Biomarkers Prev 2017;26:155-161. Published OnlineFirst February 5, 2017.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-16-0761](https://doi.org/10.1158/1055-9965.EPI-16-0761)

Cited articles This article cites 26 articles, 7 of which you can access for free at:
<http://cebp.aacrjournals.org/content/26/2/155.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/26/2/155>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.