

# HLA Class II DR-DQ and Increased Risk of Cervical Cancer among Senegalese Women<sup>1</sup>

Patricia Lin, Laura A. Koutsky, Cathy W. Critchlow, Raymond J. Apple, Stephen E. Hawes, James P. Hughes, Papa Touré, Amadou Dembele, and Nancy B. Kiviat<sup>2</sup>

Departments of Pathology, School of Medicine [N. B. K.], Epidemiology [P. L., L. A. K., C. W. C., S. E. H.], and Biostatistics [J. P. H.], University of Washington, Seattle, Washington 98195; Roche Molecular Systems, Department of Human Genetics, Alameda, California 94501 [R. J. A.]; and Institut du Cancer, Dantec Hospital, Dakar, Senegal [P. T., A. D.]

## Abstract

To examine Senegalese women to confirm and extend associations between HLA class II types and cervical cancer previously observed among African-American, Caucasian, Hispanic, and Japanese ethnic populations, 55 Senegalese women with invasive cervical carcinoma were compared with age-matched (human papillomavirus) HPV-positive ( $n = 83$ ) and HPV-negative ( $n = 107$ ) control women. PCR-based HPV and HLA typing methods were used. Data were analyzed using a global randomization test and conditional logistic regression. Although this study failed to confirm a previously reported association between cervical cancer and *DQB1\*03* alleles, the *DRB1\*1101-DQB1\*0301* haplotype was detected more frequently among cervical carcinoma cases than among controls (adjusted odds ratio, 2.6; 95% confidence interval, 1.0–7.1). Furthermore, as reported by others, we observed a negative association of borderline statistical significance between *DRB1\*13* and cervical carcinoma (adjusted odds ratio, 0.5; 95% confidence interval, 0.2–1.1). Observations from this study confirm earlier findings of a negative association between *DRB1\*13* and cervical cancer and suggest that specific *DRB1-DQB1* haplotype combinations, rather than individual *DQB1\*03* alleles, increase the risk for cervical cancer.

## Introduction

Strong epidemiological evidence suggests that persistent infection with specific “high-risk” types of genital HPV<sup>3</sup> leads to development of cervical cancer (1, 2), the third most frequent cancer in women worldwide, with approximately 371,200 new cases reported every year (3). HPV DNA has been found in

nearly 100% of cervical cancers (4, 5), but in only 17–44% of control tissues (6). However, current evidence suggests that most HPV infections are self-limited and that only a minority of untreated cervical HPV infections progress to cervical cancer (7). Why only some untreated women infected with a high-risk HPV type develop invasive cancer is unclear; however one potential cofactor may be the host’s cellular immune response to HPV infection.

HLAs are cell surface proteins that play an important role in human cell-mediated immunity, with HLA class I proteins present on all cells and HLA class II proteins present on antigen-presenting cells. Several HLA types have been found to be associated with an increased risk of various diseases. Previous studies among African-American, Caucasian, Hispanic, and Japanese ethnic populations report associations between cervical neoplasia and *DRB1\*1101-DQB1\*0301* haplotype (8, 9), *DQB1\*03* alleles (8, 10–22), *DR5* (*DRB1\*11*, *DRB1\*12*) alleles (21, 23, 24), *DQB1\*0201* (14, 17), and *DQB1\*06* (13, 15, 17, 19, 25). However, disparate findings have been reported in different populations, possibly because of differences in HPV infection status or the extent to which preneoplastic lesions were discovered and treated before cancer developed. The present case-control study examined associations between invasive cervical carcinoma and specific HLA alleles *DQB1\*03*, *DRB1\*11*, *DQB1\*0201*, *DQB1\*06* and HLA haplotype *DRB1\*1101-DQB1\*0301* among a population of Senegalese women with and without HPV infection who had never undergone cervical cytological screening or treatment of preneoplastic lesions. Thus, our ability to distinguish associations of HLA alleles with invasive cervical cancer from those between HLA alleles and HPV infection (often manifested by morphological changes consistent with low-grade squamous intraepithelial lesions which frequently regress), in addition to the vast allelic diversity found among African and African-derived populations (26), provides a unique context for the interpretation of the role of specific HLA alleles in the etiology of cervical cancer.

## Materials and Methods

**Study Population.** Between March and December of 1998, all women 35 years of age or older presenting to the University of Dakar (Senegal) Oncology Clinic, the main cancer referral clinic in Senegal, with signs or symptoms of invasive cervical cancer were evaluated by colposcopy and biopsy to confirm the presence of invasive cervical cancer. A total of 55 women with histologically confirmed invasive disease were enrolled. Two control groups, individually matched to cases by age ( $\pm 3$  years), were recruited from among 1201 women 35 years of age or older seeking primary care or who presented for family planning purposes to the Pikine Outpatient Clinic of the University of Dakar during the same time period. All cases and potential controls provided written informed consent according to procedures approved by the institutional review boards at the

Received 8/16/00; revised 7/19/01; accepted 8/16/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> Supported by NIH Grants CA 75920 and CA 62801 (both to N. B. K.).

<sup>2</sup> To whom requests for reprints should be addressed, at University of Washington HPV Research Group, Lake Union Place, Suite 300, 1914 North 34th Street, Seattle, WA 98103.

<sup>3</sup> The abbreviations used are: HPV, human papillomavirus; OR, odds ratio; CI, confidence interval; CIN, cervical intraepithelial neoplasia.

University of Dakar and the University of Washington. The HPV-positive control group consisted of 83 women with normal cervical cytology findings and positive HPV DNA cervical swab results. The HPV-negative control group included 107 women with normal cytology findings and negative HPV DNA results. Controls were matched to the cases by age because HPV DNA detection is highly associated with age (27–32) and cancer incidence. Although we attempted to match each cervical carcinoma case with two controls, only 83 HPV-positive controls could be matched to the 55 cervical cancer cases because the referral population included too few cytologically normal women with positive tests for HPV DNA. All cases were matched to at least one control, but five cases (68–83 years of age) could not be matched with an HPV-positive control. At the time of the enrollment examination, all cases and controls had an intact cervix and were not pregnant.

**Study Procedures and Collection of Specimens.** All subjects completed a short standardized interview, including questions concerning medical, gynecological, and sexual history, and underwent a physical and gynecological examination. Cervical cellular samples were obtained for cytological screening, detection of HPV DNA and for analysis of HLA class II loci (*DRB1* and *DQB1*). Colposcopically directed biopsy specimens of the uterine cervix were obtained from all women with suspected cancer.

**Detection and Typing of HPV.** Cervical swab specimens were tested for presence and type of HPV using a PCR-based assay with L1 consensus primers MY09 and MY011 and then hybridization with a biotin-labeled generic HPV probe. If the sample tested positive using the generic probe, it was amplified again with biotinylated primers (MY09/MY11/HMB01). HPV detection was then performed through probe hybridization using a reverse-line blot-strip detection method, as described by Gravitt *et al.* (33). The HPV strip method signaled the presence of HPV DNA with two control beta-globin probes and typed for 27 specific genital HPV types.

**HLA Typing.** Human DNA was extracted from cervical swab samples and prepared for PCR amplification. All samples were typed for two HLA class II loci, *DQB1* and *DRB1*. Primers were provided by Roche Molecular Systems (Alameda, California). HLA class II line-blot typing strips and all liquid detection reagents were from the Amplicor Strip Detection Reagent Kit manufactured by Dynal (Oslo, Norway). HLA class II typing used single-stranded oligonucleotide probe reverse-dot blot methods as described by Erlich *et al.* (34) and Saiki *et al.* (35).

*DQB1* was amplified using 2–5  $\mu$ l of sample with a single amplification using biotinylated primers DB378B/DB380B. Forty PCR cycles were conducted on a Perkin-Elmer GeneAmp PCR System 9600 using the following cycling conditions: 95°C denaturation for 15 s; 60°C annealing for 45 s; and 72°C extension for 15 s. Amplified DNA was denatured and hybridized onto *DQB1* reverse line-blot typing strips in 20-well typing trays immersed in 4 $\times$  saline-sodium phosphate-EDTA-0.1% SDS hybridization and 1 $\times$  saline-sodium phosphate-EDTA-0.01% SDS stringent wash solutions in a 50°C shaking water bath. Color detection was performed using streptavidin-horse-radish peroxidase color conjugate and a hydrogen peroxidase-tetramethylbenzidine mixture. Positive probe patterns of the 25 immobilized sequence-specific oligonucleotide probes were entered into a computerized *DQB1* pattern-matching program, which were interpreted or deciphered into *DQB1*-specific alleles. This typing method is able to detect and differentiate between 30 different *DQB1* alleles.

General *DRB* typing was amplified using biotinylated primers RC1000/CRX28B with the same PCR cycling conditions described above for the *DQB1* locus. Amplified PCR product was denatured and hybridized onto *DRB* general strips with 29 sequence-specific immobilized probes using the same hybridization procedures and solutions described for the *DQB1* locus. The general typing strips categorized samples into *DRB1* sero-groups, *DRB1*-specific allele possibilities, and some *DRB3*, *DRB4*, and *DRB5* types, which were not used in this analysis. Because of the complexity of the *DRB1* locus, subgroup amplification was necessary for specific *DRB1* allele differentiation. The *DRB1* general amplification defined *DRB1\*07*, *DRB1\*09*, and *DRB1\*10* alleles, which did not need to be specified further. All other samples underwent a second round of amplification using group-specific biotinylated primers. The *DR1* group was amplified with primers DB1151/DB1035. The *DR2* group was amplified using primers DB1150/DB1035. The *DR3* group was amplified using primers DB1146/DB1035. The *DR4* group was amplified with DB1034/DB1035. The *DR8* group was amplified using primers DB1148/DB1035. Group-specific amplification methods and procedures were typed using the same hybridization protocol as described for the *DQB1* and *DRB1* general loci, but used 30-probe *DRB1*-specific strips. Positive probe patterns were entered into a computerized pattern-matching program or matched using a probe-pattern-typing grid to determine specific *DRB1* allele type. This *DRB* typing method was able to differentiate between 131 different *DRB1* alleles.

Haplotype configuration was determined according to known linkage disequilibrium patterns (36). HLA typing was performed by one laboratory technician who was unaware of the subject's case-control and HPV status.

**Histopathological and Cytological Methods.** Cervical cytological samples, collected directly into PreservCyt (Cytoc Corporation, Boxborough, MA) for liquid-based cytology (Thin-Prep 2000; Cytoc Corporation), were stained and evaluated as described previously (37) without knowledge of other clinical or laboratory data. Representative H&E-stained slides were prepared from paraffin-embedded biopsies, and reviewed by the pathologist (N. K.) without knowledge of other clinical or laboratory data. Standard gynecological pathology criteria and terminology were used to classify all intraepithelial lesions and invasive cervical cancers (38).

**Statistical Analysis.** To reduce the possibility of reporting spurious associations which might have been detected solely because of the large number of comparisons needed to assess all of the individual HLA types, a two-step approach was taken in the analysis of these data. First, associations between cervical cancer and particular HLA types that had been reported previously more than once among other ethnic populations were assessed in our population by conditional logistic regression, which took into account the individual matching by age. Second, a global randomization test was used to determine whether any of the remaining HLA types (that is, those types not hypothesized *a priori* to be associated with cervical cancer) were associated with cervical cancer beyond what might be expected to occur by chance alone.

Because controls were individually matched to the cases by age, conditional logistic regression analyses were used to obtain ORs, 95% CIs, and *Ps* assessing *a priori* hypothesized associations between HLA alleles/haplotypes and invasive cervical cancer. Confounding of the association between cervical carcinoma and certain HLA alleles was ascertained by identifying characteristics that varied in distribution between cases

and controls. A factor was considered to be a potential confounder if the characteristic also differed substantially with the presence or absence of HLA alleles among controls and changed the crude OR by >10%. Analyses were performed comparing cases to HPV-positive and HPV-negative control groups separately. The control groups were combined for the final analyses because there were no significant differences between the two groups. Statistical analysis was performed using SAS v.6.12 (Cary, NC).

The association between HLA type and case-control status was also assessed by means of a randomization test. First, the observed  $\chi^2$  statistic comparing cases with each control group was calculated. The case-control indicator was then randomly permuted within each matched stratum and the  $\chi^2$  statistic was recomputed. This procedure was repeated 1000 times, providing a distribution of the  $\chi^2$  statistic under the null hypothesis of no association between case-control status and HLA haplotype. A *P* was calculated by comparing the observed  $\chi^2$  statistic with this null distribution.

We also compared the risk of cervical cancer associated with HLA homozygosity (for *DRB1* and *DQB1* alleles) with heterozygosity using conditional logistic regression analyses, because homozygosity may confer a greater cancer risk inasmuch as less variation in the HLA region may be associated with a narrower range of specific antigens recognized by the host immune system.

## Results

**Characteristics of the Study Population.** The mean age of women enrolled was 48 years (range, 35–83 years). Although the controls were age-matched ( $\pm 3$  years) to the cases, HPV-positive controls were significantly younger than cases (45.8 years *versus* 49.5 years; *P* < 0.05; Table 1). Women in both control groups were similar to cases with respect to ethnicity, marital status (including polygamous *versus* monogamous marriage), and smoking habits. A higher proportion of cases, as compared with HPV-positive controls and HPV-negative controls, were born outside of Dakar (*P* < 0.05).

Women with cervical carcinoma tended to have fewer children (mean, 4.9) than did women in the control groups (HPV-positive, mean, 5.8; HPV-negative, mean, 6.1). Women in both control groups were more likely than cancer cases to practice contraception; however, no one reported the use of condoms.

**Detection of HPV.** Of the women with cervical carcinoma, 64.7% tested positive for HPV DNA, with 37.3% positive for HPV 16, 7.8% for HPV 18, and 18.2% for other HPV types. Of the HPV-positive controls, 4.8% tested positive for HPV type 16, 10.8% tested positive for HPV type 18, and 84.3% were positive for other HPV types.

**Distribution of HLA Haplotypes among Cases and Controls.** Class II HLA distributions for the *DQB1* and *DRB1* were compared among HPV-positive controls, HPV-negative controls, and cervical cancer cases (Table 2). Of the possible 131 *DRB1* and 30 *DQB1* alleles that could be ascertained with the assay used in this study, 25 different *DRB1* alleles and 13 *DQB1* alleles were actually detected.

**Hypothesized HLA Associations between Cervical Cancer Cases and HPV-positive and HPV-negative Controls.** Comparison of cervical carcinoma cases to the HPV-positive controls with respect to the class II HLA alleles and haplotypes that were hypothesized *a priori* to be associated with cervical cancer revealed no statistically significant associations (Table 3).

Table 1 Demographic characteristics of cervical carcinoma cases and HPV-positive and HPV-negative controls

Characteristic	Cases <i>n</i> = 55	Controls	
		HPV-positive <i>n</i> = 83	HPV-negative <i>n</i> = 107
Age (mean yr $\pm$ SD)	49.5 $\pm$ 12	45.8 $\pm$ 8 <sup>a</sup>	47.9 $\pm$ 10
Ethnicity			
Wolof	22 (40.0%)	21 (51.2%) <sup>b</sup>	26 (54.2%) <sup>b</sup>
Pular	15 (27.3%)	8 (19.5%)	11 (22.9%)
Other	18 (32.7%)	12 (29.1%)	11 (23.3%)
Origin (birthplace)			
Dakar	35 (63.6%)	83 (100.0%) <sup>a</sup>	104 (97.2%) <sup>a</sup>
Other	20 (36.7%)	0 (0%)	3 (2.8%)
Marital status			
Monogamously married	13 (23.6%)	22 (26.5%)	29 (27.4%) <sup>c</sup>
Polygamously married	26 (47.3%)	43 (51.8%)	58 (54.7%)
Unmarried	16 (29.1%)	18 (21.6%)	19 (17.9%)
Mean no. of live-born infants $\pm$ SD	4.9 $\pm$ 3	5.8 $\pm$ 2 <sup>a</sup>	6.1 $\pm$ 3 <sup>a</sup>
Current method of birth control			
None	50 (90.9%)	51 (61.4%) <sup>a</sup>	83 (79.0%) <sup>a,d</sup>
Pill	2 (3.6%)	5 (6.0%)	3 (2.9%)
Injection/Norplant	3 (5.5%)	5 (6.0%)	6 (5.7%)
Spermicide	0 (0%)	1 (1.2%)	1 (1.0%)
IUD	0 (0%)	15 (18.1%)	10 (9.5%)
Ligature	0 (0%)	3 (3.6%)	1 (1.0%)
Other	0 (0%)	3 (3.6%)	1 (1.0%)

<sup>a</sup> *P* < 0.05 comparing controls with cases; for birth control method, nonusers were compared with users.

<sup>b</sup> Forty or more missing values.

<sup>c</sup> One missing value.

<sup>d</sup> Two missing values.

A comparison of the frequency of these *DQB1* and *DRB1* alleles between cases and HPV-negative controls revealed a significant association for *DQB1*\*02 (specifically *DQB1*\*0201/2). The *DRB1*\*1101-*DQB1*\*0301 haplotype was found to be associated with cervical carcinoma, particularly when compared with HPV-negative controls (adjusted OR, 7.4; 95% CI, 1.5–36.5). Although associations between cervical cancer and specific class II HLA alleles and haplotypes did not achieve statistical significance in analyses using the HPV-positive control group, ORs from these analyses were similar in magnitude to those obtained using the HPV-negative control group, suggesting that the control groups could be combined (Table 3). Although the positive association between the *DRB1*\*1101-*DQB1*\*0301 haplotype and cervical carcinoma was stronger when comparing cases with HPV-negative controls, the association persisted when cases were compared with the combined control groups (adjusted OR, 2.6; 95% CI, 1.0–7.1). The individual *DRB1*\*1101 and *DQB1*\*0301 alleles alone were not associated with cervical carcinoma.

**Other Potential Allele and Haplotype Associations Examined by Use of a Global Randomization Test.** The global randomization test was used to test whether HLA alleles or haplotypes not considered in our *a priori* hypotheses could be associated with invasive cervical cancer in this population. Using this test, no statistically significant difference in HLA allele or haplotype distribution between cases and HPV-positive controls (*P* = 0.72) or cases and HPV-negative controls (*P* = 0.14) was observed.

**HLA Homozygosity and Cervical Cancer Risk.** Women homozygous compared with those heterozygous for an associated HLA allele may be at greater risk for developing cervical disease, perhaps caused by less variation in the HLA region. In our study population, 18 (7.4%) had homozygous *DRB1*-*DQB1*

Table 2 Distribution of class II HLA alleles among cervical carcinoma cases and HPV-positive and HPV-negative controls [n (%)]

Allele	Cases (n = 55) <sup>a</sup>	Controls		DRB1-DQB1 haplotype	Cases (n = 55) <sup>a</sup>	Controls	
		HPV + (n = 83) <sup>a</sup>	HPV - (n = 107) <sup>a</sup>			HPV + (n = 83) <sup>a</sup>	HPV - (n = 107) <sup>a</sup>
<i>DRB1</i>							
0101	1 (0.9%)	0 (0.0%)	2 (0.9%)	0101-0501	1 (0.9%)	0 (0.0%)	2 (0.9%)
0102	3 (2.7%)	5 (3.0%)	4 (1.9%)	0102-0501	3 (2.7%)	5 (3.0%)	4 (1.9%)
0301	6 (5.4%)	10 (6.0%)	8 (3.8%)	0301-0201/2	6 (5.5%)	9 (5.4%)	8 (3.7%)
0302	5 (4.5%)	13 (7.8%)	16 (7.5%)	0301-0203	0 (0.0%)	1 (0.6%)	0 (0.0%)
0401	0 (0.0%)	1 (0.6%)	1 (0.5%)	0301-0501	0 (0.0%)	1 (0.6%)	0 (0.0%)
0403	0 (0.0%)	1 (0.6%)	2 (0.9%)	0302-0402	5 (4.5%)	13 (7.8%)	15 (7.0%)
0405	2 (1.8%)	5 (3.0%)	9 (4.2%)	0302-0501	0 (0.0%)	0 (0.0%)	1 (0.5%)
0701	10 (9.1%)	8 (4.8%)	5 (2.4%)	0401-0301	0 (0.0%)	1 (0.6%)	0 (0.0%)
0804	5 (4.5%)	7 (4.2%)	8 (3.8%)	0401-0302	0 (0.0%)	0 (0.0%)	1 (0.5%)
0806	2 (1.8%)	3 (1.8%)	1 (0.5%)	0403-0302	0 (0.0%)	1 (0.6%)	1 (0.5%)
0901	10 (9.1%)	6 (3.6%)	8 (3.8%)	0405-0201/2	2 (1.8%)	1 (0.6%)	2 (0.9%)
1001	17 (15.5%)	17 (10.2%)	28 (13.2%)	0405-0302	0 (0.0%)	4 (2.4%)	0 (0.0%)
1101	12 (10.9%)	22 (13.3%)	13 (6.1%)	0701-0201/2	10 (9.1%)	8 (4.8%)	4 (1.9%)
1102	5 (4.5%)	6 (3.6%)	24 (11.3%)	0701-0203	0 (0.0%)	0 (0.0%)	1 (0.5%)
1104	1 (0.9%)	1 (0.6%)	0 (0.0%)	0804-0301	5 (4.5%)	7 (4.2%)	7 (3.3%)
1201	0 (0.0%)	0 (0.0%)	3 (1.4%)	0804-0604	0 (0.0%)	0 (0.0%)	1 (0.5%)
1202	0 (0.0%)	0 (0.0%)	1 (0.5%)	0806-0602	2 (1.8%)	3 (1.8%)	1 (0.5%)
1301	3 (2.7%)	6 (3.6%)	17 (8.0%)	0901-0201	9 (8.2%)	6 (3.6%)	7 (3.3%)
1302	5 (4.5%)	16 (9.6%)	19 (9.0%)	0901-0203	1 (0.9%)	0 (0.0%)	1 (0.5%)
1303	4 (3.6%)	7 (4.2%)	4 (1.9%)	1001-0501	17 (15.5%)	17 (10.2%)	28 (13.1%)
1304	11 (10.0%)	26 (15.7%)	30 (14.2%)	1101-0203	0 (0.0%)	0 (0.0%)	1 (0.5%)
1322	1 (0.9%)	3 (1.8%)	3 (1.4%)	1101-0301	11 (10.0%)	12 (7.2%)	6 (2.8%)
1401	2 (1.8%)	0 (0.0%)	1 (0.5%)	1101-0501	1 (0.9%)	3 (1.8%)	5 (2.3%)
1503	3 (2.7%)	0 (0.0%)	1 (0.5%)	1101-0502	0 (0.0%)	1 (0.6%)	0 (0.0%)
1602	2 (1.8%)	3 (1.8%)	4 (1.9%)	1101-0602	0 (0.0%)	5 (3.0%)	0 (0.0%)
<i>DQB1</i>							
0201/2	27 (24.5%)	26 (15.7%)	24 (11.2%)	1101-0604	0 (0.0%)	1 (0.6%)	1 (0.5%)
0203	1 (0.9%)	1 (0.6%)	3 (1.4%)	1102-0301	5 (4.5%)	5 (3.0%)	23 (10.7%)
0301	37 (33.6%)	57 (34.3%)	71 (33.2%)	1102-0302	0 (0.0%)	1 (0.6%)	0 (0.0%)
0302	0 (0.0%)	8 (4.8%)	11 (5.1%)	1104-0301	1 (0.9%)	1 (0.6%)	0 (0.0%)
0402	5 (4.5%)	13 (7.8%)	15 (7.0%)	1201-0301	0 (0.0%)	0 (0.0%)	3 (1.4%)
0501	25 (22.7%)	36 (21.7%)	51 (23.8%)	1202-0602	0 (0.0%)	0 (0.0%)	1 (0.5%)
0502	2 (1.8%)	5 (3.0%)	4 (1.9%)	1301-0201/2	0 (0.0%)	0 (0.0%)	2 (0.9%)
0503	2 (1.8%)	0 (0.0%)	1 (0.5%)	1301-1302	0 (0.0%)	1 (0.6%)	1 (0.5%)
0602	4 (3.6%)	9 (5.4%)	7 (3.3%)	1301-0602	0 (0.0%)	1 (0.6%)	0 (0.0%)
0603	4 (3.6%)	5 (3.0%)	18 (8.4%)	1301-0603	3 (2.7%)	3 (1.8%)	14 (6.5%)
0604	2 (1.8%)	3 (1.8%)	8 (3.7%)	1302-0201/2	0 (0.0%)	1 (0.6%)	0 (0.0%)
0605	0 (0.0%)	1 (0.6%)	0 (0.0%)	1302-0501	2 (1.8%)	10 (6.0%)	13 (6.1%)
0609	1 (0.9%)	2 (1.2%)	1 (0.5%)	1302-0502	0 (0.0%)	1 (0.6%)	0 (0.0%)
				1302-0604	2 (1.8%)	1 (0.6%)	5 (2.3%)
				1302-0605	0 (0.0%)	1 (0.6%)	0 (0.0%)
				1302-0609	1 (0.9%)	2 (1.2%)	1 (0.5%)
				1303-0201/2	0 (0.0%)	1 (0.6%)	1 (0.5%)
				1303-0301	4 (3.6%)	6 (3.6%)	3 (1.4%)
				1304-0301	11 (10.0%)	25 (15.1%)	29 (13.6%)
				1304-0302	0 (0.0%)	1 (0.6%)	0 (0.0%)
				1304-0604	0 (0.0%)	0 (0.0%)	1 (0.5%)
				1322-0603	1 (0.9%)	2 (1.2%)	3 (1.4%)
				1322-0604	0 (0.0%)	1 (0.6%)	0 (0.0%)
				1401-0503	2 (1.8%)	0 (0.0%)	1 (0.5%)
				1503-0301	1 (0.9%)	0 (0.0%)	0 (0.0%)
				1503-0602	2 (1.8%)	0 (0.0%)	1 (0.5%)
				1602-0502	2 (1.8%)	3 (1.8%)	4 (1.9%)

<sup>a</sup> Each subject contributes two alleles (haplotypes) to the analysis, thus, the total number of alleles (haplotypes) in each column is twice the number of subjects. One HPV negative control sample could not be HLA-typed for the *DRB1* locus (it was known to be positive for alleles in the *DRB1\*03*, *\*11*, or *\*13* groups) and was excluded from the analyses.

haplotypes, whereas 7 (2.9%) were homozygous at the *DRB1* locus, 32 (13.1%) were homozygous at the *DQB1* locus, and the remaining 187 (76.6%) subjects were heterozygous at both the *DRB1* and *DQB1* loci.

Detection of homozygous *DRB1-DQB1* haplotypes was positively associated with invasive cervical cancer (adjusted OR, 5.5, 95% CI, 1.5–20.2), whereas homozygosity for *DRQ1*

alone (OR, 1.1; 95% CI, 0.3–3.7) was not significantly associated with risk of cervical cancer. None of the seven subjects homozygous for *DRB1* alone had cervical cancer, although the sample size was too small to determine whether this represented a negative association with cervical cancer.

Among the *a priori* hypothesized associations on which we focused our analysis (*DR11*, *DR13*, *DQ02*, *DQ03*, and

Table 3 Crude and adjusted ORs for the association between cervical carcinoma and *a priori* hypothesized class II HLA alleles and haplotypes

<i>DQB1</i> and <i>DRB1</i> alleles	Cases vs.	Cases vs.	Cases vs.	Cases vs.	Cases vs.
	HPV + controls Crude OR (95% CI)	HPV - controls Crude OR (95% CI)	HPV-controls Adjusted OR <sup>a</sup> (95% CI)	combined controls Crude OR (95% CI)	combined controls Adjusted OR <sup>a</sup> (95% CI)
<i>DQB1</i> *0201/2	1.4 (0.7–2.8)	2.9 (1.3–6.3)	1.1 (0.4–3.1)	2.1 (1.1–3.9)	1.2 (0.5–2.8)
<i>DQB1</i> *03	0.6 (0.3–1.2)	0.8 (0.4–1.6)		0.8 (0.4–1.4)	0.8 (0.4–1.9)
<i>DQB1</i> *0602	0.5 (0.1–2.2)	0.9 (0.2–3.3)		0.7 (0.2–2.4)	0.6 (0.1–2.7)
<i>DRB1</i> *13	0.5 (0.6–1.1)	0.5 (0.3–1.0)	0.5 (0.2–1.4)	0.5 (0.3–1.0)	0.5 (0.2–1.1)
<i>DRB1</i> *11	0.8 (0.3–1.7)	0.9 (0.4–1.8)		0.8 (0.4–1.6)	1.0 (0.4–2.2)
<i>DRB1</i> *1101- <i>DQB1</i> *0301	1.4 (0.5–3.5)	3.6 (1.2–10.6)	7.4 (1.5–36.5)	2.1 (0.9–4.9)	2.6 (1.0–7.1)

<sup>a</sup> Adjusted for birthplace, no. of live-born infants, age, and other alleles in Table.

*DQ06* alleles), we evaluated whether there was a difference in association with cervical cancer among those homozygous for those alleles *versus* those heterozygous for alleles. Using conditional logistic regression analysis to account for the individual matching of cases to controls and to control for potential confounding factors, we found that heterozygosity for *DR13* was somewhat negatively (but not significantly) associated with cervical cancer (OR, 0.5; 95% CI, 0.2–1.3), although none of the nine subjects homozygous for *DR13* had cervical cancer. Neither heterozygosity nor homozygosity for *DR11* and *DQ03* were associated with risk of cervical cancer. *DQ06* was not tested because there were only two subjects in our population that were homozygous for *DQB1*\*06 alleles. However, those homozygous for *DQB1*\*02 were significantly more likely to have cervical cancer (adjusted OR, 6.5; 95% CI, 1.1–36.9), whereas heterozygosity for *DQB1*\*02 did not seem to be associated with cervical cancer risk (adjusted OR, 1.0; 95% CI, 0.4–2.4).

## Discussion

Among this group of previously unscreened Senegalese women, we observed *DRB1*\*1101-*DQB1*\*0301 to be more common among cervical carcinoma cases compared with controls. This association has been reported in previous investigations of British women with CIN (8) and Swedish women with cervical carcinoma who were positive for HPV types other than HPV 16 (9). However, Sanjeevi *et al.* (39), in a similar study among Swedish women, did not find the same haplotype association with CIN among HPV 16-seronegative subjects.

Although we observed an association between cervical carcinoma and a specific *DRB1*-*DQB1* haplotype containing *DQB1*\*03, we were unable to detect an association between the *DQB1*\*03 allele group and cervical carcinoma found in other studies. Several studies have observed an increase in frequency of *DQB1*\*03 alleles with dysplasia among Spanish women (15), with CIN among British women (8, 10, 23), and with cervical carcinoma among African-American (17), British (12), German (11), Japanese (13), Norwegian (14), and United States (16, 40) women. One report also found women homozygous for *DQB1*\*03 were more likely to have CIN than women heterozygous for this allele group (23). However, a number of other studies reported no association between *DQB1*\*03 alleles and CIN among Europeans (18) and cervical carcinoma among northwest English (24, 41), Brazilian (42), Hispanic (19), French (20), and Swedish (9) women. The findings from our analysis support observations of no association between the *DQB1*\*03 allele and cervical cancer.

Additionally, despite our observation of an association between a *DRB1*\*11-*DQB1*\*03 haplotype and cervical carcinoma,

we did not observe an association with *DRB1*\*11 alone. A strong association of *DRB1*\*11 with CIN was observed among British women (23). Syrjänen *et al.* (21) showed an increase in frequency of *DR5* (*DRB1*\*11, *DRB1*\*12) among women with high-grade CIN and women with HPV 16, and although Glew *et al.* (24) found no significant *HLA* associations among cervical carcinoma patients, they noted that all patients possessing *DR5* had HPV 16-positive tumors.

The first report to observe an increase in frequency of *HLA*-*DR5* antigen among German women with cervical carcinoma suggested the association was attributable to linkage disequilibrium between *DR5* and *DQw3* (11). Our study suggests that the combination of *DR11* with the *DQB1*\*0301 allele is associated with cervical carcinoma, because *DRB1*\*1101 and *DQB1*\*0301 alleles alone did not show any association. Individually, each allele was observed in similar frequencies among cases and controls (*DRB1*\*1101: cases 10.9%, combined controls 9.3%; *DQB1*\*0301: cases 33.6%, combined controls 33.7%). However, the *DRB1*\*11-*DQB1*\*03 haplotype was detected more often among cases than controls. This finding suggests that *DQB1*\*03 may be a marker, which, when coupled with *DRB1*\*1101 or another gene in linkage disequilibrium with the *DRB1*\*1101-*DQB1*\*0301 haplotype, may influence the development of HPV-associated cancers. Our study was able to make this distinction because *DR11* is observed in haplotype combinations with *DQ* alleles other than *DQB1*\*0301 among African groups, whereas *DR11* is found in linkage disequilibrium with *DQB1*\*0301 in most other ethnic groups (26). Another analysis of the *HLA* influence on cervical carcinoma among African-American women (17) only conducted *DQB1* analysis, and thus was unable to test whether particular alleles in linkage disequilibrium have an (interactive) effect on the development of cervical carcinoma. Variations in *HLA* haplotype distributions among different ethnic populations may explain important differences between studies.

Our study found evidence of a negative association between *DRB1*\*13 alleles and cervical carcinoma consistent with previous reports (11, 16, 18–22). *DRB1*\*13 has been found with lower frequency among Hispanic women with cervical carcinoma (OR, 0.3; 95% CI: 0.1–0.7; *P* = 0.001; Ref. 19) and French women (OR, 0.3; *P* = 0.0004; Ref. 20) compared with controls. A decreased cancer risk associated with *DR6* (*DRB1*\*13 and *DRB1*\*14) has also been observed among German (11), Mexican (22), and Finnish women (21). Additionally, a *DR13*-serotype-associated *DQ* allele (*DQB1*\*0603) was found to have a negative association with invasive squamous cell carcinoma of the cervix in a Spanish population (15). Our study extends these previous findings by showing similar results among West African women.

We found that our crude results required adjustment for

place of birth, number of live-born infants, and age, all recognized risk factors for cervical cancer, in addition to factors that we found differed in frequency between the various *HLA* alleles. Additionally, we simultaneously adjusted our estimates for the other *a priori* hypothesized alleles and haplotypes. After adjusting for potentially confounding variables, we did not detect an association between *DQB1\*0201/2* and cervical carcinoma in our study population. This contrasts with studies that have observed a decrease in frequency of *DQB1\*0201* among African-American (17) and Norwegian women (14) with cervical cancer as compared with controls. However, we did find that homozygosity for *DQB1\*02* was positively associated with cervical cancer. We found no associations between *DQB1\*06*, *\*0602*, *\*0603*, or *\*0604* and cervical carcinoma in our study population. Other investigators have observed an increase of *DQB1\*0602* (19, 25), *DQB1\*0603* (15), *DQB1\*0604* (17), and *DQ1* (*DQB1\*05* and *DQB1\*06*; Ref. 13) among cervical carcinoma patients compared with controls. A number of studies have detected an association between *DQB1\*0602* and cervical carcinoma in the *DRB1\*1501-DQB1\*0602* haplotype (19, 42). Perhaps *DQB1\*06* is associated with cervical cancer when it is in linkage disequilibrium with *DRB\*15*. Other studies have observed an increased frequency of *DR15* among cervical carcinoma patients (18, 42); although this association remains uncertain, the results of a study by Hildesheim *et al.* (16) found *DRB1\*1501-DQB1\*0602* to be decreased among CIN patients. We were not able to assess this haplotype association, because our Senegalese population did not contain the *DRB1\*1501-DQB1\*0602* haplotype.

People homozygous for positively or negatively associated alleles may show a greater predisposition for disease than those heterozygous for such alleles, because the *HLA* locus is responsible for mounting an immune response against infections. Few studies have examined *HLA* homozygosity as it relates to cervical cancer. Of the studies that have conducted this analysis, one found homozygosity for *DQB1\*03* to be positively associated with cervical HPV infection (12), whereas another found homozygosity for *DQB1\*0302* to be associated with a higher risk of CIN (16). Our analyses did not support these previous findings.

The *DRB1* and *DQB1* distribution of alleles of the combined controls was similar to that of another *HLA* study among a Senegalese population (43). Other *HLA* studies of African and West African populations (44, 45) show similar class II *HLA* trends (Appendix). *DRB1\*1304*, extremely rare in other populations, is the most prevalent *DRB1* allele in these West African populations.

Various *HLA* alleles have been found at a higher or lower frequency among HPV-positive and HPV-negative control groups. These differences suggest that certain *HLA* alleles may play a role in the persistence or clearance of HPV. However, because *HLA* loci *DRB1* and *DQB1* are so polymorphic, there is the potential for a few alleles to differ between the two groups by chance. Several studies have noted no difference in *HLA* distribution among HPV-positive and HPV-negative controls (19, 39–40). Our comparison supports those observations, suggesting *HLA* type is not associated with susceptibility, persistence, or detection of HPV infection.

Sixty-four percent of the cervical cancer patients in our study population tested positive for HPV at the time of enrollment into the study. This percentage is lower than reported in most studies of HPV DNA detection in cervical cancer patients (5). However, because of the advanced stages of cancer among many of the cervical carcinoma cases at the time of enrollment, several tissue samples were necrotic and may have been oth-

erwise histologically inadequate for sampling and testing. Recent evidence questions the existence of true HPV negative cervical carcinomas and suggests that inadequate histological sampling, integration of HPV genome into human DNA (46), or disruption of the L1 region of HPV DNA (which is normally used for HPV detection and typing) may account for false-negative or undetectable HPV test results among cervical cancer cases (47, 48).

One of the limitations of our study was the relatively small number of available cervical carcinoma cases. However, even with the small number of cases in this study, we were still able to detect associations reported in previous studies. Additionally, in studies involving multiple comparisons, it is possible that a significant result could be found by chance. The associations reported in our study were based on *a priori* hypotheses.

The postulated role of *HLA* class II alleles in the risk of cervical neoplasia is in their ability to elaborate proteins that facilitate immune recognition and control of HPV replication. If *HLA* expression is suppressed, lost, or increased among HPV-infected individuals of a particular *HLA* type, this may suggest that the ability or inability to ward off HPV infection is related to *HLA* type. Studies have shown up-regulation of *HLA* class II expression among women with cervical carcinoma (49–52). HPV infection seems to induce *HLA* class II expression in cervical squamous epithelium, which is not normally expressed in this region. Furthermore, *HLA* class II antigens have been observed to be expressed differentially, with *DRB* expression increased relative to *DQB* antigens (52, 53). A loss of *HLA* class I expression has been observed among cervical carcinoma patients (49–52, 54, 55). Recent studies have found altered *HLA* class I antigen expression can be attributed to specific genetic mutations (55, 56). Understanding the effects of changes in *HLA* expression subsequent to HPV exposure may explain biological mechanisms in which HPV interacts with *HLA* related to the development of susceptibility to, or protection against, cervical cancer.

This study of Senegalese women supports previous findings that the *HLA* haplotype *DRB1\*1101-DQB1\*0301* is increased among cervical carcinoma cases compared with controls. We were unable to confirm any association between *DQB1\*03* and cervical carcinoma, except when coupled with *DRB1\*1101*. These observations suggest that specific *DRB1-DQB1* haplotype combinations, rather than individual *DRB1* or *DQB1* alleles, are important in cervical carcinoma susceptibility. Because *DRB1* alleles appear in linkage disequilibrium with different *DQB1* alleles among different ethnic populations, variation in haplotype combinations may help explain the different results observed in previous studies of *HLA* and cervical carcinoma risk. Our study of Senegalese women confirms and extends findings from previous studies in other ethnic groups showing associations between specific *HLA* class II molecules and invasive cervical cancer.

#### Acknowledgments

We thank the following people who helped make this study possible: Dr. Henry Erlich of Roche Molecular Systems in Alameda, CA, for his review and commentary on this paper; Charlotte Sarr, clinician in Pikine; Deana Rich, Elise Reay-Ellers, and Macoumba Touré, study coordinators in Senegal; Fatou Faye Diop, data entry personnel in Senegal; Haby Agne, laboratory personnel in Senegal; Alison Starling, in charge of data management in Seattle; and Jane Kuypers, in charge of the HPV laboratory in Seattle.

Appendix HLA class II allele and haplotype distribution among Senegalese study population controls in comparison with other HLA-typed West African populations

DRB1 allele	n	%	Senegal [43] <sup>a</sup>	DRB1-DQB1 haplotype	n	%	Gambia [44] <sup>b</sup>	Malawi [44] <sup>b</sup>
0101	2	0.5%	0.6%	0101-0501	2	0.5%	DR1-DQ5: 4.1%	4.6%
0102	9	2.4%	0.6%	0102-0501	9	2.4%		
0301	17	4.5%	9.0%	0301-0201/2	16	4.3%	DR0301-DQw2: 2.7%	7.3%
0302	29	7.7%	2.8%	0301-0203	1	0.3%		
0401	2	0.5%		0301-0501	1	0.3%		
0403	3	0.8%	0.4%	0302-0402	28	7.4%	DR0302-DQw4: 3.4%	6.4%
0405	14	3.7%	0.6%	0302-0501	1	0.3%		
0701	13	3.5%	7.5%	0401-0301	1	0.3%	DR4-DQB*0301: 0.1%	0.0%
0801			2.8%	0401-0302	1	0.3%	DR4-DQB1*0302: 4.5%	2.7%
0802			7.2%	0403-0302	3	0.8%		
0803			1.3%	0405-0201/2	3	0.8%	DR4-DQw2: 0.9%	0.3%
0804	15	4.0%		0405-0302	11	2.9%		
0806	4	1.1%		0701-0201/2	12	3.2%	DR7-DQw2: 4.3%	4.3%
0901	14	3.7%	1.4%	0701-0203	1	0.3%		
1001	45	12.0%	4.3%	0804-0301	14	3.7%	DR8-DQw7: 3.0%	4.3%
1101	35	9.3%	9.7%	0804-0604	1	0.3%	DR3-DQw6: 0.3%	0.0%
1102	30	8.0%	6.0%	0806-0602	4	1.1%		
1103			0.2%	0901-0201/2	13	3.5%	DR9-DQw2: 7.1%	10.4%
1104	1	0.3%	0.7%	0901-0203	1	0.3%		
1201	3	0.8%	0.4%	1001-0501	45	12.0%	DR10-DQw5: 8.9%	0.9%
1202	1	0.3%		1101-0203	1	0.3%	DR1101-DQw2: 0.2%	0.0%
1301	23	6.1%	4.3%	1101-0301	18	4.8%	DR1101-DQ0301: 4.2%	6.4%
1302	34	9.0%	4.5%	1101-0501	5	1.3%	DR1101-DQw5: 0.1%	0.0%
1303	11	2.9%	1.5%	1101-0502	1	0.3%		
1304	56	14.9%	27.1%	1101-0602	9	2.4%	DR1101-DQw6: 2.0%	8.5%
1305			0.5%	1101-0604	2	0.5%		
1322	6	1.6%		1102-0301	28	7.4%		
1401	1	0.3%	0.8%	1102-0302	1	0.3%		
1503	1	0.3%		1104-0301	1	0.3%		
1601			1.6%	1201-0301	3	0.8%	DR12-DQ0301: 0.4%	0.0%
1602	7	1.9%		1202-0602	1	0.3%		
Missing	2			1301-0201/2	2	0.5%		
Total (missing)	376			1301-0302	2	0.5%		
DQB1 allele	n	%		1301-0602	1	0.3%	DR13-DQw6: 3.6%	4.2%
0201/2	49	13.0%		1301-0603	17	4.5%		
0203	4	1.1%		1302-0201/2	1	0.3%	DR13-DQw2: 0.6%	0.0%
0301	128	33.9%		1302-0501	22	5.9%	DR13-DQw5: 0.3%	0.3%
0302	19	5.0%		1302-0502	1	0.3%	DR1302-DQw1: 16.4%	7.6%
0402	28	7.4%		1302-0604	6	1.6%		
0501	86	22.8%		1302-0605	1	0.3%		
0502	9	2.4%		1302-0609	3	0.8%		
0503	1	0.3%		1303-0201/2	2	0.5%	1303-0301: 2.2%	0.9%
0602	16	4.2%		1303-0301	9	2.4%	1304-0301: 27.3%	3.0%
0603	23	6.1%		1304-0301	54	14.4%		
0604	11	2.9%		1304-0302	1	0.3%		
0605	1	0.3%		1304-0604	1	0.3%		
0609	3	0.8%		1322-0603	5	1.3%		
Missing	0			1322-0604	1	0.3%		
Total (missing)	378			1401-0503	1	0.3%	DR14-DQw1: 0.5%	0.9%
				1503-0301	0	0.0%		
				1503-0602	1	0.3%	DR15-DQw6: 0.8%	13.4%
				1602-0502	7	1.9%	DR16-DQw5: 1.5%	0.3%
				Missing	2			
				Total (missing)	376			

<sup>a</sup> Tiercy et al.: Mandenka (Mandenkalu) of Senegal.

<sup>b</sup> Hill et al.: The Gambia (West Africa) and Malawi (South-Central Africa).

## References

- Schiffman, M. H., Bauer, H. M., Hoover, R. N., Glass, A. G., Cadell, D. M., Rush, B. B., Scott, D. R., Sherman, M. E., Kurman, R. J., Wacholder, S., Stanton, C. K., and Manos, M. M. Epidemiological evidence showing that human papillomavirus infection causes most cervical intraepithelial neoplasia. *J. Natl. Cancer Inst. (Bethesda)*, 85: 958–964, 1993.
- Becker, T. M., Wheeler, C. M., McGough, N. S., Parmenter, C. A., Jordan, S. W., Stidley, C. A., McPherson, S., and Dorin, M. H. Sexually transmitted diseases and other risk factors for cervical dysplasia among southwestern Hispanic and non-Hispanic white women. *JAMA*, 271: 1181–1188, 1994.
- Parkin, D. M., Laara, E., and Muir, C. S. Estimates of the worldwide frequency of sixteen major cancers in 1980. *Int. J. Cancer*, 41: 184–197, 1988.
- Stoler, M. H. A brief synopsis of the role of human papillomaviruses in cervical carcinogenesis. *Am. J. Obstet. Gynecol.*, 175: 1091–1098, 1996.
- Bosch, F. X., Manos, M. M., Muñoz, N., Sherman, M., Jansen, A. M., Peto, J., Schiffman, M. H., Moreno, V., Kurman, R., and Shah, K. V. Prevalence of

- human papillomavirus in cervical cancer: a worldwide perspective. *J. Natl. Cancer Inst. (Bethesda)*, 87: 796–802, 1995.
6. Human Papillomaviruses. IARC Monogr. Eval. Carcinog. Risks Hum. No. 64. Lyon, France: IARC, 1995.
  7. Kiviat, N. Natural history of cervical neoplasia: overview and update. *Am. J. Obstet. Gynecol.*, 175: 1099–1104, 1996.
  8. Odunsi, K., Terry, G., Ho, L., Bell, J., Cuzick, J., and Ganesan, T. S. Susceptibility to human papillomavirus-associated cervical intra-epithelial neoplasia is determined by specific *HLA DR-DQ* alleles. *Int. J. Cancer*, 67: 595–602, 1996.
  9. Allen, M., Kalantari, M., Ylitalo, N., Pettersson, B., Hagmar, B., Scheibenpflug, L., Johansson, B., Pettersson, U., and Gyllensten, U. HLA DQ-DR haplotype and susceptibility to cervical carcinoma: indications of increased risk for development of cervical carcinoma in individuals infected with HPV 18. *Tissue Antigens*, 48: 32–37, 1996.
  10. David, A. L. M., Taylor, G. M., Gokhale, D., Aplin, J. D., Seif, M. W., and Tindall, V. R. HLA-DQB1\*03 and cervical intraepithelial neoplasia type III. *Lancet*, 340: 52, 1992.
  11. Wank, R., and Thomssen, C. High risk of squamous cell carcinoma of the cervix for women with HLA-DQw3. *Nature (Lond.)*, 352: 723–725, 1991.
  12. Mehal, W. Z., Lo, Y. M., Herrington, C. S., Evans, M. F., Papadopoulos, M. C., Odunsi, K., Ganesan, T. S., McGee, J. O. D., Bell, J. I., and Fleming, K. A. Role of human papillomavirus in determining the HLA associated risk of cervical carcinogenesis. *J. Clin. Pathol.*, 47: 1077–1081, 1994.
  13. Nawa, A., Nishiyama, Y., Kobayashi, T., Wakahara, Y., Okamoto, T., Kikkawa, F., Suganuma, N., Goto, S., Kuzuya, K., and Tomoda, Y. Association of human leukocyte antigen-B1\*03 with cervical cancer in Japanese women aged 35 years and younger. *Cancer (Phila.)*, 75: 518–521, 1995.
  14. Helland, Å., Børresen, A.-L., Kristensen, G., and Rønningen, K. S. *DQA1* and *DQB1* genes in patients with squamous cell carcinoma of the cervix: relationship to human papillomavirus infection and prognosis. *Cancer Epidemiol. Biomark. Prev.*, 3: 479–486, 1994.
  15. Montoya, L., Saiz, I., Rey, G., Vela, F., and Clerici-Larradet, N. Cervical carcinoma: human papillomavirus infection and HLA-associated risk factors in the Spanish population. *Eur. J. Immunogenet.*, 25: 329–337, 1998.
  16. Hildesheim, A., Schiffman, M., Scott, D. R., Marti, D., Kissner, T., Sherman, M. E., Glass, A. G., Manos, M. M., Lorincz, A. T., Kurman, R. J., Buckland, J., Rush, B. B., and Carrington, M. Human leukocyte antigen class I/II alleles and development of human papillomavirus-related cervical neoplasia: results from a case-control study conducted in the United States. *Cancer Epidemiol. Biomark. Prev.*, 7: 1035–1041, 1998.
  17. Gregoire, L., Lawrence, W. D., Kukuruga, D., Eisenbrey, A. B., and Lancaster, W. D. Association between *HLA-DQB1* alleles and risk for cervical cancer in African-American women. *Int. J. Cancer*, 57: 504–507, 1994.
  18. Krul, E. J., Schipper, R. F., Schreuder, G. M., Fleuren, G. J., Kenter, G. G., and Melief, C. J. HLA and susceptibility to cervical neoplasia. *Hum. Immunol.*, 60: 337–342, 1999.
  19. Apple, R. J., Erlich, H. A., Klitz, W., Manos, M. M., Becker, T. M., and Wheeler, C. M. HLA DR-DQ associations with cervical carcinoma show papillomavirus-type specificity. *Nat. Genet.*, 6: 157–162, 1994.
  20. Sastre-Garau, X., Loste, M.-N., Vincent-Salomon, A., Favre, M., Mouret, E., de la Rochefordiere, A., Durand, J.-C., Tartout, E., Lepage, V., and Charron, D. Decreased frequency of *HLA-DRB1\*13* alleles in Frenchwomen with HPV-positive carcinoma of the cervix. *Int. J. Cancer*, 69: 159–164, 1996.
  21. Syrjänen, K., Nurmi, T., Mäntyjärvi, R., Ilonen, J., Syrjänen, S., Surcel, H. M., Yliskoski, M., Väyrynen, M., Chang, F., and Saarikoski, S. HLA types in women with cervical human papillomavirus (HPV) lesions prospectively followed up for 10 years. *Cytopathology*, 7: 99–107, 1996.
  22. Silva, B., Vargas-Alarcón, G., Zúñiga-Ramos, J., Rodríguez-Reyna, T. S., Hernández-Martínez, B., Osnaia, N., Kofman, S., Torres-Lobatón, A., and Granados, J. Genetic features of Mexican women predisposing to cancer of the uterine cervix. *Hum. Pathol.*, 30: 626–628, 1999.
  23. Odunsi, K., Terry, G., Ho, L., Bell, J., Cuzick, J., and Ganesan, T. S. Association between *HLA DQB1\*03* and cervical intra-epithelial neoplasia. *Mol. Med.*, 1: 161–171, 1995.
  24. Glew, S. S., Duggan-Keen, M., Ghosh, A. K., Ivanson, A., Sinnott, P., Davidson, J., Dyer, P. A., and Stern, P. L. Lack of association of HLA polymorphisms with human papillomavirus-related cervical cancer. *Hum. Immunol.*, 37: 157–164, 1993.
  25. Wank, R., Meulen, J. T., Luande, J., Eberhardt, H. C., and Pawlita, M. Cervical intraepithelial neoplasia, cervical carcinoma, and risk for patients with *HLA-DQB1\*0602, \*301, \*303* alleles. *Lancet*, 341: 1215, 1993.
  26. Lee, K. W., Johnson, A. H., Tang, T., Yu, W. Y., Karr, R. W., and Hurley, C. K. DRw11 haplotypes: continuum of DRB1 diversity augmented by unique DQ/DRw52 associations. *Hum. Immunol.*, 32: 150–155, 1991.
  27. Koutsky, L. A., Galloway, D. A., and Holmes, K. K. Epidemiology of genital human papillomavirus infection. *Epidemiol. Rev.*, 10: 122–163, 1988.
  28. Hildesheim, A., Gravitt, P., Schiffman, M. H., Kurman, R. J., Barnes, W., Jones, S., Tchabo, J.-G., Brinton, L. A., Copeland, C., Epp, J., and Manos, M. M. Determinants of genital human papillomavirus infection in low-income women in Washington, D.C. *Sex. Transm. Dis.*, 20: 279–285, 1993.
  29. Bauer, H. M., Hildesheim, A., Schiffman, M. H., Glass, A. G., Rush, B. B., Scott, D. R., Cadell, D. M., Kurman, R. J., and Manos, M. M. Determinants of genital human papillomavirus infection in low-risk women in Portland, Oregon. *Sex. Transm. Dis.*, 20: 274–278, 1993.
  30. Wheeler, C. M., Parmenter, C. A., Hunt, W. C., Becker, T. M., Greer, C. E., Hildesheim, A., and Manos, M. M. Determinants of genital human papillomavirus infection among cytologically normal women attending the University of New Mexico Student Health Center. *Sex. Transm. Dis.*, 20: 286–289, 1993.
  31. Melkert, P. W. J., Hopman, E., van den Brule, A. J. C., Risse, E. K. J., van Diest, P. J., Bleker, O. P., Helmerhorst, T., Schipper, M. E. I., Meijer, C. J. L. M., and Walboomers, J. M. M. Prevalence of HPV in cytologically normal cervical smears, as determined by the polymerase chain reaction, is age-dependent. *Int. J. Cancer*, 53: 919–923, 1993.
  32. Zahm, D. M., Nindl, I., Greinke, C., Hoyer, H., and Schneider, A. Colposcopic appearance of cervical intraepithelial neoplasia is age dependent. *Am. J. Obstet. Gynecol.*, 179: 1298–1304, 1998.
  33. Gravitt, P. E., Peyton, C. L., Apple, R. J., and Wheeler, C. M. Genotyping of 27 human papillomavirus types by using L1 consensus PCR products by a single-hybridization, reverse line blot detection method. *J. Clin. Microbiol.*, 36: 3020–3027, 1998.
  34. Erlich, H., Bugawan, T., Begovich, A., and Scharf, S. Analysis of HLA class II polymorphism using polymerase chain reaction. *Arch. Pathol. Lab. Med.*, 117: 482–485, 1993.
  35. Saiki, R. K., Walsh, P. S., Levenson, C. H., and Erlich, H. A. Genetic analysis of amplified DNA with immobilized sequence-specific oligonucleotide probes. *Proc. Natl. Acad. Sci. USA*, 86: 6230–6234, 1989.
  36. Begovich, A. B., McClure, G. R., Suraj, V. C., Helmuth, R. C., Fildes, N., Bugawan, T. L., Erlich, H. A., and Klitz, W. Polymorphism, recombination, and linkage disequilibrium within the HLA class II region. *J. Immunol.*, 148: 249–258, 1992.
  37. Coll Seck, A., Faye, M. A., Critchlow, C. W., Mbaye, A. D., Kuypers, J., Woto-Gaye, G., Langley, C., Benga De, E., Holmes, K. K., and Kiviat, N. B. Cervical intraepithelial neoplasia and human papillomavirus infection among Senegalese women seropositive for HIV-1 or HIV-2 or seronegative for HIV. *Int. J. STD AIDS*, 5: 189–193, 1994.
  38. Ferenczy, A., and Winkler, B. Anatomy and histology of the cervix. In: R. J. Kurman (ed.), *Blaustein's Pathology of the Female Genital Tract*, pp. 141–157. New York: Springer-Verlag New York, Inc., 1987.
  39. Sanjeevi, C. B., Hjelmström, P., Hallmans, G., Wiklund, F., Lenner, P., Ångström, T., Dillner, J., and Lernmark, Å. Different HLA-DR-DQ haplotypes are associated with cervical intraepithelial neoplasia among human papillomavirus type-16 seropositive and seronegative Swedish women. *Int. J. Cancer*, 68: 409–414, 1996.
  40. Neuman, R. J., Huettner, P. C., Li, L., Mardis, E. R., Duffy, B. F., Wilson, R. K., and Rader, J. S. Association between DQB1 and cervical cancer in patients with human papillomavirus and family controls. *Obstet. Gynecol.*, 95: 134–140, 2000.
  41. Duggan-Keen, M. F., Keating, P. J., Stevens, F. R., Sinnott, P., Snijders, P. J., Walboomers, J. M., Davidson, S., Hunter, R. D., Dyer, P. A., and Stern, P. L. Immunogenetic factors in HPV-associated cervical cancer: influence on disease progression. *Eur. J. Immunogenet.*, 23: 275–284, 1996.
  42. Maciag, P. C., Schlecht, N. F., Souza, P. S. A., Franco, E. L., Villa, L. L., and Pétzl-Erler, M. L. Major histocompatibility complex class II polymorphisms and risk of cervical cancer and human papillomavirus infection in Brazilian women. *Cancer Epidemiol. Biomark. Prev.*, 9: 1183–1191, 2000.
  43. Tiercy, J.-M., Sanchez-Mazas, A., Excoffier, L., Shi-Issac, X., Jeannot, M., Mach, B., and Langaney, A. HLA-DR polymorphism in a Senegalese Mandenka population: DNA oligotyping and population genetics of DRB1 specificities. *Am. J. Hum. Genet.*, 51: 592–608, 1992.
  44. Hill, A. V. S., Allsopp, C. E. M., Kwiatkowski, D., Taylor, T. E., Yates, S. N. R., Anstey, N. M., Wirima, J. J., Brewster, D. R., McMichael, A. J., Molyneux, M. E., and Greenwood, B. M. Extensive genetic diversity in the HLA class II region of Africans, with a focally predominant allele, *DRB1\*1304*. *Proc. Natl. Acad. Sci. USA*, 89: 2277–2281, 1992.
  45. Martell, R. W., Arendse, B., Jacobs, M., and Du Toit, E. D. HLA class II specificities and haplotypes in South Africa detected using polymerase chain reaction and sequence-specific oligonucleotide typing. *Tissue Antigens*, 38: 95–98, 1991.
  46. Walboomers, J. M. M., and Meijer, C. J. L. M. Do HPV negative carcinomas exist? *J. Pathol.*, 181: 253–254, 1997.



47. Walboomers, J. M., Jacobs, M. V., Manos, M. M., Bosch, F. X., Kummer, J. A., Shah, K. V., Snijders, P. J., Peto, J., Meijer, C. J., and Muñoz, N. Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *J. Pathol.*, 189: 12–19, 1999.
48. Herrington, C. S. Do HPV-negative cervical carcinomas exist?—revisited. *J. Pathol.*, 189: 1–3, 1999.
49. Brietburd, F., Ramoz, N., Salmon, J., and Orth, G. HLA control in the progression of human papillomavirus infections. *Semin. Cancer Biol.*, 7: 359–371, 1996.
50. Glew, S. S., Connor, M. E., Snijders, P. J., Stanbridge, C. M., Buckley, C. H., Walboomers, J. M., Meijer, C. J., and Stern, P. L. HLA expression in pre-invasive cervical neoplasia in relation to human papilloma virus infection. *Eur. J. Cancer*, 14: 1963–1970, 1993.
51. Hilders, C. G., Munoz, I. M., Nooyen, Y., and Fleuren, G. J. Altered HLA expression by metastatic cervical carcinoma cells as a factor in impaired immune surveillance. *Gynecol. Oncol.*, 57: 366–375, 1995.
52. Hilders, C. G., Houbiers, J. G., Krul, E. J., and Fleuren, G. J. The expression of histocompatibility-related leukocyte antigens in the pathway to cervical carcinoma. *Am. J. Clin. Pathol.*, 101: 5–12, 1994.
53. Glew, S. S., Duggan-Keen, M., Cabrera, T., and Stern, P. L. HLA class II antigen expression in human papillomavirus-associated cervical cancer. *Cancer Res.*, 52: 4009–4016, 1992.
54. Bontkes, H. J., Walboomers, J. M., Meijer, C. J., Helmerhorst, T. J., and Stern, P. L. Specific HLA class I down-regulation is an early event in cervical dysplasia associated with clinical progression (Letter). *Lancet*, 351: 187–188, 1998.
55. Koopman, L. A., Corver, W. E., van der Slik, A. R., Giphart, M. J., and Fleuren, G. J. Multiple genetic alterations cause frequent and heterogeneous human histocompatibility leukocyte antigen class I loss in cervical cancer. *J. Exp. Med.*, 191: 961–976, 2000.
56. Brady, C. S., Bartholomew, J. S., Burt, D. J., Duggan-Keen, M. F., Glenville, S., Telford, N., Little, A-M., Davidson, J. A., Jimenez, P., Ruiz-Cabello, F., Garrido, F., and Stern, P. L., Multiple mechanisms underlie HLA dysregulation in cervical cancer. *Tissue Antigens*, 55: 401–411, 2000.

# Cancer Epidemiology, Biomarkers & Prevention

AACR American Association  
for Cancer Research

## HLA Class II DR-DQ and Increased Risk of Cervical Cancer among Senegalese Women

Patricia Lin, Laura A. Koutsky, Cathy W. Critchlow, et al.

*Cancer Epidemiol Biomarkers Prev* 2001;10:1037-1045.

**Updated version** Access the most recent version of this article at:  
<http://cebp.aacrjournals.org/content/10/10/1037>

**Cited articles** This article cites 54 articles, 10 of which you can access for free at:  
<http://cebp.aacrjournals.org/content/10/10/1037.full#ref-list-1>

**Citing articles** This article has been cited by 4 HighWire-hosted articles. Access the articles at:  
<http://cebp.aacrjournals.org/content/10/10/1037.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://cebp.aacrjournals.org/content/10/10/1037>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.