

Sulfotransferase 1A1 Polymorphism, Endogenous Estrogen Exposure, Well-done Meat Intake, and Breast Cancer Risk¹

Wei Zheng,² Dawen Xie, James R. Cerhan,
Thomas A. Sellers, Wanqing Wen, and Aaron R. Folsom

Vanderbilt-Ingram Cancer Center and Vanderbilt Center for Health Services Research, Vanderbilt University, Nashville, Tennessee 37232-8300 [W. Z., W. W.]; School of Public Health, University of South Carolina, Columbia, South Carolina [D. X.]; Mayo Clinic Cancer Center, Rochester, Minnesota 55905 [J. R. C., T. A. S.]; and School of Public Health, University of Minnesota, Minneapolis, Minnesota 55454 [A. R. F.]

Abstract

Sulfotransferase (SULT) 1A1 is involved in the inactivation of estrogens and bioactivation of heterocyclic amines and polycyclic aromatic hydrocarbons. A G→A transition at codon 213 (CGC/Arg to CAC/His) of the *SULT1A1* gene was reported recently, and individuals homozygous for the His allele have a substantially lower activity of this enzyme than those with other genotypes. We hypothesized that the His allele may be a risk factor for breast cancer, particularly among women who had risk factors related to higher endogenous estrogen exposure. This hypothesis was investigated in a case-control study conducted in a cohort of postmenopausal Iowa women who completed a mailed questionnaire in 1986 on lifestyle factors including information on major breast cancer risk factors. DNA samples and information related to well-done meat intake were obtained from breast cancer cases diagnosed between 1992 and 1994 and a random sample of cancer-free cohort members. Multivariate analysis was performed on data from 156 cases and 332 controls who donated a blood sample. The frequency of the His allele was 41.6% in cases and 34.1% in controls ($P = 0.03$), and the risk of breast cancer was increased with the number of His alleles (P for trend = 0.02). Compared with women with the Arg/Arg genotype, an 80% elevated risk was observed among women homozygous for the His allele (95% confidence interval, 1.0–3.2; $P = 0.04$). This positive association was more pronounced among women who drank alcohol and had a high body mass index, early age at menarche, and late age at menopause, factors related to high endogenous estrogen exposure, than among those who did not have these risk factors. The risk of breast cancer was elevated with increasing doneness level of red meat intake among

women with the Arg/Arg genotype (P for trend, 0.01) or the Arg/His genotype (P for trend, 0.10), whereas this association was not evident for women with the His/His genotype. The results from this study suggest that homozygosity for the *SULT1A1* His²¹³ allele may be a risk factor for breast cancer, and its effect may be modified by the exposure level of endogenous estrogens and heterocyclic amines.

Introduction

Sulfonate conjugation is an important pathway in the metabolism of a variety of endogenous and exogenous compounds, including estrogens and other mammary carcinogens (1–3). This reaction is catalyzed by SULTs,³ a superfamily of multifunctional enzymes including six cytosolic SULTs that have been identified in human tissues (1–3). Some of these enzymes share substantial amino acid sequence identity and overlap considerably in substrate specificity and tissue distribution (1–3). SULT1A1 is one of the most important members in this enzyme family due to its extensive tissue distribution and abundance (1–3). This enzyme has a substantially higher activity than other SULTs in catalyzing the sulfonation of 4-nitrophenol, a commonly used assay in biochemical pharmacogenetic studies for testing the activity of thermostable phenol SULTs (1–3). SULT1A1 catalyzes the sulfonation of estrogens to form water-soluble and biologically inactive estrogen sulfates, reducing the level of estrogen exposure in their target tissues (1, 2). Although sulfonation is, in general, considered a detoxification reaction, several SULTs, particularly SULT1A1, are involved in the bioactivation of certain procarcinogens, including heterocyclic amines and PAHs (3–6), well-documented mammary carcinogens (7, 8). The sulfate esters of these compounds are highly reactive and can bind to DNA to form adducts (3, 6). Therefore, it is possible that SULTs may be related to the risk of breast cancer through both their role in the inactivation of estrogens and bioactivation of environmental mammary carcinogens.

The polymorphic nature of SULTs has long been recognized (1–3, 9). Because of a strong correlation of enzyme activity and thermal stability between SULTs from platelets and those from other organs such as liver, brain, and intestine, platelets were used, for convenience, in many previous biochemical pharmacogenetic studies of human SULTs (1, 2, 9). These studies have demonstrated a large individual variation (about 50-fold in some studies) in the activity of platelet SULTs in humans (1, 2, 9, 10). It was reported recently that a large portion of this variability could be explained by a newly identified common polymorphism (a G→A transition) in the coding

Received 4/12/00; revised 11/9/00; accepted 11/16/00.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported in part by USPHS Grants RO1CA39742 and RO1CA64277 from the NIH and by Supplementary Grant OWH-284 from the National Action Plan on Breast Cancer, Office on Women's Health, USPHS.

² To whom requests for reprints should be addressed, at Vanderbilt University Medical Center, Center for Health Services Research, Medical Center East, 6th Floor, Nashville, TN 37232-8300.

³ The abbreviations used are: SULT, sulfotransferase; CI, confidence interval; OR, odds ratio; PAH, polycyclic aromatic hydrocarbon; BMI, body mass index; WHR, waist:hip ratio.

region (nucleotide 638) of the *SULT1A1* gene (10). This base change in the gene sequence results in an arginine to histidine substitution at codon 213, and individuals homozygous for the *His* allele had only about 15% of the *SULT* activity in platelets of those with other genotypes. The *His* allele is common in the Caucasian population, and the frequency of this allele was reported to be about 30–35% (11, 12). Given the role of *SULT1A1* in the metabolism of estrogens and environmental carcinogens, we hypothesized that the *SULT1A1* polymorphism and its interaction with relevant lifestyle factors may be associated with the risk of breast cancer. We have designed a PCR-based RFLP assay to investigate the association of the *SULT1A1* polymorphism with breast cancer risk in a case-control study conducted in the Iowa Women's Health Study.

Materials and Methods

The Iowa Women's Health Study is a prospective cohort study of 41,836 women, ages 55–69 years, who completed a self-administered questionnaire in January 1986. This cohort of women has been followed for mortality and cancer incidence through computer linkage of study participants with Iowa death certificate files, the National Death Index, and cancer diagnosis data collected by the Iowa State Health Registry, part of the Surveillance, Epidemiology, and End Results Program of the National Cancer Institute (13–16). The baseline questionnaire focused on anthropometric, dietary, and other major risk factors for cancer, including family history of cancer, prior medical conditions, cigarette smoking, reproductive factors, and hormone use. Circumferences of the waist and hip were measured according to a validated protocol. These were used to construct a WHR. Details on the methodology of the cohort study have been published elsewhere (13–16).

From 1995 to 1996, a case-control study was conducted in the Iowa Women's Health Study to collect DNA samples and additional information on meat consumption habits (17–21). Eligible cases for this case-control study were cohort members who were diagnosed with breast cancer between January 1, 1992 and December 31, 1994 ($n = 456$). Controls were randomly selected from women who were cancer free as of January 1, 1992 ($n = 900$). Of the 900 controls, 24 were excluded from the control group because they were later found either to have a breast cancer diagnosis ($n = 3$) or to have been selected to participate in other Iowa Women's Health Study ancillary projects ($n = 21$). All eligible women were asked to complete a self-administered questionnaire on meat consumption habits during the reference year (1991, 1992, or 1993). The reference year for cases was the year immediately before breast cancer diagnosis. Controls were divided randomly into three approximately equal groups for each reference year and answered the same questionnaire. In addition, a series of color photographs used to represent the various doneness levels of hamburger, beefsteak, and bacon was included in the questionnaire to facilitate the assessment of usual doneness levels of meat consumed by the study participants (17). Of all women selected for the study, 273 cases and 657 controls responded, representing approximately 60% and 75% response rates, respectively. The major reasons for nonparticipation were refusal (29.1% of cases and 18.7% of controls), inability to locate subject (4.9% of cases and 3.8% of controls), and subject death before contact (5.7% of cases and 2.5% of controls).

Exfoliated buccal cell samples were collected through the mail as a source of genomic DNA for genotyping assays for this study. About 96.6% of women (267 cases and 631 controls) who completed the supplemental questionnaire provided a sam-

ple of exfoliated buccal cells using the cytobrush method (22). Genomic DNA from buccal cells were extracted using the NaOH method described by Richards *et al.* (22). Blood samples were also collected through the mail from 488 women (156 cases and 332 controls), reflecting an overall response rate of 53% (57% for cases and 50% for controls; Refs. 18–20). Specifically, a blood collection kit including vacutainer tubes, biological specimen packaging containers and envelopes, and instructions was mailed to all women who agreed to donate a blood sample. Study participants were instructed to contact their physicians to have their blood drawn and to return samples via overnight express mail using preaddressed, prepaid envelopes provided by the study. Because of an unsatisfactory quality and quantity of DNA extracted from buccal cells, this study was restricted to those whose blood samples were collected.

Genomic DNA was extracted from peripheral blood leukocytes using a standard protocol and stored at a low temperature for subsequent assays (18–20). Most assays for the *SULT1A1* genotypes were completed in the summer of 1998 using a PCR RFLP-based assay. According to the published sequence of the human *SULT1A1* gene (10, 23), we designed two primers (forward primer, 5'-GGGTCTCTAGGAGAGGTGGC; reverse primer, 5'-GCTGTGGTCCATGAACCTCCT) to amplify a 270-bp fragment of exon 7 that included the polymorphic site (codon 213, *His/CAC* to *Arg/CGC*) of the gene. The PCR reactions were performed on a Perkin-Elmer GeneAmp System 9700 according to the manufacturer's protocol. Specifically, these reactions were carried out in 50 μ l of 20 mM Tris-HCl (pH 8.4), 50 mM KCl, 1.5 mM MgCl₂, 0.2 mM deoxynucleotide triphosphate, and 1 unit of Taq polymerase. The reactions were heated to 94°C for 1 min followed by 35 cycles of 94°C for 30 s, 62°C for 30 s, and 72°C for 30 s. At the end, the reactions were extended 7 min at 72°C. The PCR products (270 bp) were digested with *HhaI* and analyzed by gel electrophoresis [3% (2:1) Nusiev/SeaKem agarose]. Digestion of each PCR product with *HhaI* gives rise to 155- and 115-bp fragments for the *Arg/CGC* allele and a single 270-bp fragment for the *His/CAC* allele. The *SULT1A1* genotypes were successfully identified for all but one case and four controls, resulting in 155 cases and 328 controls for data analysis.

ORs were calculated to measure the strength of the association of breast cancer risk with *SULT1A1* genotypes. Unconditional logistic regression was used to control for potential confounding variables assessed in the baseline survey in 1986 and to derive adjusted ORs and 95% CIs. In addition to the study variables, covariates included in multiple logistic models were age (continuously), WHR (<0.85, \geq 0.85), and live births (<4, \geq 4), variables that were significantly associated with the risk of breast cancer in this study population. Trend tests for dose-response relationships across levels of the exposure variables were performed by treating ordinal score variables (with values of 1, 2, 3, ...) as continuous variables in logistic regression models. Tests for interaction used the likelihood ratio test by adding the interaction term(s) (genotype by exposure) to the model that already included the main effect variables (genotype, exposure) and confounding variables. Reported *P* values are based on two-tailed probability tests.

BMI, age at menarche, age at menopause, and alcohol consumption were evaluated as potential modifiers of *SULT1A1* genotypes because they are related to endogenous estrogen exposure. Although WHR, age at first live birth, and parity are also hormonally related, their association with breast cancer risk is thought to be mainly through mechanisms other than cumulative endogenous estrogen exposure (24, 25). These

Table 1 Comparison of cases and controls by selected demographic and risk factors among postmenopausal Iowa women

	Participants who provided a blood sample			All study participants		
	Cases (n = 156)	Controls (n = 332)	OR (95% CI)	Cases (n = 273)	Controls (n = 657)	OR (95% CI)
Demographic and major risk factors						
Age \geq 65 years at baseline	33	64	1.1 (0.7–1.8)	70	144	1.2 (0.9–1.7)
Education \geq high school	81	153	1.3 (0.9–1.9)	128	277	1.2 (0.9–1.6)
First-degree relatives with breast cancer	26	38	1.5 (0.9–2.7)	47	68	1.8 (1.2–2.7)
WHR \geq 0.85	71	109	1.7 (1.2–2.5)	127	230	1.6 (1.2–2.2)
BMI \geq 26	85	169	1.2 (0.8–1.7)	159	323	1.4 (1.1–1.9)
Menarche at \leq 13 years	116	240	1.1 (0.7–1.7)	203	470	1.2 (0.8–1.6)
Menopause at \geq 55 years	18	31	1.3 (0.7–2.3)	30	66	1.1 (0.7–1.7)
Live birth $<$ 4	106	180	1.8 (1.2–2.7)	180	376	1.4 (1.1–1.9)
First live birth \geq 22 years	91	177	1.2 (0.8–1.8)	155	354	1.1 (0.8–1.5)
Alcohol drinking	66	152	0.9 (0.6–1.3)	118	294	0.9 (0.7–1.2)
Meat doneness level						
Rare/medium	34	122	1.0 (reference)	60	224	1.0 (reference)
Mostly well done	41	73	1.8 (1.1–1.9)	71	155	1.5 (1.03–2.2)
Consistently well/very well done	65	103	2.0 (1.3–3.1)	113	212	1.7 (1.2–2.4)
			<i>P</i> for trend, 0.001			<i>P</i> for trend, $<$ 0.001

variables were also considered in stratified analyses. Exposure to well-done meat was measured by assessing usual doneness levels of hamburgers, beefsteaks, and bacon. Doneness levels of rare or medium, well-done, and very well-done meat were given scores of 1, 2, or 3, respectively, for each food to describe the eating habits of participants on the basis of their responses to the color photographs in the questionnaire. A doneness score, defined as the sum of the usual doneness level for each of these three meats, was then calculated (17). The scores ranged from 3 to 9, with 3 for consistently rare/medium doneness, and 9 for consistently very well-done meat. To enhance the stability of risk estimates, several levels of the doneness score were combined to form the following three groups: (a) rare/medium, scores of 3–4; (b) mostly well-done meat, score of 5; and (c) consistently well- or very well-done meat, scores of 6–9 (17). Information on *SULT1A1* genotype was analyzed along with estrogen-related factors and well-done meat intake to evaluate potential joint effects of genetic and lifestyle factors in the etiology of breast cancer.

Results

Case-control differences in the distribution of selected demographic and major risk factors are shown in Table 1. Information related to well-done meat intake, a relevant environmental exposure for the current analyses, is also presented in Table 1. The risk of breast cancer was found to be positively associated with age, education, family history of breast cancer, WHR, BMI, and reproductive factors, although with this sample size, only the ORs for WHR and number of live births were statistically significant. A clear dose-response relationship between meat doneness level and breast cancer risk was observed. The results were similar to those observed from all study participants, including those who did not provide a blood sample.

Table 2 presents the frequencies of *SULT1A1* alleles and genotypes by case-control status and the association of *SULT1A1* genotypes with breast cancer risk. A substantially higher percentage of cases (41.6%) than controls (34.1%) carried the *His* allele ($P = 0.02$), and more cases (18.7%) than controls (13.4%) were homozygous for this allele. Compared to women with the *Arg/Arg* genotype, the risk of breast cancer was elevated with an increasing number of *His* alleles (P for

Table 2 *SULT1A1* allele and genotype frequencies and adjusted ORs for breast cancer among postmenopausal Iowa women, 1992–1994

<i>SULT1A1</i> (codon 213)	Cases (n = 155)	Controls (n = 328)	Statistics
Allele frequencies (%) ^a			
<i>Arg</i>	58.4	65.9	$P = 0.02$
<i>His</i>	41.6	34.1	
Genotype frequencies (%) ^b			
<i>Arg/Arg</i>	35.5	45.0	$P = 0.09$
<i>Arg/His</i>	45.8	41.3	
<i>His/His</i>	18.7	13.4	
Adjusted ORs and 95% CI			
<i>Arg/Arg</i>	55	148	1.0 (reference) ^c
<i>Arg/His</i>	71	136	1.4 (0.9–2.2)
<i>His/His</i>	29	44	1.8 (1.0–3.2)
			P for trend, 0.027

^a Number of alleles/number of chromosomes.

^b Number of participants with genotype/total number of participants.

^c Adjusted for age, waist:hip ratio, and number of live births.

trend = 0.027), and the adjusted OR for the *His/His* genotype was statistically significant (OR, 1.8; 95% CI, 1.0–3.2; $P = 0.04$).

To evaluate the potential modifying effect of *SULT1A1* genotypes with selected lifestyle factors, stratified analyses were performed (Table 3). With a few exceptions, the risk of breast cancer was elevated for those who carried the *His* allele compared to women with the *Arg/Arg* genotype in virtually all strata defined by the lifestyle factors. The positive association of breast cancer with the *His/His* genotype was stronger among women who had factors reflecting higher endogenous estrogen exposure, such as high BMI, later age at menopause, earlier age at menarche, and regular alcohol consumption, than those who did not have these factors. The trend tests for the gene dose effect were statistically significant only among those who had these risk factors. Although age at first live birth, parity, and WHR are also established risk factors for breast cancer, their association is believed to be mainly through mechanisms (e.g., promoting mammary cell differentiation or insulin resistance) other than cumulative estrogen exposure. The positive

Table 3 Association of *SULT1A1* genotype with breast cancer, stratified by selected lifestyle factors among postmenopausal Iowa women

Variables (by median) ^a	<i>SULT1A1</i> -Arg/Arg		<i>SULT1A1</i> -Arg/His		<i>SULT1A1</i> -His/His		<i>P</i> for trend test
	Cases/controls	OR ^b	Cases/controls	OR (95% CI) ^c	Cases/controls	OR (95% CI) ^c	
BMI							
<26.18 kg/m ²	24/64	1.0	36/72	1.3 (0.7–2.5)	12/28	1.1 (0.5–2.6)	0.60
≥26.18 kg/m ²	31/84	1.0	35/64	1.5 (0.8–2.7)	17/16	2.9 (1.3–6.5)	<0.01
WHR							
<0.82	18/73	1.0	38/69	2.2 (1.2–4.3)	12/21	2.3 (1.0–5.6)	0.02
≥0.82	37/75	1.0	33/67	1.0 (0.6–1.8)	17/23	1.5 (0.7–3.2)	0.37
Age at menopause (yrs)							
<50	24/74	1.0	38/62	2.1 (1.1–3.8)	10/23	1.4 (0.6–3.4)	0.17
≥50	31/74	1.0	33/74	1.1 (0.6–2.0)	19/21	2.2 (1.0–4.7)	0.07
Age at menarche (yrs)							
>13	12/37	1.0	22/44	1.5 (0.6–3.5)	6/11	1.7 (0.5–5.5)	0.32
≤13	43/111	1.0	49/92	1.4 (0.9–2.3)	23/33	1.9 (1.0–3.5)	0.04
Alcohol drinking							
No	36/78	1.0	40/74	1.3 (0.7–2.3)	13/25	1.3 (0.6–2.8)	0.43
Yes	19/70	1.0	31/62	1.8 (0.9–3.6)	16/19	3.4 (1.4–7.9)	<0.01
Age at first live birth (yrs)							
<22	19/73	1.0	33/60	2.1 (1.1–4.2)	12/21	2.2 (0.9–5.3)	0.03
≥22	36/75	1.0	38/76	1.1 (0.6–1.9)	17/23	1.6 (0.7–3.3)	0.30
Parity							
>2	35/103	1.0	44/91	1.5 (0.9–2.5)	15/31	1.5 (0.7–3.0)	0.19
≤2	20/45	1.0	27/45	1.4 (0.7–2.8)	14/13	2.4 (1.0–6.1)	0.07

^a Strata related to high endogenous estrogen exposure are shown in bold.

^b Reference group.

^c Adjusted for age, WHR, and number of live births, except where stratified.

Table 4 Stratified analyses of the association of *SULT1A1* genotype and well-done meat intake with breast cancer risk among postmenopausal Iowa women

Meat doneness level	<i>SULT1A1</i> genotypes			<i>P</i> for trend test
	Arg/Arg	Arg/His	His/His	
A.	No. of cases/controls			
Rare/medium	7/58	17/45	10/17	
Mostly well done	14/29	17/32	10/10	
Consistently well done ^a	25/48	31/44	8/11	
B.	ORs ^b for genotypes			
Rare/medium	1.0 (reference)	3.1 (1.2–8.1)	5.1 (1.7–15.7)	<0.01
Mostly well done	1.0 (reference)	1.1 (0.5–2.6)	2.1 (0.7–6.3)	0.27
Consistently well done ^a	1.0 (reference)	1.6 (0.8–3.2)	1.6 (0.5–4.6)	0.23
	<i>P</i> for interaction, 0.40			
	ORs ^b for meat doneness level			
Rare/medium	1.0 (reference)	1.0 (reference)	1.0 (reference)	
Mostly well done	4.0 (1.4–11.1)	1.4 (0.6–3.1)	1.7 (0.5–6.1)	
Consistently well done ^a	3.6 (1.4–9.3)	1.8 (0.9–3.8)	1.0 (0.3–3.7)	
Trend tests	<i>P</i> = 0.01	<i>P</i> = 0.10	<i>P</i> = 0.98	

^a Including consistently well done or very well done.

^b Adjusted for age, WHR, and number of live births.

association with the *His/His* genotype was stronger in women who had an early age at first live birth, lower parity, or low WHR than it was in those who did not have these factors. Although these data suggested an interactive effect, tests for multiplicative interaction were not statistically significant, perhaps due to the small sample size.

To evaluate potential modifying effects of *SULT1A1* genotype on the association between well-done meat intake and breast cancer risk, cases and controls were tabulated according to the joint distribution of these two factors (Table 4). In the stratified analysis by meat doneness level, a substantially elevated risk of breast cancer was found to be associated with the

His/His genotype (low enzyme activity) only among women who consumed meat rare or medium. The ORs for the association with the *His/His* genotype were 5.1 (95% CI, 1.7–15.7) among women who consumed meats rare/medium and only 1.6 (95% CI, 0.5–4.6) among those who consumed meats consistently well or very well done. The effect of meat doneness level was evaluated in the stratified analysis by *SULT1A1* genotypes. The risk of breast cancer was elevated only among women who carried the *Arg/Arg* genotype (high enzyme activity). These data again suggested an effect modification, although the interaction test based on the multiplicative model was not statistically significant.

Discussion

To our knowledge, no study has been published thus far on the association between the *SULT1A1* polymorphism and breast cancer risk. The data from this study suggest that the *R213H* polymorphism of the *SULT1A1* gene may influence the risk of breast cancer. Homozygosity for the *His* allele was associated with an 80% elevated risk of breast cancer, and the risk was further elevated among women who had some risk factors related to higher endogenous estrogen exposure. We also found that the *SULT1A1* polymorphism may modify the association between well-done meat intake and breast cancer risk; the positive association with usual doneness level of meat intake was more evident among women who carried the *Arg* allele than those who carried the *His* allele. Our findings are new and consistent with the role of *SULT1A1* in the metabolism of estrogens and environmental carcinogens, such as heterocyclic amines and PAH (1–6), well-documented mammary carcinogens in animal models (4, 7, 8).

Our finding of a positive association between the *His* allele (encoding a low activity allozyme) and breast cancer risk is consistent with the role of *SULT1A1* in the inactivation of estrogens, the hormones believed to play a central role in the etiology of breast cancer (24, 25). This finding was further supported by the observation that the positive association with the *His* allele was more evident among women who had risk factors related to higher endogenous estrogen exposure. It has been reported that the level of endogenous estrogen is positively related to age at menopause and inversely related to age at menarche (24, 25). Alcohol drinking also increases blood estrogen levels (24–26). After menopause, adipose tissue is the major source of estrogen production from androgens (24, 25, 27), and high body weight, often measured using BMI, has been demonstrated to be a risk factor for postmenopausal breast cancer (19, 23). It was intriguing to find that the association of breast cancer with the *His/His* genotype was stronger among women who had a high BMI, earlier age at menarche, later age at menopause, and a regular alcohol consumption history than it was among those who did not have these factors.

The strong association of breast cancer risk with the *His* allele observed in this study among women with a low WHR is puzzling because a high WHR is a risk factor for breast cancer. In a cross-sectional study of 88 participants of the Iowa Women's Health Study, we found that WHR was positively correlated with blood level of insulin and inversely related to the blood level of sex hormone-binding globulin (28). However, WHR was not correlated with blood estrogens ($r = -0.16$ for estrone; $r = 0.13$ for estradiol), even free estradiol ($r = 0.14$). Recently, Tchernof *et al.* (29) reviewed published data from previous epidemiological studies and concluded that there is an increase in central body fatness during the transition period of menopause, and this increase may be due to estrogen deficiency as a result of a progressive loss of ovarian function in estrogen production. Intervention studies have shown that estrogen replacement therapy reduces central fat distribution in postmenopausal women (29). The association of breast cancer with the *His/His* genotype also appears to be stronger among women who had an earlier age at first live birth and lower parity than it is among those who did not have these factors. It appears that this observation cannot simply be explained by the cumulative estrogen exposure hypothesis. During a pregnancy, particularly during the first pregnancy, breast cells proliferate rapidly and differentiate into mature cells prepared for lactation (24, 25). Mature breast cells have a longer cell cycle and spend more time in G_1 , the phase that allows for DNA repair, and are thus

less susceptible to carcinogens (24). Early age at first live birth is believed to reduce the susceptibility of mammary tissue at an early stage of a women's life, and additional live births may confer further benefits by increasing mature cells (and thus decreasing susceptible cells) of the breast.

Burned meat contains a wide variety of potentially carcinogenic compounds (4, 7, 8, 30), including heterocyclic amines and PAH procarcinogens that can be activated by *SULT1A1* and other *SULT* variants (2–4). Despite some inconsistent findings (31), high intakes of fried foods and a preference for well-done meats have been shown to be associated with an increased risk of breast cancer in several epidemiological studies (17, 21, 32–37), including a prospective study reported recently (37). We recently reported a dose-response relationship between doneness levels of meat consumption and the risk of breast cancer, with a nearly 5-fold elevated risk observed among women who consistently consumed very well-done meat (17). We showed in this report that the positive association was evident only among women with the *Arg/Arg* or *Arg/His* genotype and not among those with the *His/His* genotype. This finding is biologically plausible because the former two genotypes are associated with higher phenol *SULT* activity and hence greater bioactivation of procarcinogens than the latter genotype.

A consideration in this study may be the low response rate for blood sample collection. The survival rate for breast cancer, however, is very high in our study population. As such, only 5.7% of patients died before we contacted them for the supplemental survey and sample collection. Therefore, the potential influence of selective survival due to genotypes is unlikely to be substantial in our study. The response rate for buccal cell collection for studying genetic factors was reasonably high, indicating that the low response in blood collection is unlikely to be related to the genetic characteristics of study subjects. Women who donated a blood sample to the study were not atypical because virtually all established risk factors for breast cancer were found to be positively associated with the risk of breast cancer in our study, although some of the associations were not statistically significant due to the small sample size. Data on factors related to endogenous estrogen exposure were collected before cancer diagnosis. The prospective nature of the study design for these factors eliminates potential recall bias. The information on well-done meat intake was collected in this case-control study retrospectively. An individual's intake preference for meat doneness level, however, is likely to be related to a personal habit, which may be recalled with relatively high accuracy. At the time this study was conducted, the hypothesis for a potential link between well-done meat intake and breast cancer risk was relatively new. Therefore, there is no reason to speculate that the breast cancer patients would differentially recall intake of well-done meat as compared with controls, particularly because consumption of well-done hamburger is recommended in the news media to reduce the risk of *Escherichia coli* infection.

In summary, this case-control study found that postmenopausal women who were homozygous for the *His* allele at codon 213 of the *SULT1A1* gene may be at an increased risk of breast cancer, particularly if they have risk factors related to higher endogenous estrogen exposure. Our data also suggest a potentially modifying effect of the *SULT1A1* polymorphism on the association of well-done meat intake and breast cancer. The dual effect of *SULT1A1* in the inactivation of estrogens and activation of environmental carcinogens complicates the association between *SULT1A1* genotype and breast cancer risk. Because endogenous estrogen exposure is believed to play a

more important role in the etiology of breast cancer than heterocyclic amine and PAH exposure, it is conceivable that low activity of *SULT1A1* (the *His* allele) may be a risk factor for breast cancer for most women. The high frequency of this risk allele and the involvement of this enzyme in estrogen, heterocyclic amine, and PAH metabolism suggest that *SULT1A1* polymorphism may be important in the etiology of breast cancer. The sample size of this study is small, and additional studies will be needed to replicate the findings.

Acknowledgments

We thank Drs. Deborah R. Gustafson, Lawrence Kushi, and Kristin Anderson for their contributions to the Iowa Women's Health Study, Dr. Zhong Lin Deng for performing part of the genotyping assays, Dr. Rashmi Sinha for providing copies of the food photographs for assessing meat doneness levels, and Dr. Richard Weinshilboum for helpful discussion of this project.

References

- Weinshilboum, R. M., Otterness, D. M., Aksoy, I. A., Wood, T. C., Her, C., and Raftogianis, R. B. Sulfation and sulfotransferases. I. Sulfotransferase molecular biology: cDNAs and genes. *FASEB J.*, *11*: 3–14, 1997.
- Falany, C. N. Enzymology of human cytosolic sulfotransferases. *FASEB J.*, *11*: 206–216, 1997.
- Glatt, H. R. Sulfation and sulfotransferases. IV. Bioactivation of mutagens via sulfation. *FASEB J.*, *11*: 314–321, 1997.
- Lewis, A. J., Walle, U. K., King, R. S., Kadlubar, F. F., Falany, C. N., and Walle, T. Bioactivation of the cooked food mutagen *N*-hydroxy-2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine by estrogen sulfotransferase in cultured human mammary epithelial cells. *Carcinogenesis (Lond.)*, *19*: 2049–2053, 1998.
- Nowell, S. A., Ozawa, S., Ambrosone, C. B., MacLeod, S. L., Kadlubar, F. F., and Lang, N. P. Relationship of *SULT1A1* genotype to sulfotransferase activity phenotype in platelet cytosol: effect of genotype on heterocyclic amine activation. *Proc. Am. Assoc. Cancer Res.*, *41*: 551, 2000.
- Miller, J. A. Sulfonation in chemical carcinogenesis: history and present status. *Chem. Biol. Interact.*, *92*: 329–341, 1994.
- Nagao, M., Ushijima, T., Wakabayashi, K., Ochiai, M., Kushida, H., Sugimura, T., Hasegawa, R., Shirai, T., and Ito, N. Dietary carcinogens and mammary carcinogenesis: induction of rat mammary carcinomas by administration of heterocyclic amines in cooked foods. *Cancer (Phila.)*, *74* (Suppl.): 1063–1069, 1994.
- Adamson, R. H. Mutagens and carcinogens formed during cooking of foods and methods to minimize their formation. In: V. T. DeVita Jr., S. Hellman, and S. A. Rosenberg (eds.), *Cancer Prevention*, pp. 1–7. Philadelphia: J. B. Lippincott, 1990.
- Price, R. A., Spielman, R. S., Lucena, A. L., Van Loon, J. A., Maidak, B. L., and Weinshilboum, R. M. Genetic polymorphism for human platelet thermostable phenol sulfotransferase (TS PST) activity. *Genetics*, *122*: 905–914, 1989.
- Raftogianis, R. B., Wood, T. C., Otterness, D. M., Van Loon, J. A., and Weinshilboum, R. M. Phenol sulfotransferase pharmacogenetics in humans: association of common *SULT1A1* alleles with TS PST phenotype. *Biochem. Biophys. Res. Commun.*, *239*: 298–304, 1997.
- Coughtrie, M. W. H., Gilissen, R. A. H. J., Shek, B., Strange, R. C., Fryer, A. A., Jones, P. W., and Bamber, D. E. Phenol sulphotransferase *SULT1A1* polymorphism: molecular diagnosis and allele frequencies in Caucasian and African populations. *Biochem. J.*, *337*: 45–49, 1999.
- Ozawa, S., Tang, Y.-M., Yamazoe, Y., Kato, R., Lang, N. P., and Kadlubar, F. F. Genetic polymorphisms in human liver phenol sulfotransferases involved in the bioactivation of *N*-hydroxy derivatives of carcinogenic arylamines and heterocyclic amines. *Chem. Biol. Interact.*, *109*: 237–248, 1998.
- Folsom, A. R., Kaye, S. A., Sellers, T. A., Hong, C., Cerhan, J. R., Potter, J. D., and Prineas, R. J. Body fat distribution and 5-year risk of death in older women. *J. Am. Med. Assoc.*, *269*: 483–487, 1993.
- Bisgard, K. M., Folsom, A. R., Hong, C., and Sellers, T. A. Mortality and cancer rates in non-respondents to a prospective study of older women: 5-year follow-up. *Am. J. Epidemiol.*, *139*: 990–1000, 1994.
- Sellers, T. A., Kushi, L. H., Potter, J. D., Kaye, S. A., Nelson, C. L., McGovern, P. G., and Folsom, A. R. Effect of family history, body-fat distribution, and reproductive factors on the risk of post-menopausal breast cancer. *N. Engl. J. Med.*, *326*: 1323–1329, 1992.
- Kushi, L. H., Sellers, T. A., Potter, J. D., Nelson, C. L., Munger, R. G., Kaye, S. A., and Folsom, A. R. Dietary fat and post-menopausal breast cancer. *J. Natl. Cancer Inst. (Bethesda)*, *84*: 1092–1099, 1992.
- Zheng, W., Gustafson, D. R., Sinha, R., Cerhan, J. R., Moore, D., Hong, C., Anderson, K. E., Kushi, L. H., Sellers, T. A., and Folsom, A. R. Well-done meat intake and the risk of breast cancer. *J. Natl. Cancer Inst. (Bethesda)*, *90*: 1724–1729, 1998.
- Zheng, W., Deitz, A. C., Campbell, D. R., Wen, W. Q., Cerhan, J. R., Sellers, T. A., Folsom, A. R., and Hein, D. W. *N*-Acetyltransferase 1 (NAT1) genetic polymorphism, cigarette smoking, well-done meat intake, and breast cancer risk. *Cancer Epidemiol. Biomark. Prev.*, *8*: 233–239, 1999.
- Deitz, A. C., Zheng, W., Leff, M. A., Gross, M., Wen, W. Q., Doll, M. A., Xiao, G. H., Folsom, A. R., and Hein, D. W. *N*-Acetyltransferase-2 (NAT2) acetylation polymorphism, well-done meat intake, and breast cancer risk among postmenopausal women. *Cancer Epidemiol. Biomark. Prev.*, *9*: 905–910, 2000.
- Zheng, W., Xie, D., Heimeil, A. Y., Wen, W. Q., Cerhan, J. R., Gross, M., Gustafson, D. R., and Folsom, A. R. Breast cancer and *GSTM1* and *GSTT1* polymorphisms: evidence for potential gene-environment interaction. In: *The 2000 Keystone Symposia—Molecular Epidemiology: A New Tool in Cancer Prevention*, p. 47. Taos, NM: Keystone Symposia, 2000.
- Sinha, R., Gustafson, D. R., Kulldorff, M., Wen, W. Q., Cerhan, J. R., and Zheng, W. 2-Amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine, a carcinogen in high-temperature-cooked meat, and breast cancer risk. *J. Natl. Cancer Inst. (Bethesda)*, *92*: 1352–1354, 2000.
- Richards, B., Skoletsky, J., Shuber, A. P., Balfour, R., Stern, R. C., Dorkin, H. L., Parad, R. B., Witt, D., and Klinger, K. W. Multiplex PCR amplification from *CFTR* gene using DNA prepared from buccal brushes/swabs. *Hum. Mol. Genet.*, *2*: 159–163, 1993.
- Raftogianis, R. B., Her, C., and Weinshilboum, R. M. Human phenol sulfotransferase pharmacogenetics: *STP1* gene cloning and structural characterization. *Pharmacogenetics*, *6*: 473–487, 1996.
- Willett, W. C., Rockhill, B., Hankinson, S. E., Hunter, D. J., and Colditz, G. A. Epidemiology and nongenetic causes of breast cancer. In: J. R. Harris, M. E. Lippman, M. Morrow, and C. K. Osborne (eds.), *Diseases of the Breast*, pp. 175–220. Philadelphia: Lippincott Williams & Wilkins, 1999.
- Henderson, B. E., Pike, M. C., Bernstein, L., and Ross, R. K. Breast cancer. In: D. Schottenfeld and J. F. Fraumeni Jr. (eds.), *Cancer Epidemiology and Prevention*, 2nd ed., pp. 1022–1039. New York: Oxford University Press, 1996.
- Schatzkin, A., and Longnecker, M. P. Alcohol and breast cancer. *Cancer (Phila.)*, *74*: 1101–1110, 1994.
- Ballard-Barbash, R. Anthropometry and breast cancer. *Cancer (Phila.)*, *74*: 1090–1100, 1994.
- Kaye, S. A., Folsom, A. R., Soler, J. T., Prineas, R. J., and Potter, J. D. Associations of body mass and fat distribution with sex hormone concentrations in postmenopausal women. *Int. J. Epidemiol.*, *20*: 151–156, 1991.
- Tchernof, A., Poehlman, E. T., and Despres, J. P. Body fat distribution, the menopause transition, and hormone replacement therapy. *Diabetes & Metab.*, *26*: 12–20, 2000.
- El-Bayoumy, K. Environmental carcinogens that may be involved in human breast cancer etiology. *Chem. Res. Toxicol.*, *5*: 585–590, 1992.
- Gertig, D. M., Hankinson, S. E., Hough, H., Spiegelman, D., Colditz, G. A., Willett, W. C., Kelsey, K. T., and Hunter, D. J. *N*-Acetyltransferase-2 genotypes, meat intake and breast cancer risk. *Int. J. Cancer*, *80*: 13–17, 1999.
- Knekt, P., Steineck, G., Jrvinen, R., and Hakulinen, A. A. Intake of fried meat and risk of cancer: a follow-up study in Finland. *Int. J. Cancer*, *59*: 756–760, 1994.
- Phillips, R. L. Role of lifestyle and dietary habits in risk of cancer among Seventh-Day Adventists. *Cancer Res.*, *35*: 3513–3522, 1975.
- Iscovich, J. M., Iscovich, R. B., Howe, G., Shiboski, S., and Kaldor, J. M. A case-control study of diet and breast cancer in Argentina. *Int. J. Cancer*, *44*: 770–776, 1989.
- Matos, E. L., Thomas, D. B., Sobel, N., and Vuoto, D. Breast cancer in Argentina: case-control study with special reference to meat eating habits. *Neoplasma (Bratisl.)*, *38*: 357–366, 1991.
- De Stefani, E., Ronco, A., Mendilaharsu, M., Guidobono, M., and Deneo-Pellegrini, H. Meat intake, heterocyclic amines, and risk of breast cancer: a case-control study in Uruguay. *Cancer Epidemiol. Biomark. Prev.*, *6*: 573–581, 1997.
- Visvanathan, K., Strickland, P., Bell, D. A., Watson, M. A., Rothman, N., Hoffman, S., and Helzlsouer, K. J. Association of NAT2, *GSTM1*, *GSTP1*, *GSTT1*, flame-broiled food and the risk of breast cancer: a nested case-control study. *Proc. Am. Assoc. Cancer Res.*, *41*: 804, 2000.

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Sulfotransferase 1A1 Polymorphism, Endogenous Estrogen Exposure, Well-done Meat Intake, and Breast Cancer Risk

Wei Zheng, Dawen Xie, James R. Cerhan, et al.

Cancer Epidemiol Biomarkers Prev 2001;10:89-94.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/10/2/89>

Cited articles This article cites 32 articles, 5 of which you can access for free at:
<http://cebp.aacrjournals.org/content/10/2/89.full#ref-list-1>

Citing articles This article has been cited by 8 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/10/2/89.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/10/2/89>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.