

Serum Oxidized Low-Density Lipoprotein Levels and Risk of Colorectal Cancer: A Case-Control Study Nested in the Japan Collaborative Cohort Study

Koji Suzuki,¹ Yoshinori Ito,¹ Kenji Wakai,^{3,4} Miyuki Kawado,² Shuji Hashimoto,² Hideaki Toyoshima,⁵ Masayo Kojima,⁶ Shinkan Tokudome,⁶ Norihiko Hayakawa,⁷ Yoshiyuki Watanabe,⁸ Koji Tamakoshi,⁵ Sadao Suzuki,⁶ Kotaro Ozasa,⁸ Akiko Tamakoshi,⁴ and Japan Collaborative Cohort Study Group

¹Department of Public Health, Fujita Health University School of Health Sciences; ²Department of Hygiene, Fujita Health University School of Medicine, Toyoake, Japan; ³Division of Epidemiology and Prevention, Aichi Cancer Center Research Institute; Departments of ⁴Preventive Medicine/Biostatistics and Medical Decision Making and ⁵Public Health/Health Information Dynamics, Nagoya University Graduate School of Medicine; ⁶Department of Health Promotion and Preventive Medicine, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan; ⁷Department of Epidemiology, Research Institute for Radiation Biology and Medicine, Hiroshima University, Hiroshima, Japan; and ⁸Department of Epidemiology for Community Health and Medicine, Kyoto Prefectural University of Medicine Graduate School of Medical Science, Kyoto, Japan

Abstract

Oxidative stress plays an important role in carcinogenesis, but few epidemiologic studies have examined associations with risk of colorectal cancer. Relationships between serum levels of oxidized low-density lipoprotein (oxLDL) and oxLDL antibody (oLAB) and colorectal cancer risk were investigated in a case-control study nested in the Japan Collaborative Cohort

Study for Evaluation of Cancer Risk. Serum samples and lifestyle information were collected at baseline from 39,242 men and women between 1988 and 1990. Of these, 161 incidents and deaths from colorectal cancer were identified through 1999, and 395 controls were matched for gender, age, and study area. Measurements were taken of serum oxLDL levels in 119 cases and 316 controls and serum oLAB levels in 153 cases and 376 controls. Odds ratios (95% confidence intervals) across quartiles, adjusted for confounding factors, were 1.55 (0.70-3.46), 1.90 (0.84-4.28), and 3.65 (1.50-8.92) for oxLDL ($P_{\text{trend}} = 0.004$) and 0.98 (0.54-1.80), 0.75 (0.39-1.48), and 1.68 (0.90-3.13) for oLAB ($P_{\text{trend}} = 0.140$). Further adjustment for serum total cholesterol and α -tocopherol did not materially change these associations. Odds ratio (95% confidence interval) of the highest quartile of serum oxLDL compared with the lowest quartile was 3.40 (1.09-10.58; $P_{\text{trend}} = 0.045$). Analyses restricted to colon cancer cases and corresponding controls yielded similar relationships between serum oxLDL and oLAB levels and risk. In conclusion, higher levels of serum oxLDL may increase risk of colorectal cancer. (Cancer Epidemiol Biomarkers Prev 2004;13(11):1781-7)

Received 2/23/04; revised 5/24/04; accepted 5/28/04.

Grant support: Ministry of Education, Culture, Sports, Science, and Technology of Japan Scientific Research on Priority Areas grant-in-aid 14031222 and Scientific Research grants-in-aid 61010076, 62010074, 63010074, 1010068, 2151065, 3151064, 4151063, 5151069, 6279102, and 11181101.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Note: The present members of the JACC Study and their affiliations are as follows: A. Tamakoshi (Nagoya University Graduate School of Medicine; present Chairman of the JACC Study Group), M. Mori (Sapporo Medical University School of Medicine), Y. Motohashi (Akita University School of Medicine), I. Tsuji (Tohoku University Graduate School of Medicine), Y. Nakamura (Jichi Medical School), H. Iso (Institute of Community Medicine, University of Tsukuba), H. Mikami (Chiba Cancer Center), Y. Inaba (Juntendo University School of Medicine), Y. Hoshiyama (Showa University School of Medicine), H. Suzuki (Niigata University Graduate School of Medical and Dental Sciences), H. Shimizu (Gifu University School of Medicine), H. Toyoshima (Nagoya University Graduate School of Medicine), S. Tokudome (Nagoya City University Graduate School of Medical Sciences), Y. Ito (Fujita Health University School of Health Sciences), S. Hashimoto (Fujita Health University School of Medicine), S. Kikuchi (Aichi Medical University School of Medicine), A. Koizumi (Graduate School of Medicine and Faculty of Medicine, Kyoto University), T. Kawamura (Kyoto University Center for Student Health), Y. Watanabe and T. Miki (Kyoto Prefectural University of Medicine Graduate School of Medical Science), C. Date (Faculty of Human Environmental Sciences, Mukogawa Women's University), K. Sakata (Wakayama Medical University), T. Nose (Tottori University Faculty of Medicine), N. Hayakawa (Research Institute for Radiation Biology and Medicine, Hiroshima University), T. Yoshimura (Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan), K. Fukuda (Kurume University School of Medicine), N. Okamoto (Kanagawa Cancer Center), H. Shio (Moriyama Municipal Hospital), Y. Ohno (Asahi Rosai Hospital; former Chairman of the JACC Study Group), T. Kitagawa (Cancer Institute of the Japanese Foundation for Cancer Research), T. Kuroki (Gifu University), and K. Tajima (Aichi Cancer Center Research Institute). Past members of the JACC Study are listed in ref. 7 along with their affiliations at the time of participation, with the exception of the following: T. Shimamoto (Institute of Community Medicine, University of Tsukuba), H. Tanaka (Medical Research Institute, Tokyo Medical and Dental University), S. Hisamichi (Tohoku University Graduate School of Medicine), M. Nakao (Kyoto Prefectural University of Medicine), T. Suzuki (Research Institute, Osaka Medical Center for Cancer and Cardiovascular Diseases), T. Hashimoto (Wakayama Medical University), and T. Ishibashi (Asama General Hospital).

Requests for reprints: Koji Suzuki, Department of Public Health, Fujita Health University School of Health Sciences, 1-98 Dengakugakubo, Kutsukake-cho, Toyoake, Aichi 470-1192, Japan. Phone: 81-562-93-9424; Fax: 81-562-93-4595. E-mail: ksuzuki@fujita-hu.ac.jp

Copyright © 2004 American Association for Cancer Research.

Introduction

Reactive oxygen species (ROS) cause oxidation of lipids, proteins, and DNA *in vivo* (1, 2), and free radical and lipid peroxides have been considered very important in carcinogenesis (3). Some studies have reported high lipid peroxidation in human colorectal cancer tissue (4, 5). However, few epidemiologic studies have investigated relationships between lipid peroxidation and colorectal cancer.

Oxidized low-density lipoprotein (oxLDL) is generated by the actions of ROS *in vivo*. The oxLDL is taken up by macrophages, which develop into foam cells, and oxLDL antibody (oLAB) is present in both atherosclerotic lesions and plasma (6). Thus, oxLDL is believed to play a critical role in the development and progression of

atherosclerosis (7). Serum oxLDL levels may be considered as a biomarker reflecting the state of oxidative stress and lipid metabolism *in vivo*. Experimental studies have indicated that oxLDL increases intracellular levels of ROS and lipid peroxidation products (thiobarbituric acid reactive substances; ref. 8). The oLAB plays a positive role in maintaining low levels of serum oxLDL.

Various lifestyle factors such as physical activity and diets reportedly affect oxLDL (9-14). Regular physical activity has been found to increase LDL resistance to oxidation and decrease plasma oxLDL concentration (9). Another study identified correlations between weight reduction and decreased oxLDL (10). Some epidemiologic studies have reported that physical activity (15-17) displays significant inverse associations with colorectal cancer and that obesity (18, 19) is associated with increased risk of colorectal cancer.

Vitamin E and lycopene have been shown to display powerful antioxidant properties, reducing LDL oxidation and oxidative damage to plasma proteins (11). Supplementation with antioxidant nutrients (vitamin E, vitamin C, and carotenoids) has been shown to protect LDL from oxidation (12-14). High dietary carotenoid intake possibly decreases the risk of colorectal cancer (20) and a meta-analysis (21) of five prospective nested case-control studies indicated that high plasma levels of α -tocopherol were associated with a modest decrease in the subsequent incidence of colorectal cancer.

Given the results of these previous studies, we hypothesize that serum oxLDL levels represent a biomarker reflecting oxidative stress and lifestyle factors such as physical activity and diet as related to colorectal cancer.

To the best of our knowledge, no studies have identified relationship between oxLDL and risk of colorectal cancer. We therefore examined correlations between serum levels of oxLDL and oLAB and risk of colorectal cancer in a case-control study nested in a large-scale Japanese cohort.

Materials and Methods

Study Subjects and Serum Samples. Study subjects were recruited in the Japan Collaborative Cohort (JACC) Study for Evaluation of Cancer Risk sponsored by Monbukagakusyo (Ministry of Education, Culture, Sports, Science, and Technology of Japan; ref. 22). This study involves 110,792 residents who were ages 40 to 79 years at baseline from 45 areas all over Japan. An epidemiologic survey of lifestyle factors was conducted using a self-administered questionnaire about health conditions and lifestyles such as medical history, smoking habits, and alcohol consumption. Details of this study have been published elsewhere (22).

In addition to the questionnaire survey, participants in the JACC Study provided peripheral blood samples at health screening checkups sponsored by municipalities between 1988 and 1990. A total of 39,242 subjects (35.4% of respondents to the questionnaire survey) provided blood samples. Sera were separated from samples at laboratories in or near the surveyed municipalities as soon as possible after sampling. Serum derived from each subject was divided into three to five tubes (100-500 μ L/tube) and stored at -80°C until analyzed.

Written informed consent for participation was obtained individually from subjects, with the exception of those in a few study areas in which informed consent was provided at the group level after the aim of the study and confidentiality of the data had been explained to community leaders. This study was approved by the Ethical Committee of Medical Care and Research at Fujita Health University.

Case Ascertainment and Control Selection. Subjects who died or moved away from study areas were identified using population registries, and causes of death were confirmed from death certificates. Incident cases of cancer could be identified by linkage with cancer registries in 24 of the 45 study areas. Follow-up for death was conducted from baseline to the end of 1999, and follow-up for incidence was conducted from baseline to the end of 1997, excluding three study areas (from baseline to the end of 1994, 1995, and 1996, respectively). Only 4% of subjects were lost to follow-up due to moving during the study period.

Death and incidence of colorectal cancer were defined by the codes "C18," "C19," and "C20" in the *International Statistical Classification of Diseases and Related Health Problem, 10th Revision* (23). During follow-up, 76 deaths from colorectal cancer [colon (C18), $n = 50$; rectum (C19 and 20), $n = 26$] and 185 incident cases of colorectal cancer (colon, $n = 123$; rectum, $n = 62$) were identified from subjects who had provided serum samples at baseline. Of these, 23 subjects with a history of colorectal and other cancers at baseline were excluded. For each case of colorectal cancer, two or three controls were selected from the remaining population without incident cancer or previous history of cancer, matching for gender, age (± 3 years), and study area. A total of 49 cases and 56 controls without sufficient samples for measurement of serum levels of both oxLDL and oLAB were excluded from analysis. Following these exclusions, subjects without corresponding cases or controls were also excluded. Finally, serum levels of either oxLDL or oLAB could be measured in 161 cases (111 colon cancer cases and 50 rectum cancer cases) and 395 controls in this study. Of these, sufficient serum samples for determination of oxLDL and oLAB were available for 103 cases and 279 controls and 135 cases and 330 controls, respectively. For analyses using only incident cases and corresponding controls, the subjects were 82 cases and 216 controls for oxLDL and 111 cases and 266 controls for oLAB, respectively. Incident and dead cases were analyzed together to maximize sample size for main analysis.

Biochemical Analyses of Sera. All samples were analyzed by trained staff blinded to case-control status in 2001. Serum oxLDL and oLAB were determined by enzyme-linked immunoassay using commercially available kits (oxLDL: Oxidized LDL ELISA kit, Mercodia, Uppsala, Sweden; oLAB: oLAB ELISA kit, Biomedica, Vienna, Austria) in our laboratory. With regard to intraassay and interassay reproducibility, coefficients of variation for oxLDL (24) and oLAB (25) were $<10\%$. Serum α -tocopherol levels were measured separately using high-performance liquid chromatography (26) in our laboratory. Serum total cholesterol was measured using an autoanalyzer at a single laboratory (SRL,

Hachioji, Japan). Values for oxLDL and oLAB could not be measured in all serum samples, because some initial samples yielded insufficient sera and other various substances were also measured from the same samples.

Serum samples of subjects had been stored for ~10 years until assay. Distribution of mean \pm SD values for serum oxLDL levels in study controls [males: 36.1 ± 11.1 units/L ($n = 144$); females: 39.0 ± 11.4 units/L ($n = 172$)] was similar to that in our previous study (24) using fresh sera [males: 41.6 ± 12.2 units/L ($n = 158$); females: 42.7 ± 13.9 units/L ($n = 158$)]. Distributions of serum oLAB were also similar, with median values (25th-75th percentiles) at 191.0 (128.0-241.0) units/L in males ($n = 179$) and 192.0 (142.0-304.0) units/L in females ($n = 197$) for the present study compared with 170.7 (130.9-301.2) units/L in males ($n = 158$) and 209.0 (152.6-312.5) units/L in females ($n = 158$) for the previous study (25). Subjects in this and our previous study were Japanese ages 40 to 79 years, and the same ELISA kits were used. Serum levels of oxLDL and oLAB had thus not changed substantially during long-term storage.

Statistical Analyses. Body mass index (BMI) was calculated as body weight (kg) divided by height (m) squared. Baseline characteristics were compared between cases and controls using χ^2 tests. Mean differences for serum total cholesterol levels and BMI between cases and controls were examined using t tests. Because serum oxLDL, oLAB, and α -tocopherol levels are log normally distributed (25, 26), mean differences between cases and controls were examined using t tests after converting serum levels of oxLDL, oLAB, and α -tocopherol to logarithmic values. Relationships among serum levels of oxLDL, oLAB, total cholesterol, and α -tocopherol were examined using Spearman correlation coefficients. α -Tocopherol was included in this analysis because it binds to LDL and may be associated with decreased risk of colorectal cancer (21).

Conditional logistic regression models with gender, age, and study area strata were applied to calculate odds ratios (OR) and 95% confidence intervals (95% CI) for colorectal cancer. ORs were computed according to quartile levels of serum oxLDL and oLAB. Cases were categorized into four groups according to the quartile in controls for serum oxLDL and oLAB. To test for linear trends in ORs over quartiles, each quartile was coded as 0, 1, 2, or 3 and then incorporated into logistic models as a single variable.

Potential confounding was considered by smoking habits (never, former, or current smokers and unknown), drinking habits (never, former, or current drinkers and unknown), intake frequency of green leafy vegetables (1-2 times/mo or less, 1-2 times/wk or more, and unknown), time spent in sports or physical exercise (little, 1 h/wk or more, and unknown), family history of colorectal cancer (yes, no, and unknown), and BMI (<20.0 , 20.0 - 24.9 , or ≥ 25.0 kg/m² and unknown). Moreover, ORs for colorectal cancer by serum levels of oxLDL and oLAB were also computed after adjustment for the above confounding factors and quartiles of serum total cholesterol and α -tocopherol, because LDL binds to cholesterol and α -tocopherol. Elevated serum cholesterol levels are linked with increased colon cancer risk (27), and α -tocopherol is an antioxidant that inhibits mutagenesis and cell transformation (21). We therefore

calculated these ORs to know the risk in relation to serum oxLDL and oLAB independent of α -tocopherol and total cholesterol.

Two-sided P s < 0.05 were considered statistically significant. All statistical analyses were done using the Statistical Analysis System.

Results

Table 1 summarizes baseline characteristics of study subjects. No significant differences between cases and controls were observed for age distribution, smoking and drinking habits, family history of colorectal cancer, intake frequency of green leafy vegetables, or time spent in sports or physical exercise.

Table 2 compares serum levels of oxLDL, oLAB, total cholesterol, and α -tocopherol and BMI between cases and controls. Serum oxLDL levels were significantly higher in cases than in controls. BMI and serum levels of oLAB, α -tocopherol, and total cholesterol did not differ significantly between cases and controls.

Table 3 shows relationships among serum levels of oxLDL, oLAB, total cholesterol, and α -tocopherol in control subjects. Serum oxLDL levels were significantly and positively correlated with serum levels of total cholesterol and α -tocopherol in both genders. Serum

Table 1. Baseline characteristics of colorectal cancer cases and controls

	Cases (%)	Controls (%)	P (χ^2 test)
<i>n</i>	161 (100.0)	395 (100.0)	
Male	75 (46.6)	187 (47.3)	
Female	86 (53.4)	208 (52.7)	
Age (y)			
40-49	14 (8.7)	36 (9.1)	0.743
50-59	48 (29.8)	131 (33.2)	
60-69	65 (40.4)	159 (40.3)	
70-79	34 (21.1)	69 (17.5)	
Smoking habit			
Current smoker	37 (23.0)	96 (24.3)	0.674
Ex-smoker	23 (14.3)	51 (12.9)	
Nonsmoker	93 (57.8)	218 (55.2)	
Unknown	8 (5.0)	30 (7.6)	
Drinking habit			
Current drinker	75 (46.6)	168 (42.5)	0.503
Ex-drinker	2 (1.2)	12 (3.0)	
Nondrinker	78 (48.4)	195 (49.4)	
Unknown	6 (3.7)	20 (5.1)	
Family history of colorectal cancer			
Yes	10 (6.2)	14 (3.5)	0.161
No	151 (93.8)	381 (96.5)	
Intake frequency of green leafy vegetables			
1-2 times/mo or less	23 (14.3)	57 (14.4)	0.100
1-2 times/wk or more	126 (78.3)	325 (82.3)	
Unknown	12 (7.5)	13 (3.3)	
Time spent in sport or physical exercise			
Little	97 (60.2)	247 (62.5)	0.591
1 h/wk or more	57 (35.4)	125 (31.6)	
Unknown	7 (4.3)	23 (5.8)	

Table 2. Serum levels of oxLDL, oLAB, total cholesterol, and α -tocopherol and BMI for colorectal cancer cases and controls

	Cases		Controls		P
	n		n		
oxLDL (units/L), median (25th-75th percentiles)	119	39.2 (31.6-47.8)	316	36.2 (29.2-44.7)	0.045
oLAB (units/L), median (25th-75th percentiles)	153	201.0 (142.0-312.0)	376	192.0 (135.5-272.5)	0.120
Total cholesterol (mmol/L), mean \pm SD	159	5.22 \pm 0.96	382	5.17 \pm 0.98	0.225
α -Tocopherol (μ mol/L), median (25th-75th percentiles)	155	21.87 (15.67-27.35)	377	21.69 (17.30-26.75)	0.834
BMI (kg/m ²), mean \pm SD	158	23.1 \pm 3.4	380	23.2 \pm 2.8	0.779

oLAB levels displayed no correlation with serum levels of oxLDL, total cholesterol, or α -tocopherol in either gender.

Table 4 shows ORs and 95% CIs for colorectal cancer by serum levels of oxLDL and oLAB after adjusting for confounding factors. ORs (95% CIs) across quartiles for serum oxLDL adjusted for gender, age, and study area (OR1) were 1.21 (0.57-2.55), 1.49 (0.71-3.14), and 2.34 (1.03-5.30; $P_{\text{trend}} = 0.030$). OR (95% CI) for serum oxLDL adjusted for gender, age, study area, smoking and drinking habits, intake frequency of green leafy vegetables, time spent in sports or physical exercise, family history of colorectal cancer, and BMI (OR2) was significantly higher in the highest quartile compared with the lowest quartile [3.65 (1.50-8.92); $P_{\text{trend}} = 0.004$]. OR1 and OR2 for oLAB tended to be higher in the highest quartile of serum oLAB but not significantly (OR1, 1.66; 95% CI, 0.91-3.01; $P_{\text{trend}} = 0.148$; OR2, 1.68; 95% CI, 0.90-3.13; $P_{\text{trend}} = 0.140$).

When the analysis was limited to incident cases and corresponding controls, the higher risk was still found in relation to higher serum levels of oxLDL. OR2s (95% CIs) for colorectal cancer across quartiles of serum oxLDL were 3.11 (1.09-8.87), 2.25 (0.79-6.39), and 4.77 (1.50-15.10; $P_{\text{trend}} = 0.027$). OR2s (95% CIs) for colorectal cancer across quartiles of serum oLAB were 0.67 (0.32-1.41), 0.89 (0.41-1.92), and 1.22 (0.51-2.62; $P_{\text{trend}} = 0.412$).

Associations of serum oxLDL and oLAB with risk of colorectal cancer were also evaluated after further adjustment for quintiles of total cholesterol and α -tocopherol (OR3). However, no substantial change in results was observed. When evaluated by gender, no apparent difference between males and females was noted.

The same analyses were attempted using only colon cancer cases ($n = 80$ for oxLDL and $n = 106$ for oLAB) and corresponding controls ($n = 215$ for oxLDL and $n = 261$ for oLAB). OR3s (95% CIs) for colon cancer across

quartiles of serum oxLDL were 2.97 (0.97-9.06), 1.90 (0.55-6.59), and 4.68 (1.19-18.38; $P_{\text{trend}} = 0.062$). A similar trend was observed for serum oLAB levels: OR3s (95% CIs) across quartiles were 1.75 (0.73-4.20), 1.69 (0.68-4.15), and 2.20 (0.90-5.37; $P_{\text{trend}} = 0.119$).

Furthermore, modified data sets excluding cases diagnosed within 2 years from baseline were also analyzed. Results of these analyses were consistent with those of analyses without exclusion (data not shown).

Discussion

The present investigation represents the first prospective study to examine associations between serum oxLDL and risk of colorectal cancer. Significant positive associations were observed between serum oxLDL levels and risk of colorectal cancer. There was no association between serum oLAB levels and risk of colorectal cancer. Risk of colorectal cancer was higher in the presence of higher levels of serum oxLDL, independent of confounders. The mechanisms involved in this association between oxLDL and colorectal cancer remain unclear.

The adjustment for lifestyle factors, family history, and BMI somewhat strengthened the positive association between oxLDL and risk of colorectal cancer. This may not be in line with our initial hypothesis that serum oxLDL levels represent a marker reflecting lifestyles related to the cancer. Serum oxLDL may be a predictor of the risk independently of other risk factors.

There are some reports that studied the association between serum or plasma oxLDL levels and coronary heart disease (7, 28). It is well known that oxLDL is found in monocyte-derived macrophages in atherosclerosis lesions and that plasma oxLDL levels were significantly higher in patients with coronary artery disease (28). Several studies have been carried out on the modified forms of oxLDL, which are prepared by oxidizing LDL under various conditions *in vitro* (28). However, there is little information about oxLDL present *in vivo* (28).

We have also studied associations between serum carotenoids levels and risk of colorectal cancer in this prospective epidemiologic study. We found inverse associations of some carotenoids with colorectal cancer risk in men.⁹ Crohn disease is a chronic inflammatory disorder and is associated with increased risk of colon cancer (29). Although the etiology of Crohn disease is unknown, patients with Crohn disease have increased

Table 3. Spearman correlation coefficients (no. subjects) among serum levels of oxLDL, oLAB, total cholesterol, and α -tocopherol among control subjects

	oxLDL	oLAB
Males		
oLAB	0.066 (136)	
Total cholesterol	0.525* (138)	-0.023 (172)
α -Tocopherol	0.397* (140)	0.011 (169)
Females		
oLAB	0.021 (161)	
Total cholesterol	0.429* (166)	-0.093 (191)
α -Tocopherol	0.227* (168)	0.006 (192)

* $P < 0.001$.⁹ Unpublished data.

Table 4. ORs and 95% CIs for colorectal cancer risk by serum levels of oxLDL and oLAB

	Range	Cases	Controls	OR1	95% CI	<i>P</i> _{trend}	OR2	95% CI	<i>P</i> _{trend}	OR3	95% CI	<i>P</i> _{trend}
oxLDL (units/L)												
Q1	≤29.1	22	79	1.00	—	0.030	1.00	—	0.004	1.00	—	0.038
Q2	29.2-36.1	26	79	1.21	0.57-2.55		1.55	0.70-3.46		1.15	0.49-2.72	
Q3	36.2-44.6	31	79	1.49	0.71-3.14		1.90	0.84-4.28		1.38	0.54-3.51	
Q4	≥44.7	40	79	2.34	1.03-5.30		3.65	1.50-8.92		3.10	1.04-9.23	
oLAB (units/L)												
Q1	≤135.4	34	94	1.00	—	0.148	1.00	—	0.140	1.00	—	0.212
Q2	135.5-191.9	41	96	1.14	0.64-2.01		0.98	0.54-1.80		1.11	0.58-2.11	
Q3	192.0-272.4	28	92	0.87	0.46-1.64		0.75	0.39-1.48		0.74	0.36-1.52	
Q4	≥272.5	50	94	1.66	0.91-3.01		1.68	0.90-3.13		1.69	0.85-3.35	

NOTE: OR1: OR adjusted for gender, age, and study area; OR2: OR adjusted for gender, age, study area, smoking and drinking habits, intake frequency of green leafy vegetables, time spent in sport or physical exercise, family history of colorectal cancer, and BMI; OR3: OR adjusted for gender, age, study area, smoking and drinking habits, intake frequency of green leafy vegetables, time spent in sport or physical exercise, family history of colorectal cancer, BMI, and serum levels of total cholesterol and α -tocopherol.

production of ROS (29). It was reported that lipid peroxidation and F₂ isoprostane was significantly higher in patients with Crohn disease than in healthy control subjects (30).

Various potentially toxic oxidized lipids are contained in oxLDL such as lipid peroxides, oxysterol, and aldehydes (31). These oxidized lipids elicit oxidative stress and lipid peroxidation (31). As oxLDL reduces antioxidant enzymes such as Cu/Zn superoxide dismutase (32) and glutathione peroxidase (33) and ROS degradation is decreased following increases in oxLDL (31), ROS levels are elevated. Lipid peroxidation is initiated by ROS attacks, generating large amounts of reactive products that have been implicated in tumor initiation and promotion (34). Increased levels of malondialdehyde, a major genotoxic carbonyl compound generated by lipid peroxidation (34), have been reported in tumor tissue from colorectal cancer patients compared with normal mucosa from the same individuals (35).

In another experimental study (8), oxLDL-induced oxidative stress enhanced p53 DNA binding activity and p53 protein synthesis. As a tumor suppressor, p53 is induced by various kinds of cell stress (36) to protect the cell. Genetic information is protected by the functions of p53, including induction of cell cycle arrest or apoptosis after DNA damage and maintenance of genomic stability (37). Given the above, high levels of oxLDL might induce excess stress against the cell. This stress may induce DNA damage and mutation, because oxidative stress is known to cause such damage (38). Mutation of *p53* gene is found in >50% of all human cancers and >75% of colorectal adenocarcinomas (39). Mutation of the *p53* gene is known to play crucial roles in tumor development and progression (37).

Cyclooxygenase-2 (COX-2) expression is reportedly induced by oxLDL in a murine macrophage-like cell line (40) and human monocytes (41). COX is an enzyme that initiates the conversion of arachidonic acid into all of the prostaglandins and thromboxanes (42). Lipid peroxidation is necessary for initiation of COX activity (43), and reactive oxygen intermediates (ROI) induce COX-2 (44). Levels of arachidonic acid (45) and prostaglandin E₂ (46) are higher in colon tumor than in normal colonic mucosa. Prostaglandin E₂, a major product of COX, stimulates proliferation and growth of human colorectal cancer cells (47).

Analysis of COX-2 expression (induced by cytokines, growth factors, and mitogens) has revealed elevated levels in up to 90% of sporadic colon carcinomas and 40% of colonic adenomas but no elevation in normal colonic epithelium (48). Recent clinical epidemiologic studies have shown that COX inhibitors such as aspirin and other nonsteroidal anti-inflammatory agents exert preventive effects on colorectal cancer (49, 50). Such inhibition of COX-2 is considered to lead to decreased incidence of colorectal cancer, although the mechanisms are not fully understood.

Functions of oxLDL such as increasing oxidative stress and inducing COX-2 expression might play an important role in colorectal carcinogenesis. At the very least, oxidative stress is increased in subjects with high levels of serum oxLDL, and oxidative stress should be related to colorectal carcinogenesis.

Epidemiologic studies showed the close association between insulin resistance and colon cancer risk (51). The consumption of excess dietary energy results in the development of insulin resistance with increased circulating levels of insulin, triglycerides, and nonesterified fatty acids. These circulating factors subject colonic epithelial cells to a proliferative stimulus and also expose them to reactive oxygen intermediates. Other study reported that LDL oxidizability is increased in insulin resistance subjects compared with healthy subjects (52). These long-term exposures are expected to result in the promotion of colon cancer.

Serum oLAB levels were not significantly associated with risk of colorectal cancer. Serum oLAB is generated from immunoresponses against oxLDL. Serum oLAB levels, in addition to serum oxLDL levels, may therefore also depend on various lifestyle factors such as dietary intake of antioxidants and smoking habits. Plasma oLAB levels are reported to show a negative correlation with plasma oxLDL levels in healthy subjects (53), and oLAB may play a role in maintaining low levels of blood oxLDL. Wide ranges of serum oLAB levels might reflect interindividual differences in immune responses rather than in oxLDL generation. The immune system is also affected by various lifestyle factors such as smoking habits. We considered that almost no relationship between serum oLAB and oxLDL in controls was derived from interindividual differences in immune responses. Interindividual difference in immune responses may

have also attenuated the association between serum oLAB and risk of colorectal cancer.

Although oxLDL is an oxidant and α -tocopherol is an antioxidant, our results show positive association between serum oxLDL and α -tocopherol levels. We suggest that this association was observed because serum LDL binds to α -tocopherol (54). Similarly, serum oxLDL is positively associated with serum cholesterol levels.

Cases included both colon and rectal cancers. Risk for colon cancer only was increased with high serum oxLDL levels after adjusting for gender, age, study area, and potential confounders. The sample population for rectum cancer cases was too small to analyze associations between serum levels of oxLDL and oLAB and risk of rectum cancer. These associations warrant further study.

In conclusion, the present study showed that increased levels of serum oxLDL represent a risk factor for colorectal cancer among Japanese. Although further investigations are needed to clarify the role of oxLDL in tumorigenesis for colorectal cancer, serum oxLDL levels may be one biomarker for predicting risk of colorectal cancer.

Acknowledgments

We thank K. Aoki (Emeritus Professor, Nagoya University Graduate School of Medicine and former Chairman of the JACC Study Group) and H. Sugano (former Director of the Cancer Institute of the Japanese Foundation for Cancer Research) for great contribution to the initiation of the study.

References

- Channon KM. Oxidative stress and coronary plaque stability. *Arterioscler Thromb Vasc Biol* 2002;22:1751–2.
- Saintot M, Astre C, Pujol H, Gerber M. Tumor progression and oxidant-antioxidant status. *Carcinogenesis* 1996;17:1267–71.
- Guyton KZ, Kensler TW. Oxidative mechanisms in carcinogenesis. *Br Med Bull* 1993;49:523–44.
- Keshavarzian A, Zapeda D, List T, Mobarhan S. High levels of reactive oxygen metabolites in colon cancer tissue: analysis by chemiluminescence probe. *Nutr Cancer* 1992;17:243–9.
- Otamiri T, Sjadahl R. Increased lipid peroxidation in malignant tissues of patients with colorectal cancer. *Cancer* 1989;64:422–5.
- Witztum JL. The oxidation hypothesis of atherosclerosis. *Lancet* 1994;344:793–5.
- Witztum JL, Steinberg D. Role of oxidized low density lipoprotein in atherogenesis. *J Clin Invest* 1991;88:1785–92.
- Maziere C, Meignotte A, Dantin F, Conte MA, Maziere JC. Oxidized LDL induces an oxidative stress and activates the tumor suppressor p53 in MRC5 human fibroblasts. *Biochem Biophys Res Commun* 2000;276:718–23.
- Elosua R, Molina L, Fito M, et al. Response of oxidative stress biomarkers to a 16-week aerobic physical activity program, and to acute physical activity, in healthy young men and women. *Atherosclerosis* 2003;167:327–34.
- Vasankari T, Fogelholm M, Kukkonen-Harjula K, et al. Reduced oxidized low-density lipoprotein after weight reduction in obese premenopausal women. *Int J Obes Relat Metab Disord* 2001;25:205–11.
- Esterbauer H, Gebicki J, Puhl H, Jurgens G. The role of lipid peroxidation and antioxidants in oxidative modification of LDL. *Free Radic Biol Med* 1992;13:341–90.
- Chopra M, O'Neill ME, Keogh N, Wortley G, Southon S, Thurnham DI. Influence of increased fruit and vegetable intake on plasma and lipoprotein carotenoids and LDL oxidation in smokers and non-smokers. *Clin Chem* 2000;46:1818–29.
- Calzada C, Bizzotto M, Paganga G, et al. Levels of antioxidant nutrients in plasma and low density lipoproteins: a human volunteer supplementation study. *Free Radical Res* 1995;23:489–503.
- Abbey M, Nestel PJ, Baghurst PA. Antioxidant vitamins and low-density-lipoprotein oxidation. *Am J Clin Nutr* 1993;58:525–32.
- Martinez ME, Giovannucci E, Spiegelman D, Hunter DJ, Willett WC, Colditz GA. Leisure-time physical activity, body size, and colon cancer in women. Nurses' Health Study Research Group. *J Natl Cancer Inst* 1997;89:948–55.
- Nilsen TI, Vatten LJ. Prospective study of colorectal cancer risk and physical activity, diabetes, blood glucose and BMI: exploring the hyperinsulinemia hypothesis. *Br J Cancer* 2001;84:417–22.
- Thune J, Lund E. Physical activity and risk of colorectal cancer in men and women. *Br J Cancer* 1996;73:1134–40.
- Garland C, Shekelle RB, Barrett-Connor E, Criqui MH, Rossof AH, Paul O. Dietary vitamin D and calcium and risk of colorectal cancer: a 19-year prospective study in men. *Lancet* 1985;325:307–9.
- Nomura A, Heilbrun LK, Stemmermann GN. Body mass index as a predictor of cancer in men. *J Natl Cancer Inst* 1985;74:319–23.
- Chapter 4.10. Colon, rectum. In: Food, nutrition and the prevention of cancer: a global perspective. Washington (DC): World Cancer Research; 1997. p. 216–51. Fund in association with American Institute for Cancer Research.
- Longnecker MP, Martin-Moreno JM, Knekt P, et al. Serum α -tocopherol concentration in relation to subsequent colorectal cancer: pooled data from five cohorts. *J Natl Cancer Inst* 1992;84:430–5.
- Ohno Y, Tamakoshi A, JACC Study Group. Japan Collaborative Cohort Study for Evaluation of Cancer Risk sponsored by Monbusho (JACC Study). *J Epidemiol* 2001;11:144–50.
- WHO. International Statistical Classifications of Diseases and Related Health Problems. 10th Revision. Vol. I. Geneva: WHO; 1992. p. 1–1243.
- Suzuki K, Ito Y, Ochiai J, et al., for the JACC Study Group. Relationship between obesity and serum markers of oxidative stress and inflammation in Japanese. *Asian Pac J Cancer Prev* 2003;4:259–66.
- Suzuki K, Ito Y, Ochiai J, et al., for the JACC Study Group. The relationship between smoking habits and serum levels of 8-OHdG, oxidized LDL antibodies, Mn-SOD and carotenoids in rural Japanese residents. *J Epidemiol* 2003;13:29–37.
- Ito Y, Ochiai J, Sasaki R, et al. Serum concentrations of carotenoids, retinol, and α -tocopherol in healthy persons determined by high performance liquid chromatography. *Clin Chem Acta* 1990;194:131–44.
- Quadrilatero J, Hoffman-Goetz L. Physical activity and colon cancer. A systematic review of potential mechanisms. *J Sports Med Phys Fitness* 2003;43:121–38.
- Itabe H. Oxidized low-density lipoproteins: what is understood and what remains to be clarified. *Biol Pharm Bull* 2003;26:1–9.
- Orlando RC. Mechanisms of epithelial injury and inflammation in gastrointestinal diseases. *Rev Gastroenterol Disord* 2002;2 Suppl 2:S2–8.
- Wendland BE, Aghdassi E, Tam C, et al. Lipid peroxidation and plasma antioxidant micronutrients in Crohn disease. *Am J Clin Nutr* 2001;74:259–64.
- Salvyre R, Auge N, Benoist H, Negre-Salvyre A. Oxidized low-density lipoprotein-induced apoptosis. *Biochim Biophys Acta* 2002;1585:213–21.
- Li W, Yuan XM, Brunk UT. OxLDL-induced macrophage cytotoxicity is mediated by lysosomal rupture and modified by intralysosomal redox-active iron. *Free Radic Res* 1998;29:389–98.
- Rosenblat M, Aviram M. Macrophage glutathione content and glutathione peroxidase activity are inversely related to cell-mediated oxidation of LDL: *in vitro* and *in vivo* studies. *Free Radic Biol Med* 1998;24:305–17.
- Marnett LJ. Oxyradicals and DNA damage. *Carcinogenesis* 2000;21:361–70.
- Hendrickse CW, Kelly RW, Radley S, Donovan IA, Keighley MR, Neoptolemos JP. Lipid peroxidation and prostaglandins in colorectal cancer. *Br J Surg* 1994;81:1219–23.
- Khanna KK, Lavin MF. Ionizing radiation and UV induction of p53 protein by different pathways in ataxia-telangiectasia cells. *Oncogene* 1993;8:3307–12.
- Nakamura Y. Isolation of p53-target genes and their functional analysis. *Cancer Sci* 2004;95:7–11.
- Hussain SP, Amstad P, Raja K, et al. Increased p53 mutation load in noncancerous colon tissue from ulcerative colitis: a cancer-prone chronic inflammatory disease. *Cancer Res* 2000;60:3333–7.
- Pontsler AV, St. Hilaire A, Marathe GK, Zimmerman GA, McIntyre TM. Cyclooxygenase-2 is induced in monocytes by peroxisome proliferator activated receptor γ and oxidized alkyl phospholipids from oxidized low density lipoprotein. *J Biol Chem* 2002;277:13029–36.
- Fyrnys B, Claus R, Wolf G, Deigner HP. Oxidized low density lipoprotein stimulates protein kinase C (PKC) activity and expression of PKC-isotypes via prostaglandin-H-synthase in P388D1 cells. *Adv Exp Med Biol* 1997;407:93–8.
- Kinzler KW, Vogelstein B. Lessons from hereditary colorectal cancer. *Cell* 1996;87:159–70.

42. Taketo MM. Cyclooxygenase-2 inhibitors in tumorigenesis. *J Natl Cancer Inst* 1998;90:1529–36.
43. Capdevila JH, Morrow JD, Belosludtsev YY, Beauchamp DR, DuBois RN, Falck JR. The catalytic outcomes of the constitutive and the mitogen inducible isoforms of prostaglandin H2 synthase are markedly affected by glutathione and glutathione peroxidase(s). *Biochemistry* 1995;34:3325–37.
44. Feng L, Xia Y, Garcia GE, Hwang D, Wilson CB. Involvement of reactive oxygen intermediates in cyclooxygenase-2 expression induced by interleukin-1, tumor necrosis factor- α , and lipopolysaccharide. *J Clin Invest* 1995;95:1669–75.
45. Nicholson ML, Neoptolemos JP, Clayton HA, Talbot IC, Bell PR. Increased cell membrane arachidonic acid in experimental colorectal tumors. *Gut* 1991;32:413–8.
46. Hendrickse CW, Kelly RW, Radley S, Donovan IA, Keighley MR, Neoptolemos JP. Lipid peroxidation and prostaglandins in colorectal cancer. *Br J Surg* 1994;81:1219–23.
47. Sheng H, Shao J, Washington MK, DuBois RN. Prostaglandin E₂ increases growth and motility of colorectal carcinoma cells. *J Biol Chem* 2001;276:18075–81.
48. Eberhart CE, Coffey RJ, Radhika A, Giardiello FM, Ferrenbach S, DuBois RN. Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology* 1994;107:1183–8.
49. Kawamori T, Rao CV, Seibert K, Reddy BS. Chemopreventive activity of celecoxib, a specific cyclooxygenase-2 inhibitor, against colon carcinogenesis. *Cancer Res* 1998;58:409–12.
50. Reddy BS, Rao CV, Seibert K. Evaluation of cyclooxygenase-2 inhibitor for potential chemopreventive properties in colon carcinogenesis. *Cancer Res* 1996;56:4566–9.
51. Bruce WR, Giacca A, Medline A. Possible mechanisms relating diet and risk of colon cancer. *Cancer Epidemiol Biomarkers Prev* 2000;9:1271–9.
52. Schwenke DC, D'Agostino RB Jr, Goff DC Jr, Karter AJ, Rewers MJ, Wagenknecht LE. Differences in LDL oxidizability by glycemic status: the insulin resistance atherosclerosis study. *Diabetes Care* 2003;26:1449–55.
53. Shoji T, Nishizawa Y, Fukumoto M, et al. Inverse relationship between circulating oxidized low density lipoprotein (oxLDL) and anti-oxLDL antibody levels in healthy subjects. *Atherosclerosis* 2000;148:171–7.
54. Parks EJ, German JB, Davis PA, et al. Reduced oxidative susceptibility of LDL from patients participating in an intensive atherosclerosis treatment program. *Am J Clin Nutr* 1998;68:778–85.

Serum Oxidized Low-Density Lipoprotein Levels and Risk of Colorectal Cancer: A Case-Control Study Nested in the Japan Collaborative Cohort Study

Koji Suzuki, Yoshinori Ito, Kenji Wakai, et al.

Cancer Epidemiol Biomarkers Prev 2004;13:1781-1787.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/13/11/1781>

Cited articles This article cites 48 articles, 13 of which you can access for free at:
<http://cebp.aacrjournals.org/content/13/11/1781.full#ref-list-1>

Citing articles This article has been cited by 2 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/13/11/1781.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/13/11/1781>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.