

# Physical Activity and Risk of Endometrial Cancer: A Report from the Shanghai Endometrial Cancer Study

Charles E. Matthews,<sup>1</sup> Wang Hong Xu,<sup>2</sup> Wei Zheng,<sup>1</sup> Yu Tang Gao,<sup>2</sup> Zhi Xian Ruan,<sup>2</sup> Jia Rong Cheng,<sup>2</sup> Yong Bing Xiang,<sup>2</sup> and Xiao Ou Shu<sup>1</sup>

<sup>1</sup>Division of General Internal Medicine, Department of Medicine, Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, TN and <sup>2</sup>Department of Epidemiology, Shanghai Cancer Institute, Shanghai, People's Republic of China

## Abstract

We evaluated the type and amount of physical activity associated with risk of endometrial cancer. In this population-based case-control study, in-person interviews were completed among 832 incident endometrial cancer cases and 846 age-matched controls. Physical activity from exercise, household activities, and transportation was assessed in adolescence and adulthood, as was lifetime occupational activity. Logistic regression was used to estimate adjusted odds ratios (OR) and 95% confidence limits (95% CL). Women reporting exercise participation in both adolescence and adulthood were at nearly a 40% reduced risk (OR, 0.63; 95% CL, 0.42-0.95), compared with women reporting no exercise in either life period. Postmenopausal women who initiated exercise in adult-

hood were also at reduced risk (OR, 0.76; 95% CL, 0.56-1.02). Reductions in risk were also observed for common lifestyle activities, including household activity (both life periods) and walking for transportation (adulthood). Examination of the independent and combined effect of exercise and lifestyle activities revealed that women with less active lifestyles but who reported exercise were at 35% reduced risk (OR, 0.65; 95% CL, 0.41-1.02), whereas nonexercisers with more active lifestyles were at 40% to 45% reduced risk. These findings suggest that both lifestyle activities of lower intensity (e.g., walking and doing household chores) and intentional exercise can reduce endometrial cancer risk. (Cancer Epidemiol Biomarkers Prev 2005;14(4):779-85)

## Introduction

Endometrial cancer accounts for about 5% of all cancers among women in more developed countries of the world (1), and it is the most prevalent gynecologic cancer and fourth most frequently diagnosed cancer among women in the United States (2). In Shanghai, China, the age-adjusted incidence of endometrial cancer increased by 75% between 1972 to 1974 and 1993 to 1994 (3, 4). These striking increases in the incidence of this disease in Shanghai parallel marked weight gain among women in China that has been linked, in part, to physical inactivity (5, 6).

Nongenetic factors have been estimated to account for nearly 80% of endometrial cancer cases (7), and physical inactivity has been found an important and potentially modifiable risk factor for this disease. For example, four (7-10) of seven (57%; refs. 11-13) prospective studies and 9 (14-22) of 10 (90%; ref. 23) case-control studies have suggested that lower levels of physical activity increase risk for endometrial cancer. The average risk reduction in these studies has been roughly 30% to 50%, and only one of these reports has suggested that higher levels of activity increase risk (11). Whereas a number of studies have reported benefit from both occupational (9, 20, 24) and nonoccupational activity (10, 14, 18, 20), our understanding of the amount and intensity of activity that may confer benefit and the influence of changes in activity patterns over the life course remain incomplete.

Accordingly, the purpose of this research was to examine the relationship between newly diagnosed endometrial cancer and a broad range of physical activity behaviors (i.e.,

household, occupation, walking and cycling for transportation, and exercise). In particular, we considered the type/intensity and duration of activity and the effect of changes in exercise behaviors between adolescence and adulthood on risk.

## Materials and Methods

The Shanghai Endometrial Cancer Study was designed to evaluate the genetic and lifestyle factors associated with endometrial cancer risk. Incident cases who were permanent residents of urban Shanghai (age 30-69 years) were identified from the Shanghai Cancer Registry between January 1997 and December 2001. The diagnosis of each case was confirmed by medical chart review and a review of the available pathology slides by senior study pathologists. In-person interviews were completed for 832 of 982 (85%) eligible cases. Among case nonparticipants, 73 (7%) declined participation, 36 (4%) were deceased, 22 (2%) could not be located, 10 (1%) were absent during the study period, and 9 (1%) were excluded for miscellaneous reasons. Controls, frequency matched to cases by age ( $\pm 5$  years), were randomly selected from permanent female residents using the Shanghai Resident Registry. Women who had a hysterectomy were not eligible. Eligible controls ( $n = 1,165$ ) were identified for possible participation and 846 women (73%) were interviewed for the study. Among control nonparticipants, 277 (24%) declined participation and 42 (4%) were absent during the study period. Trained retired nurses and physicians recruited, consented, and completed structured in-person interviews to gather information about demographic factors, menstrual and reproductive history, hormone use, usual dietary intake, prior disease history, tobacco and alcohol use, family history of cancer, and height and weight history during adolescence and adulthood. Each participant also had a number of anthropometric indices measured using a standard protocol (current weight, height, and waist and hip circumference; ref. 25).

Received 9/8/04; revised 11/11/04; accepted 12/13/04.

Grant support: ROICA92585 (X.O. Shu).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Charles E. Matthews, Division of General Internal Medicine, Vanderbilt University School of Medicine, Medical Center East, Suite 6100, Nashville, TN 37232-8300. Phone: 615-936-2145; Fax: 615-936-1269. E-mail: charles.matthews@vanderbilt.edu

Copyright © 2005 American Association for Cancer Research.

Physical activity patterns were also assessed during the in-person interview. Regular exercise and sport participation was evaluated for adulthood (past 10 years) and adolescence (13-19 years). Up to five exercise activities were reported in each life period, and quantitative data were obtained for each (i.e., type, duration, years of participation). These data were summarized in terms of intensity [metabolic equivalents (MET)], duration (hours/wk), years of participation, as an average energy expenditure in the period (MET-hours per week per year) using standard methods (26). Women reported each occupation they held for at least 3 years during their lifetime and the calendar year in which they started and ended each job. Occupations were classified into high, medium, or low levels of estimated sitting time and activity level using job codes. For example, occupations categorized into the high sitting time and low-activity categories were clerical workers, government and business clerks, and accountants, whereas occupations categorized into the low sitting time and higher-activity categories were sales clerks, cooks, and machine operators. For each occupation, participants also reported the average time they spent "standing or walking" (hours/d) and classified each occupation into one of four activity categories (i.e., heavy, medium, light, or nonphysical work). Multiplying years spent in each occupation by the specific activity variable and summing the result over all occupations calculated summary occupational indices. Time spent in several common nonoccupational (or lifestyle activities) were also evaluated in both adolescence and adulthood [i.e., walking for transportation (minutes/d), cycling for transportation (minutes/d), and housework (hours/d)]. Summary energy expenditure values (MET-hours/d) for lifestyle activities were estimated using the following MET values: housework, 2.0 METs; walking, 3.3 METs; and cycling, 4.0 METs, using relevant compendium values as a reference (26). One MET-hour/d is roughly equivalent to 1 kcal per kg per day or about 15 minutes of participation in a moderate intensity (4 METs) activity for an average adult (27). We have evaluated a similar physical activity assessment instrument among women in Shanghai and found it reasonable measure of physical activity in this population (28).

**Statistical Analysis.** Unconditional logistic regression models were used to estimate odds ratios (OR) and their 95% confidence limits (95% CL), test for multiplicative interaction, and adjust for confounding (29). Potential confounding factors evaluated included age, education, family income, age at menarche, oral contraceptive use, number of pregnancies, menopausal status, age at menopause, height, body mass index, waist-to-hip ratio, and family history of cancer. Hormone replacement therapy is very uncommon among Chinese women and its use was not related to endometrial cancer risk (i.e., 4.2% for cases and 4.0% for controls). The quantile distributions among controls were used to select cut points to classify continuous exposure variables. Linear trends in the risk estimates for physical activity were tested statistically by entering the categorized ordinal variables for the physical activity exposures in the models. Multiplicative interactions were examined by fitting the two exposure variables and their cross-product terms in the logistic model. Body mass index was evaluated as both a confounder and an effect modifier, and stratified analyses were completed by prediagnosis menopausal status.

## Results

The average age of participants was 55.3 years and there were no differences in age by disease status ( $P = 0.65$ ). Women with more education, later age at menopause, family history of cancer, greater height, greater body mass index, and a higher

waist-to-hip ratio were at increased risk; whereas women with a later age at menarche, more pregnancies, reporting oral contraceptive use in formulations primarily containing both estrogen and progesterone, and that currently smoked or ever reported alcohol consumption were at reduced risk (Table 1). Family income, a family history of endometrial cancer, and hormone replacement therapy use were not associated with risk. Among the population-based controls, the majority of women (50-60%) reported no exercise in adolescence or adulthood, and fewer than 10% reported exercise in both life periods. The prevalence of exercise in adolescence among premenopausal women was about twice that of postmenopausal women (20.7% versus 8.6%). Whereas exercise participation in adulthood was more prevalent among postmenopausal compared with premenopausal women (32.8% versus 12.3%), about two thirds of women (67.2%) reported no exercise participation in adulthood (Fig. 1).

Exercise participation in both adolescence and adulthood was associated with a 30% to 40% reduction in risk for both premenopausal and postmenopausal women. Among these women, 75% reported at least 6 years of exercise participation and >2 MET-hours per day per year of exercise energy expenditure or about 30 minutes/d of moderate intensity activity (4 METs). Total exercise energy expenditure accumulated in both life periods was inversely associated with endometrial cancer risk (Table 2). Examination of exercise effects using mutually exclusive classifications in each life period (Table 2, *bottom rows*) indicated that postmenopausal women who initiated exercise in adulthood were at about a 25% reduced risk (OR, 0.76; 95% CL, 0.56-1.02) compared with women who reported no exercise participation. Exercise only in adolescence reduced risk by about 20%, although the confidence intervals included the null.

In general, we found no consistent associations between occupational physical activity and endometrial cancer, using either job code information or methodologically stronger self-reports of individual occupational activity exposures (e.g., self-rating and time standing and walking; Table 3). Women with jobs thought to require more sitting time were associated with increased risk among premenopausal women but reduced risk among postmenopausal women (Table 3). These results remained unchanged even after further controlling for exercise participation in adolescence and adulthood (data not shown).

Evaluation of common nonexercise or lifestyle activities, revealed inverse associations between endometrial cancer and household activity (both life periods) and walking for transportation (adulthood). There was no apparent association with cycling for transportation, but only about 20% of women reported this activity in adulthood. Total household activity in both life periods was also inversely associated with risk ( $P_{\text{trend}} < 0.01$ ). Inverse relationships were present for total lifestyle activity accumulated in each life period, and maximal reductions in risk of 30% to 40% were evident in comparing the upper to the lower quartiles of activity (Table 4, *bottom row*).

In detailed analyses by menopausal status, there was a suggestion of stronger effects of household and walking activities in adulthood among premenopausal compared with postmenopausal women (roughly 50% versus 30% reduced risk), although this effect was not evident when total lifestyle activity in adulthood was evaluated. Total lifestyle activity reported in adolescence was associated with reduced risk among both postmenopausal and premenopausal women (OR<sub>premenopausal</sub>, 0.83; 95% CL, 0.49-1.40; OR<sub>postmenopausal</sub>, 0.51; 95% CL, 0.35-0.74), although the point estimate did not reach significance for premenopausal women. We found no evidence of striking differences in risk by menopausal status for walking for transportation in adolescence or adulthood. To delineate the independent and combined effects of exercise and lifestyle activity reported in adulthood, we evaluated the

joint effect of these exposures using a common referent group (i.e., nonexercisers in the lower quartile of lifestyle activity; Fig. 2). Women with the least active lifestyles (<7.30 MET-hours/d), but that reported exercise participation were at

**Table 1. Descriptive comparisons between cases (*n* = 832) and controls (*n* = 846) on endometrial cancer risk, the Shanghai Endometrial Cancer Study (1997-2001)**

	Case/controls*	OR (95% CI)
<b>Education</b>		
No formal education	75/108	1.00
Elementary	129/126	1.47 (1.01-2.16)
Junior high school	296/286	1.49 (1.07-2.09)
High school	207/227	1.31 (0.93-1.86)
Post-high school/college	125/99	1.82 (1.22-2.70)
<b>Family income (Yuan)</b>		
<10,000	95/89	1.00
10,000-15,000	199/194	0.96 (0.68-1.36)
15,000-20,000	166/155	1.00 (0.70-1.44)
20,000-30,000	197/244	0.76 (0.54-1.07)
>30,000	174/163	1.00 (0.70-1.43)
<b>Reproductive factors</b>		
Age at menarche (y)		
9-12	108/78	1.00
13-14	334/301	0.81 (0.58-1.12)
15	167/178	0.68 (0.48-0.98)
>16	221/287	0.56 (0.40-0.79)
No. pregnancies		
0/1	199/144	1.00
2	199/208	0.69 (0.52-0.93)
3	194/207	0.68 (0.51-0.91)
4	141/157	0.65 (0.48-0.89)
>5	99/130	0.55 (0.39-0.77)
Menopause		
No	327/300	1.00
Yes	505/546	0.85 (0.70-1.03)
Age at menopause (y)		
17-46	69/132	1.00
47-48	90/117	1.45 (0.97-2.16)
49-50	146/155	1.78 (1.23-2.57)
>51	199/142	2.64 (1.84-3.79)
Oral contraceptive use		
Never	685/639	1.00
Ever	147/207	0.66 (0.52-0.84)
<b>Lifestyle factors</b>		
Current smoker		
No	823/816	1.00
Yes	7/25	0.28 (0.12-0.65)
Drink ever		
No	812/804	1.00
Yes	20/42	0.47 (0.27-0.81)
Family history of cancer		
No	535/613	1.00
Yes	289/228	1.45 (1.18-1.79)
<b>Body size</b>		
Height (cm)		
140-152	148/184	1.00
153-156	201/195	1.26 (0.94-1.69)
157-160	246/247	1.22 (0.92-1.61)
160-174	233/218	1.31 (0.99-1.74)
Body mass index (kg/m <sup>2</sup> )		
14.5-21.3	120/211	1.00
21.4-23.5	166/198	1.40 (1.04-1.90)
21.6-26.1	214/216	1.66 (1.24-2.21)
26.2-41.9	323/216	2.50 (1.90-3.31)
Waist/hip ratio		
0.60-0.77	99/203	1.00
0.78-0.80	136/190	1.44 (1.04-1.99)
0.81-0.84	237/215	2.22 (1.64-2.99)
0.85-1.21	356/234	3.06 (2.29-4.08)

Abbreviation: 95% CI, 95% confidence interval.

\**n* values for cases and controls may be <832/846 due to missing data for some variables.

lower risk (OR, 0.65; 95% CL, 0.41-1.02). Similarly, women with more active lifestyles ( $\geq 9.85$  MET-hours/d) but that did not exercise were at 40% to 45% lower risk ( $P < 0.05$ ).

We also carefully evaluated our main results for effect modification and confounding by body mass index and found no significant multiplicative interactions ( $P > 0.05$ ) or major differences in stratum specific risk estimates for our main results. Similarly, body mass index did not seem a strong confounding factor with the physical activity/endometrial cancer association in these data.

## Discussion

The present findings are consistent with the growing body of literature indicating that high levels of physical activity are associated with reduced risk for endometrial cancer (8, 18, 19, 30). In this research, long-term exercise participation and high levels of common nonexercise or lifestyle activities (e.g., housework and walking for transportation) were associated with 30% to 40% reductions in risk. Our finding that higher levels of household activity and walking support the idea that broad range of lower intensity activities provide important benefit in terms of endometrial cancer risk.

The biological mechanisms through which physical activity influences the natural history of endometrial cancer are believed to be related to hormonal and insulin mediated pathways. Hyperinsulinemia has been linked to reduced sex hormone-binding globulin, which may cause increased estrogen and androgen exposures in endometrial tissue (31). Hyperinsulinemia also increases circulating insulin-like growth factor-I directly, whereas also reducing its binding protein-1 in endometrial tissue, both of which may increase tissue levels of bioavailable insulin-like growth factor-I (31). Higher levels of physical activity are associated with reduced insulin levels (32) and insulin sensitivity (33, 34) and therefore would be predicted to reduce exposure to bioavailable hormones and growth factors. The permissive influence of insulin on hormonal and growth factor exposure is plausible in both premenopausal and postmenopausal women. However, among postmenopausal women, where estrogen exposure is primarily determined by conversion of androgens from adipose tissue (35), physical activity effects on disease risk may be derived through its influence on energy balance and adipose accumulation. Higher physical activity levels have been associated with lower levels of adiposity in Chinese women (6), and reductions in adiposity via increased physical activity levels have been shown to reduce free estradiol levels (36).

There are several strengths and limitations to this research that should be considered when interpreting our results. First, like other case-control studies, recall bias is a concern. However, the consistency of our findings with a number of prospective studies (7-10) suggests that recall bias did not seem a major concern. Selection bias is also possible. Whereas this population-based study with true community controls had reasonably high response rates for both cases (85%) and controls (73%), the modest differences in these rates between comparison groups indicates some potential for the introduction of bias. However, if selection bias were present, we believe that it would be expressed most acutely in the exercise exposure data because participants in research studies are often more health conscious than nonparticipants. Our finding of a protective association in both exercise and nonexercise activity commonly done by most participants argues against a selection bias in our results. The validity of self-reported physical activity is also a concern. Our validation work with a nearly identical version of the physical activity assessment used in this research supports the ability of this instrument to stratify women into high and low activity levels for the physical activity domains we examined in this research, even light to moderate intensity




Figure 1. Prevalence of exercise participation among control women by life period and menopausal status.

household activity (28). A notable strength of this instrument was that it examined a broad range of physical activities and particularly household activities. Household activity is an important source of energy expenditure among women (37).

It is not immediately clear why we did not observe an inverse association between occupational physical activity and endometrial cancer, as have a number of previous studies (9, 20). It may be that the measures of occupational activity we used in this research were not sufficiently precise to identify the association. Indeed, our finding of apparent modification of the occupational sitting time variable by menopausal status was unexpected and may be due to the inherent limitations in using job codes to estimate physical activity levels. It is also

possible that occupational activity levels in this study were relatively homogenous and that the major differences in their overall activity patterns were determined by nonoccupational activity. We also observed a possible differential effect on the type of adolescent physical activity by menopausal status. That is, the effect of exercise in adolescence seemed to reduce risk for premenopausal but not postmenopausal women (Table 2, top), and the effect for total lifestyle activity reported in adolescence seemed stronger among postmenopausal compared with premenopausal women (see Results). There are several possible explanations for these findings. It may be that domain-specific activity levels differed by life period and age in these women. For example, 20.7% of premenopausal women

Table 2. Exercise in adolescence, adulthood, and in mutually exclusive life periods in relation to endometrial cancer risk, the Shanghai Endometrial Cancer Study (1997-2001)

	Cases/controls*	All women, OR <sup>†</sup> (95% CI)	Premenopausal, OR <sup>†</sup> (95% CI)	Postmenopausal, OR <sup>†</sup> (95% CI)
<b>Adolescence (MET-h/d/y)</b>				
None	679/666	1.00	1.00	1.00
0.1-0.9	57/46	1.17 (0.76-1.80)	1.24 (0.67-2.31)	1.00 (0.53-1.89)
1.0-1.7	20/46	0.39 (0.22-0.68)	0.50 (0.20-1.24)	0.35 (0.17-0.74)
1.8-3.5	43/41	0.89 (0.56-1.44)	0.91 (0.46-1.82)	0.85 (0.43-1.69)
3.6-17.6	33/47	0.63 (0.39-1.04)	0.37 (0.17-0.77)	1.17 (0.57-2.38)
<i>P</i> <sub>trend</sub>		0.02	0.02	0.42
<b>Adult (MET-h/d/y)</b>				
None	579/559	1.00	1.00	1.00
0.1-0.5	59/83	0.67 (0.46-0.97)	0.75 (0.36-1.54)	0.62 (0.40-0.98)
0.6-1.2	73/67	0.93 (0.64-1.36)	1.32 (0.58-2.96)	0.85 (0.55-1.31)
1.3-2.6	67/69	0.86 (0.58-1.26)	0.81 (0.35-1.85)	0.86 (0.56-1.34)
2.7-11.3	54/69	0.76 (0.50-1.13)	0.75 (0.32-1.75)	0.77 (0.48-1.22)
<i>P</i> <sub>trend</sub>		0.12	0.54	0.19
<b>Total (MET-h/d/y)</b>				
None	479/450	1.00	1.00	1.00
0.1-0.7	100/105	0.84 (0.60-1.16)	1.22 (0.68-2.19)	0.67 (0.45-1.00)
0.8-1.8	101/101	0.89 (0.64-1.24)	0.98 (0.55-1.78)	0.83 (0.56-1.24)
1.9-3.4	84/94	0.74 (0.53-1.05)	0.78 (0.43-1.40)	0.73 (0.48-1.13)
3.5-19.1	68/96	0.60 (0.42-0.86)	0.45 (0.25-0.83)	0.69 (0.44-1.10)
<i>P</i> <sub>trend</sub>		<0.01	0.02	0.06
<b>Mutually exclusive life periods</b>				
Never	479/450	1.00	1.00	1.00
Adolescent-only	100/109	0.79 (0.57-1.10)	0.81 (0.52-1.27)	0.76 (0.45-1.27)
Adult-only	200/216	0.81 (0.63-1.05)	0.99 (0.59-1.68)	0.76 (0.56-1.02)
Adolescent + adult	53/70	0.63 (0.42-0.95)	0.57 (0.28-1.17)	0.69 (0.41-1.14)
<i>P</i> <sub>trend</sub>		0.01	0.23	0.03

Abbreviation: 95% CI, 95% confidence interval.

\*n values for all cases and controls.

†Adjusted for age, age at menarche, menopausal status and age, number of pregnancies, oral contraceptive use, current smoking, ever drinking, family history of cancer, education, height, and body mass index.

**Table 3. Occupational physical activity and endometrial cancer risk, the Shanghai Endometrial Cancer Study (1997-2001)**

Job code indices	Cases/controls*	All women, OR <sup>†</sup> (95% CI)	Premenopausal, OR <sup>†</sup> (95% CI)	Postmenopausal, OR <sup>†</sup> (95% CI)
<b>Sitting</b>				
Q1 (less)	190/206	1.00	1.00	1.00
Q2	205/210	0.98 (0.73-1.33)	1.32 (0.81-2.14)	0.81 (0.55-1.19)
Q3	214/213	0.96 (0.71-1.31)	1.34 (0.83-2.18)	0.75 (0.50-1.14)
Q4 (more)	223/216	0.93 (0.67-1.30)	1.90 (1.04-3.49)	0.64 (0.42-0.99)
<i>P</i> <sub>trend</sub>		0.67	0.05	0.05
<b>Activity energy expenditure</b>				
Q1 (less)	213/212	1.00	1.00	1.00
Q2	209/213	0.80 (0.60-1.08)	0.67 (0.42-1.07)	0.96 (0.65-1.43)
Q3	221/203	1.05 (0.78-1.42)	0.95 (0.60-1.50)	1.15 (0.76-1.72)
Q4 (more)	189/203	0.86 (0.63-1.18)	0.75 (0.41-1.38)	0.89 (0.60-1.32)
<i>P</i> <sub>trend</sub>		0.70	0.57	0.68
<b>Self-rating of exertion</b>				
Q1 (less)	208/204	1.00	1.00	1.00
Q2	221/221	0.99 (0.74-1.32)	1.00 (0.63-1.57)	1.08 (0.73-1.61)
Q3	203/205	0.97 (0.72-1.32)	0.79 (0.49-1.26)	1.15 (0.77-1.74)
Q4 (more)	199/215	0.87 (0.64-1.20)	0.79 (0.46-1.37)	0.96 (0.64-1.44)
<i>P</i> <sub>trend</sub>		0.42	0.28	0.82
<b>Standing and walking (h/d)</b>				
0-1.6	339/332	1.00	1.00	1.00
1.7-3.3	218/212	0.96 (0.74-1.25)	0.92 (0.59-1.41)	0.96 (0.68-1.33)
3.4-5.5	144/177	0.81 (0.61-1.07)	0.85 (0.54-1.34)	0.78 (0.54-1.14)
5.6-13.3	131/123	1.09 (0.79-1.50)	1.20 (0.67-2.14)	1.03 (0.70-1.53)
<i>P</i> <sub>trend</sub>		0.78	0.93	0.70

Abbreviation: 95% CI, 95% confidence interval.

\**n* values for all cases and controls.

†Adjusted for age, age at menarche, menopausal status and age, number of pregnancies, oral contraceptive use, current smoking, ever drinking, family history of cancer, education, height, and body mass index.

reported exercise in adolescence but only 8.6% of postmenopausal women did so. In terms of the possible differential effect of adolescent lifestyle activities by menopausal status (OR<sub>premenopausal</sub>, 0.83; 95% CL, 0.49-1.40; OR<sub>postmenopausal</sub>, 0.51; 95% CL, 0.35-0.74), in detailed analyses, older women (postmenopausal) reported engaging in more lifestyle activity in adolescence than younger women (i.e., 7.0 versus

6.6 hours/d), although these differences were not significant (*P* = 0.10). Another plausible interpretation of these data is that the nonsignificant reduction in risk among premenopausal women is in fact a weaker effect. However, reduced sample size in stratified analyses, particularly for the smaller group of premenopausal women, limits our ability to make precise inference for stratified effects.


**Table 4. Lifestyle physical activity and endometrial cancer risk, the Shanghai Endometrial Cancer Study (1997-2001)**

	Adolescence		Adulthood	
	Cases/controls*	OR <sup>†</sup> (95% CI)	Cases/controls	OR <sup>†</sup> (95% CI)
<b>Household activity (h/d)</b>				
0	242/200	1.00	0-2	331/274
1	210/213	0.78 (0.59-1.05)	3	219/226
2	177/184	0.74 (0.55-1.00)	4	154/183
>3	199/247	0.67 (0.50-0.90)	>5	128/162
<i>P</i> <sub>trend</sub>		0.01		<0.01
<b>Walking for transportation (min/d)</b>				
0-25	140/151	1.00	0-26	210/182
26-52	238/242	1.00 (0.73-1.38)	27-55	236/223
53-65	245/240	0.97 (0.71-1.33)	56-60	223/242
>66	207/211	0.98 (0.71-1.35)	>61	163/199
<i>P</i> <sub>trend</sub>		0.84		<0.01
<b>Cycling for transportation (min/d)</b>				
No	805/814	1.00	0	648/667
Yes	25/31	0.77 (0.43-1.37)	2-29	40/27
			30-50	67/74
			>51	76/75
<i>P</i> <sub>trend</sub>		—		0.44
<b>Total lifestyle activity (MET-h/d)</b>				
0-3.63	229/184	1.00	0-7.29	217/189
3.64-6.20	240/245	0.71 (0.54-0.95)	7.30-9.83	286/232
6.21-8.80	166/172	0.72 (0.53-0.99)	9.84-12.75	145/213
>8.81	197/245	0.63 (0.46-0.84)	>12.76	184/212
<i>P</i> <sub>trend</sub>		<0.01		<0.01

Abbreviation: 95% CI, 95% confidence interval.

\**n* values for all cases and controls.

†Adjusted for age, age at menarche, menopausal status and age, number of pregnancies, oral contraceptive use, current smoking, ever drinking, family history of cancer, education, height, and body mass index.



**Figure 2.** Lifestyle activity and endometrial cancer risk by exercise participation in adulthood. Odds ratio was adjusted for age, age at menarche, menopausal status and age, number of pregnancies, oral contraceptive use, current smoking, ever drinking, family history of cancer, education, height, and body mass index.

Terry et al. (7) estimated that nongenetic factors, including physical activity, were responsible for nearly 80% of endometrial cancer cases in the Swedish Twins cohort study. Physical activity done at work, home, and for transportation has been associated with reductions in risk by about 30% to 40% in most (8, 9, 14, 15, 18, 20) but not all studies (11, 12, 23). A limitation of much of this literature has been the lack of detailed analyses focusing on the type and amount of wide variety of activities on risk. Quantitative estimates of the type and duration activity associated with reduced risk is needed to translate epidemiologic findings into meaningful information for public health officials to use in making evidence-based recommendations for cancer risk reduction (38).

For household activity and walking for transportation in adulthood, we observed graded inverse relationships between these activities and endometrial cancer risk, suggesting that benefit may be derived from modest amounts of activity. Women that reported >26 minutes/d of walking for transportation or >2 hours/d of household activity were at reduced risk. Importantly, we found that women with the lowest levels of household and transportation related activity but that participated in regular exercise were at about 35% lower risk. Similarly, nonexercisers with more active lifestyles were also at reduced risk.

Our finding that participation in exercise and sport activities reduce risk significantly was consistent with a number of other studies (7, 17, 18). The clear benefit from consistent participation across the life course in this research supports the idea that young women should be encouraged to participate in sport and exercise activities in youth and maintain this pattern of activity into middle and older ages. Our finding that fewer than 10% of women in this population reported this pattern of exercise over their lifetime (Fig. 1) suggests that the population-attributable risk for never exercisers would be substantial. We are aware of only one other study that has examined the effect of increasing physical activity over time on subsequent endometrial cancer risk. Moradi et al. (9), examining patterns of change in occupational activity, reported that women who moved from sedentary to highly active occupations were at similar risk as women who remained consistently in highly active occupations. Our finding that postmenopausal women who initiated exercise in adulthood were at a 25% lower risk of endometrial cancer (OR, 0.76; 95% CL, 0.56-1.02) than women who never exercised, suggests that it is possible that women can make changes in their physical activity patterns that will result in risk reduction. However, given the width of the confidence interval in this risk estimate, our results should be interpreted

with some caution. We and others have reported that changes in activity levels in later life can reduce breast cancer risk (39-41). Additional research that carefully evaluates changes in activity patterns over the life course is needed to extend our understanding of the potential for physical activity to be recognized as a truly *modifiable* risk factor for cancer.

In conclusion, these findings are consistent with the growing body of literature indicating that high levels of physical activity are associated with reduced risk for endometrial cancer. In this research, long-term exercise participation and high levels of common nonexercise activities were associated with 30% to 40% reductions in risk. Our findings for household activity and walking for transportation support the idea that a broad range of lower intensity activities may provide important benefit. Given the low-level exercise participation among middle-aged and older women in this and other populations (42, 43), this research has important public health consequences in highlighting the potential for both exercise and lifestyle activities to reduce endometrial cancer risk.

## Acknowledgments

We thank the field staff of the Shanghai Endometrial Cancer Study for their tireless and exceptional work in making this research possible.

## References

1. Ferlay J, Bray F, Pisani P, Parkin DM. GLOBOCAN 2000: cancer incidence, mortality and prevalence worldwide, version 1.0. IARC Cancer Base No. 5 ed. Lyon: IARC Press; 2001.
2. American Cancer Society. Cancer facts and figures, 2001. Atlanta (GA): American Cancer Society; 2001.
3. Jin F, Shu XO, Devesa SS, Zheng W, Blot WJ, Gao YT. Incidence trends for cancers of the breast, ovary, and corpus uteri in urban Shanghai, 1972-89. *Cancer Causes Control* 1993;4:355-60.
4. Jin F, Devesa SS, Chow WH, et al. Cancer incidence trends in urban Shanghai, 1972-1994: an update. *Int J Cancer* 1999;83:435-40.
5. Hu G, Hu G, Pekkarinen H, Hanninen O, Tian H, Jin R. Comparison of dietary and non-dietary risk factors in overweight and normal-weight Chinese adults. *Br J Nutr* 2002;88:91-7.
6. Paeratakul S, Popkin BM, Keyou G, Adair LS, Stevens J. Changes in diet and physical activity affect the body mass index of Chinese adults. *Int J Obes Relat Metab Disord* 1998;22:424-31.
7. Terry P, Baron JA, Weiderpass E, Yuen J, Lichtenstein P, Nyren O. Lifestyle and endometrial cancer risk: a cohort study from the Swedish Twin Registry. *Int J Cancer* 1999;82:38-42.
8. Furberg AS, Thune I. Metabolic abnormalities (hypertension, hyperglycemia and overweight), lifestyle (high energy intake and physical inactivity) and endometrial cancer risk in a Norwegian cohort. *Int J Cancer* 2003;104:669-76. Erratum in: *Int J Cancer* 2003 May 10;104:799.
9. Moradi T, Nyren O, Bergstrom R, et al. Risk for endometrial cancer in relation to occupational physical activity: a nationwide cohort study in Sweden. *Int J Cancer* 1998;76:665-70.

10. Schouten LJ, Goldbohm RA, van den Brandt PA. Anthropometry, physical activity, and endometrial cancer risk: results from the Netherlands cohort study. *J Natl Cancer Inst* 2004;96:1635-8.
11. Pukkala E, Poskiparta M, Apter D, Vihko V. Life-long physical activity and cancer risk among Finnish female teachers. *Eur J Cancer Prev* 1993; 2:369-76.
12. Colbert LH, Lacey JV Jr, Schairer C, Albert P, Schatzkin A, Albanes D. Physical activity and risk of endometrial cancer in a prospective cohort study (United States). *Cancer Causes Control* 2003;14:559-67.
13. Zheng W, Shu XO, McLaughlin JK, Gao WH, Gao YT, Blott WJ. Occupational physical activity and the incidence of cancer of the breast, corpus uteri, and ovary in Shanghai. *Cancer* 1993;71:3620-4.
14. Sturgeon SR, Brinton LA, Berman ML, et al. Past and present physical activity and endometrial cancer risk. *Br J Cancer* 1993;68:584-9.
15. Goodman MT, Hankin JH, Wilkens LR, et al. Diet, body size, physical activity, and the risk of endometrial cancer. *Cancer Res* 1997;57:5077-85.
16. Olson SH, Vena JE, Dorn JP, et al. Exercise, occupational activity, and risk of endometrial cancer. *Ann Epidemiol* 1997;7:46-53.
17. Moradi T, Weiderpass E, Signorello LB, Persson I, Nyren O, Adami HO. Physical activity and postmenopausal endometrial cancer risk (Sweden). *Cancer Causes Control* 2000;11:829-37.
18. Littman AJ, Voigt LF, Beresford SA, Weiss NS. Recreational physical activity and endometrial cancer risk. *Am J Epidemiol* 2001;154:924-33.
19. Salazar-Martinez E, Lazcano-Ponce EC, Lira-Lira GG, et al. Case-control study of diabetes, obesity, physical activity and risk of endometrial cancer among Mexican women. *Cancer Causes Control* 2000;11:707-11.
20. Levi F, La Vecchia C, Negri E, Franceschi S. Selected physical activities and the risk of endometrial cancer. *Br J Cancer* 1993;67:846-51.
21. Kalandidi A, Tzonou A, Lipworth L, Gamatsi I, Filippa D, Trichopoulos D. A case-control study of endometrial cancer in relation to reproductive, somatometric, and life-style variables. *Oncology* 1996;53:354-9.
22. Hirose K, Tajima K, Hamajima N, et al. Subsite (cervix/endometrium)-specific risk and protective factors in uterus cancer. *Jpn J Cancer Res* 1996;87:1001-9.
23. Shu XO, Hatch MC, Zheng W, Gao YT, Brinton LA. Physical activity and risk of endometrial cancer. *Epidemiology* 1993;4:342-9.
24. Furberg AS, Thune I. Energy intake, energy expenditure and BMI influence the risk of endometrial cancer in a prospective study in Norway. *IARC Sci Publ* 2002;156:247-8.
25. Xu WH, Matthews CE, Xiang YB, et al. Effect of adiposity and fat distribution on endometrial cancer risk in Shanghai women. *Cancer Epidemiol Biomarkers Prev*. In review 2004.
26. Ainsworth B, Haskell W, Whitt M, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* 2000;32:S498-516.
27. Ainsworth B, Haskell W, Leon A, et al. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc* 1993;25:71-80.
28. Matthews CE, Shu XO, Gong Y, et al. Reproducibility and validity of the Shanghai women's health study physical activity questionnaire. *Am J Epidemiol* 2003;158:1114-22.
29. Hosmer DW, Lemeshow S. *Applied logistic regression*. New York: John Wiley and Sons; 1989.
30. Friedenreich CM. Physical activity and cancer prevention: from observational to intervention research. *Cancer Epidemiol Biomarkers Prev* 2001;10:287-301.
31. Kaaks R, Lukanova A, Kurzer MS. Obesity, endogenous hormones, and endometrial cancer risk: a synthetic review. *Cancer Epidemiol Biomarkers Prev* 2002;11:1531-43.
32. Irwin ML, Mayer-Davis EJ, Addy CL, et al. Moderate-intensity physical activity and fasting insulin levels in women: the Cross-Cultural Activity Participation Study. *Diabetes Care* 2000;23:449-54.
33. Mayer-Davis EJ, D'Agostino R, Karter AJ, et al. Intensity and amount of physical activity in relation to insulin sensitivity: the Insulin Resistance Atherosclerosis Study. *JAMA* 1998;279:669-74.
34. Ivy JL, Zderic TW, Fogt DL. Prevention and treatment of non-insulin dependent diabetes. *Exerc Sport Sci Rev* 1999;27:1-35.
35. IARC. *Weight control and physical activity*. Handbook. 6th ed. Lyon: IARC; 2002.
36. McTiernan A, Tworoger SS, Ulrich CM, et al. Effect of exercise on serum estrogens in postmenopausal women: a 12-month randomized clinical trial. *Cancer Res* 2004;64:2923-8.
37. Ainsworth BE. Issues in the assessment of physical activity in women. *Res Q Exerc Sport* 2000;71:37-42.
38. Byers T, Nestle M, McTiernan A, et al. American Cancer Society guidelines on nutrition and physical activity for cancer prevention: reducing the risk for cancer with healthy food choices and physical activity. *CA Cancer J Clin* 2002;52:92-119.
39. Matthews CE, Shu XO, Jin F, et al. Lifetime physical activity and breast cancer risk in the Shanghai Breast Cancer Study. *Br J Cancer* 2001;84: 994-1001.
40. Friedenreich CM, Courneya KS, Bryant HE. Influence of physical activity in different age and life periods on the risk of breast cancer. *Epidemiology* 2001;12:604-12.
41. Verloop J, Rookus MA, van der Kooy K, van Leeuwen FE. Physical activity and breast cancer risk in women aged 20-54 years. *J Natl Cancer Inst* 2000;92:128-35.
42. U.S.D.H.H.S. Physical activity trends-United States, 1990-1998. *MMWR Morb Mortal Wkly Rep* 2001;50:166-9.
43. United States Department of Health and Human Services. Prevalence of physical activity, including lifestyle activities among adults-United States, 2000-2001. *MMWR Morb Mortal Wkly Rep* 2003;52:764-9.

## Physical Activity and Risk of Endometrial Cancer: A Report from the Shanghai Endometrial Cancer Study

Charles E. Matthews, Wang Hong Xu, Wei Zheng, et al.

*Cancer Epidemiol Biomarkers Prev* 2005;14:779-785.

**Updated version** Access the most recent version of this article at:  
<http://cebp.aacrjournals.org/content/14/4/779>

**Cited articles** This article cites 38 articles, 5 of which you can access for free at:  
<http://cebp.aacrjournals.org/content/14/4/779.full#ref-list-1>

**Citing articles** This article has been cited by 5 HighWire-hosted articles. Access the articles at:  
<http://cebp.aacrjournals.org/content/14/4/779.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://cebp.aacrjournals.org/content/14/4/779>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.