

Ki-67 Expression in Benign Breast Ductal Cells Obtained by Random Periareolar Fine Needle Aspiration

Qamar J. Khan, Bruce F. Kimler, Julie Clark, Trina Metheny, Carola M. Zalles, and Carol J. Fabian

University of Kansas Medical Center, Kansas City, Kansas

Abstract

Ki-67 expression in ductal cells obtained by random periareolar fine needle aspiration (RPFNA) is currently being used as a response biomarker in phase II breast cancer chemoprevention trials; however, Ki-67 in RPFNA has not been well studied as a risk predictor for cancer, which would support its use as a response indicator. We examined the expression of Ki-67 in RPFNA specimens with hyperplasia \pm atypia obtained from 147 women at high risk for development of breast cancer. Median Ki-67 was 1.4% (range 0-24%). Ki-67 was higher in specimens from women <50 versus those ≥ 50 (median 2% versus 0.6%; $P = 0.006$) and from premenopausal women versus postmenopausal women

($P = 0.037$); however, hormone replacement therapy (predominately low-dose estrogen without progestins) had no effect. By univariate analysis, Ki-67 was positively correlated with ductal cell number ($P = 0.001$) and hyperplasia with atypia ($P = 0.007$). By multivariable analysis, the proportion of ductal cells expressing Ki-67 was again predicted by cell number, which, in turn, was predicted by cytologic atypia. The association of Ki-67 expression with cytologic atypia, a known risk factor for development of breast cancer, provides preliminary justification for its use as a response biomarker in phase II chemoprevention trials. (Cancer Epidemiol Biomarkers Prev 2005;14(4):786-9)

Introduction

Risk biomarkers that are subject to modulation are used as response indicators in early prevention trials (1-3). We have previously shown that specimens obtained from high-risk women by random periareolar fine needle aspiration (RPFNA) and assessed as hyperplasia with atypia are associated with a 5-fold increase in the short-term risk of subsequent development of ductal carcinoma *in situ* or invasive cancer (4). RPFNA cytomorphology is being used as a primary response end point in phase II prevention trials; however, intra- and inter-interpretive variance can substantially increase study subject number requirements. Quantitative biomarkers, such as Ki-67, minimize interpretive variance, reducing the number of subjects required for a prevention study (3). The rationale for use of Ki-67 in proliferative breast disease as a risk biomarker stems from the hypothesis that the majority of breast cancers evolve from proliferative lesions (5-8), and a cross-sectional study of Ki-67 in benign biopsies where Ki-67 was higher in hyperplastic foci from women who later developed cancer than from women who had not developed cancer (medians 3.8% versus 0.8%; ref. 9). The observation that early reduction in Ki-67 predicts clinical response to tamoxifen in cancer treatment trials provides the rationale for use of Ki-67 as a response biomarker (10, 11). In the absence of prospective trials assessing Ki-67 in benign breast disease and subsequent breast cancer incidence, a positive association between RPFNA atypia and Ki-67 would provide preliminary support for use of Ki-67 as a risk and response biomarker in early prevention trials (1, 7).

Materials and Methods

Study Cohort. The study cohort consisted of high-risk women undergoing baseline eligibility assessment for one of several phase II chemoprevention trials at the University of Kansas Medical Center Breast Cancer Prevention Center. Women eligible for RPFNA were those whose 5-year Gail risk was $\geq 1.7\%$, whose relative risk of developing breast cancer was at least five times that of the general population for their age based on a 5-year Gail risk assessment, whose prior breast biopsy had exhibited atypical hyperplasia or lobular carcinoma *in situ*, who were known to carry a BRCA1/2 mutation, or who had a prior contralateral treated breast cancer. Women must have a normal mammogram at the time of aspiration and have been at least 6 months from change in hormone replacement therapy (HRT), ingestion of any selective estrogen receptor modulator or aromatase inhibitor, and 1 year from pregnancy, lactation, or any prior chemotherapy.

Eligibility Criteria. Ki-67 assessment was attempted only for specimens estimated to have sufficient (≥ 500) ductal cells on each of the slides designated for cytomorphology and Ki-67.

Gail Risk Calculation. The projected probability of developing *in situ* or invasive cancer at 5 years was calculated for each subject according to the Gail model estimated at the time of aspiration (<http://bcra.nci.nih.gov/brc/>; ref. 12).

RPFNA Procedure. Subjects with a prior history of breast cancer (invasive or ductal carcinoma *in situ*) had RPFNA done only on the uninvolved breast. RPFNA was done as previously described (4). For premenopausal women, all RPFNAs were done on days 1 to 12 (follicular portion) of the menstrual cycle.

Specimen Handling. Material from all breast aspiration sites for each woman was pooled in a 15 mL conical tube containing 9 mL Cytolyt (Cytoc, Boxborough, MA) and 1 mL of 10% neutral buffered formalin. Conical tubes were placed on a Verimix Rocker (Barnstead International, Dubuque, IA) at low speed. Cells were then washed with Cytolyt, processed to a pellet, placed in PreservCyt (Cytoc) for 48 hours, and then

Received 3/29/04; revised 11/12/04; accepted 12/6/04.

Grant support: Contract NO1-CN-15135 from Chemoprevention Branch, Cancer Prevention Research Program-Cancer Control, Division of Cancer Prevention, National Cancer Institute, NIH, and grant BCTR0100732 from Susan G. Komen Breast Cancer Research Foundation.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Carol J. Fabian, University of Kansas Medical Center, 3901 Rainbow Boulevard, Kansas City, KS 66160. Phone: 913-588-7791; Fax: 913-588-3679. E-mail: cfabian@kumc.edu

Copyright © 2005 American Association for Cancer Research.

Table 1. Correlation of Ki-67 values with demographic and cytologic characteristics

Characteristic	n	Median (%)	Range (%)	P*
Total	147	1.4	0-23.6	
Age				
≤50 y	89	2.0	0-23.6	0.006
>50 y	58	0.6	0-16.0	
Menopause/HRT				
Pre	67	2.0	0-15.8	0.037
Post, HRT	47	1.0	0-23.6	
Post, no HRT	33	1.0	0-12.0	
5-y Gail†				
<Median	71	2.2	0-23.6	0.039
>Median of 2.2%	71	1.0	0-16	
Cytology				
Hyperplasia	110	1.1	0-23.6	0.007
Atypia	37	2.8	0-16.0	
Cytology				
Benign	73	1.0	0-23.6	0.028
Indeterminate/atypia	74	1.6	0-16.0	
Masood score				
≤12	37	0.8	0-12.0	0.025
13	29	1.2	0-11.0	
14	47	1.4	0-23.6	
≥15	34	2.6	0-13.4	
Cell number per slide				
<1,000	77	1.0	0-23.6	0.001
1,000-4,999	41	2.2	0-13.4	
≥5,000	29	2.4	0-16.0	

*Mann-Whitney test or Kruskal-Wallis test for distribution of Ki-67 values.

†Gail risk could not be calculated for five subjects.

processed to three to four slides using a standard Thin Prep 2000 (Cytec) nongynecologic protocol. At least two slides were Papanicolaou stained, with one used for morphology and one for Ki-67 staining.

RNase-Free Papanicolaou Staining Procedure. Slides for both cytomorphology and Ki-67 were Papanicolaou stained under RNase-free conditions with hematoxylin, OG-6, and EA-65 (all from Richard Allen Scientific, Kalamazoo, MI) and were prepared on the ThinPrep Processor.

Cytomorphology Assessment. Cytomorphology was assessed by a single cytopathologist (C.M. Zalles) and classified by three different methods: traditional (13), Masood semi-quantitative index (14), and the 1996 National Cancer Institute Consensus Panel Criteria (15). The traditional method classifies preparations as nonproliferative, apocrine metaplasia, epithelial hyperplasia, hyperplasia with atypia, or carcinoma (13). The Masood index scoring system assigns a score of 1 to 4 points to each of six morphologic characteristics: cellular arrangement, cellular pleomorphism, prevalence of myoepithelial cells, anisonucleosis, nucleoli, and chromatin clumping. A score of 6 to 10 is generally associated with nonproliferative specimens, 11 to 14 with hyperplasia without atypia, 15 to 18 with hyperplasia with atypia, and 19 to 24 with malignancy (14). National Cancer Institute Consensus Panel Criteria classify specimens as benign, atypical/indeterminate, suspicious/probably malignant, malignant, and unsatisfactory (15). Cytologic assessments were made without knowledge of the results of the Ki-67 assessment.

Ki-67 Assessment. Only slides containing >500 epithelial cells were processed and stained for Ki-67. Antigen retrieval was done with 10 mmol/L citrate buffer (pH 6) in a BioCare (Walnut Creek, CA) decloaking chamber for 2 minutes at 120°C. Slides were then stained with MIB-1 monoclonal antibody (M7240; DakoCytomation, Carpinteria, CA) at a 1:20 dilution in a Dako Autostainer. A categorical estimate of the number of ductal epithelial cells present on the Ki-67 slide was made as 500 to 1,000; 1,000 to 5,000; or >5,000.

Hyperplastic cell clusters were preferentially assessed and the number of cells with unequivocal nuclear staining out of 500 cells assessed was recorded. Slides were hand scored by two readers and assessments recorded on separate case report forms; however, in case of a difference between the two readers, the scores were averaged. As we have previously reported, agreement between two readers using this method was excellent (Cronbach's $\alpha = 0.99$; ref. 16).

Statistical Analysis. Frequencies of categorical variables were assessed using χ^2 analysis. Continuous variables were assessed using the Mann-Whitney nonparametric test. Multivariate analyses were done using stepwise linear regression.

Results

One hundred and forty-seven high-risk women who underwent RPFNA between March 2003 and August 2003 and had sufficient ductal cells (≥ 500) in the slide designated for Ki-67 were included in this analysis. Median age was 49 years (range 28-78 years). Eighty-nine (60%) women were <50 years of age. Sixty-seven (45%) women were premenopausal and 80 (55%) were postmenopausal. Among postmenopausal women, 47 (59%) women were on some form of HRT, including 26 on estrogen alone, 4 on estrogen plus testosterone, 11 on estrogen plus progestins (3 with testosterone as well), 5 on poorly absorbed vaginal estrogens, and 1 on a soy supplement.

Ki-67 Expression. For the entire cohort, the median level of Ki-67 expression was 1.4% (range 0-23.6%) as shown in Table 1. There was excellent agreement and low interobserver variance between the two readers for the 147 specimens ($R^2 = 0.99$).

Correlation of Ki-67 Expression with Demographic Factors and Gail Risk. Median Ki-67 was 2.0% in specimens from women <50 years of age compared with 0.6% in specimens from women ≥ 50 years ($P = 0.006$, Mann-Whitney test). Median Ki-67 was 2% among specimens from premenopausal women compared with 1% in specimens from postmenopausal women ($P = 0.037$). There was no difference in Ki-67 expression for specimens from postmenopausal women receiving HRT versus those not on HRT, but only 23% of women on HRT were on estrogen plus a progestin. Lastly, there was a marginally significant difference for Ki-67 expression between specimens from women with a 5-year Gail risk below the median of 2.2% versus those with a Gail risk above the median (median Ki-67 values of 2.2% versus 1.0%, $P = 0.039$).




Figure 1. Frequency of specimens exhibiting Ki-67 expression greater than or equal to specific percent positive values as a function of cytology (hyperplasia versus hyperplasia with atypia). This shows that a greater proportion of specimens classified as hyperplasia with atypia exhibited specific levels of Ki-67 staining than did specimens classified as hyperplasia.

Correlation of Ki-67 Expression with Cytologic Morphology and Cell Number

Traditional Cytologic Assessment. Ki-67 was higher in women classified as having RPFNA epithelial hyperplasia with atypia than those with hyperplasia without atypia (Fig. 1). Median Ki-67 was 2.8% in specimens characterized as hyperplasia with atypia compared with 1.1% in specimens of hyperplasia without atypia ($P = 0.007$; Table 1).

Masood Score and Consensus Panel. Median Masood score for the study cohort was 14, with a range of 10 to 17. As shown in Table 1, median Ki-67 was 0.8% for Masood scores ≤ 12 , and 1.2%, 1.4%, and 2.6% for Masood scores of 13, 14, and ≥ 15 , respectively (Fig. 2). Globally, these values are statistically significantly different ($P = 0.025$ by Kruskal-Wallis test) with a specific pairwise comparison being significant for ≤ 12 versus ≥ 15 ($P = 0.002$; Fig. 2). Specimens classified as indeterminate/atypia had a significantly higher Ki-67 than those who were benign ($P = 0.028$).

Cell Number on Slide. For the 77 specimens with 500 to 1,000 cells in the Ki-67 slide, the median Ki-67 was 1.2%; for the 41 specimens with 1,000 to 5,000 cells, Ki-67 was 2.2%; and for the 29 specimens with $>5,000$ cells, Ki-67 was 2.4%. The distribution of Ki-67 values was statistically significantly different between the three groups ($P = 0.001$).

Despite the general correlations of Ki-67 with cytomorphology, there was still considerable heterogeneity. For example, 21% of specimens with both evidence of atypia and $>5,000$ cells per slide exhibited no Ki-67 staining.

Prediction of Ki-67 Expression: Multivariable Analysis. Regression analysis was conducted to ascertain which of the above variables, associated with Ki-67 by univariate analysis (Table 1), would best predict for Ki-67 expression. The variables examined in the regression analysis were age, menopause status, HRT use, Gail risk, cell category number, and cytomorphology assessment by traditional, National Cancer Institute Consensus Panel criteria, and Masood score. Only cell number category was a significant ($P = 0.004$) determinant of Ki-67 in the multivariable analysis. However, if cell number was excluded from the regression analysis,

traditional cytomorphology category was the only variable to be associated with Ki-67 expression ($P = 0.027$). Cytomorphology and cell count were highly associated in univariate analysis ($P = 0.002$).

Discussion

The positive association between RPFNA hyperplasia with atypia, an established risk biomarker, and Ki-67 expression provides preliminary evidence for use of Ki-67 in benign breast tissue as a risk indicator and response biomarker. A prospective study with adequate numbers of both premenopausal and postmenopausal women would be required for definitive evidence that proliferation (Ki-67) is indeed a risk factor that might complement cytomorphology assessment.

Our values for Ki-67 in cytology specimens with hyperplasia are similar to those reported by other authors for histologic preparations (9, 17-20). Further, our study results were similar to those reported by others in which higher Ki-67 values were reported in premenopausal than in postmenopausal women (19). All of our specimens from premenopausal women were obtained during the follicular phase. Higher values might be anticipated had premenopausal women with proliferative breast changes been sampled during the luteal phase of the menstrual cycle (20). Given the small size of this pilot study, it is not clear whether the association between cytomorphology and Ki-67 holds for both premenopausal and postmenopausal women. Perhaps the greatest use for Ki-67 assessment in benign breast tissue will not be for risk stratification but as a response biomarker in early phase prevention trials (2, 3). For the majority of our ongoing pilot and phase II prevention trials in which Ki-67 is the primary response biomarker, we now require a 2% level of Ki-67 staining for a woman to enter the treatment portion of the study. This seems logical given that cytologic atypia is associated with a median value Ki-67 of 2.8%. Our prior studies indicate that 70% of high-risk women will exhibit proliferative cytology and our current study indicates that 40% of high-risk women with proliferative cytology (57% with atypia, 27% with hyperplasia alone) can be expected to exhibit this $\geq 2\%$ level of Ki-67 staining.

In summary, our observation that increased Ki-67 expression is associated with the established risk biomarker of RPFNA evidence of cytologic atypia provides preliminary support for its current use as a risk biomarker and response end point in early breast cancer chemoprevention trials.

References

- Kelloff GJ, Boone CW, Crowell JA, et al. Risk biomarkers and current strategies for cancer chemoprevention. *J Cell Biochem* 1996;25:1-14.
- Fabian CJ, Kimler BF, Elledge RM, Grizzle WE, Beenken SW, Ward JH. Models for early chemoprevention trials in breast cancer. *Hematol Oncol Clin North Am* 1998;12:993-1017.
- Fabian CJ, Kimler BF, Mayo MS, Grizzle WE, Masood S, Ursin G. Clinical approaches to discovering and testing new breast cancer prevention drugs. In: Kelloff GJ, Hawk ET, Sigman CC, editors. *Cancer chemoprevention, Volume 2: Strategies for cancer chemoprevention*. Totowa, NJ: Humana Press; 2005. p. 213-37.
- Fabian CJ, Kimler BF, Zalles CM, et al. Short-term breast cancer prediction by random periareolar fine-needle aspiration cytology and the Gail risk model. *J Natl Cancer Inst* 2000;92:1217-27.
- Dupont WD, Page DL. Risk factors for breast cancer in women with proliferative breast disease. *N Engl J Med* 1985;312:146-51.
- Allred DC, Medina D. Introduction: models of premalignant breast disease. *J Mammary Gland Biol Neoplasia* 2000;5:339-40.
- Kelloff GJ, Sigman CC, Johnson KM, et al. Perspectives on surrogate end points in the development of drugs that reduce the risk of cancer. *Cancer Epidemiol Biomarkers Prev* 2000;9:127-37.
- Allred DC, Mohsin SK, Fuqua SA. Histological and biological evolution of human premalignant breast disease. *Endocr Relat Cancer* 2001; 8:47-61.




Figure 2. Top, relationship between Ki-67 expression and Masood semiquantitative cytology index score. A score of 6 to 10 is generally associated with nonproliferative specimens, 11 to 14 with hyperplasia without atypia, and 15 to 18 with hyperplasia with atypia (12). Bottom, relationship between Ki-67 expression and estimated number of ductal cells per cytology slide.

9. Shaaban AM, Sloane JP, West CR, Foster CS. Breast cancer risk in usual ductal hyperplasia is defined by estrogen receptor α and Ki 67 expression. *Am J Pathol* 2002;160:597–604.
10. Markis A, Powles TJ, Allred DC, et al. Changes in hormone receptors and proliferation markers in tamoxifen treated breast cancer patients and the relationship with response. *Breast Cancer Res Treat* 1998;48:11–20.
11. Chang J, Powles TJ, Allred DC, et al. Prediction of clinical outcome from primary tamoxifen by expression of biologic markers in breast cancer patients. *Clin Cancer Res* 2000;6:616–21.
12. Gail MH, Brinton LA, Byar DP, et al. Projecting individualized probabilities of developing breast cancer for white females who are being examined annually. *J Natl Cancer Inst* 1989;81:1879–86.
13. Zalles C, Kimler BF, Kamel S, McKittrick R, Fabian CJ. Cytologic patterns in random aspirates from women at high and low risk for breast cancer. *Breast J* 1995;1:343–9.
14. Masood S, Frykberg ER, McLellan GL, Scalapino MC, Mitchum DG, Bullard JB. Prospective evaluation of radiologically directed fine needle aspiration biopsy of nonpalpable breast lesions. *Cancer* 1990;66:1480–7.
15. The uniform approach to breast fine-needle aspiration biopsy. National Cancer Institute Fine-Needle Aspiration of Breast Workshop Subcommittees. *Diagn Cytopathol* 1997;16:295–311.
16. Fabian CJ, Clark J, Metheny T, et al. Proportion of benign ductal epithelial cells expressing Ki-67/MIB 1 in random periareolar fine needle aspirates from women at high risk of breast cancer [abstract R1878]. *Proc Am Assoc Cancer Res* 2003;44:427.
17. Harper Wynne C, Sacks NP, Shenton K, et al. Effects of aromatase inhibitor letrozole on normal breast epithelial cell proliferation and metabolic indices in postmenopausal women. A pilot study for breast cancer prevention. *Cancer Epidemiol Biomarkers Prev* 2002;11:614–21.
18. Shoker BS, Jarvis C, Clarke RB, et al. Estrogen receptor-positive proliferating cells in the normal and precancerous breast. *Am J Pathol* 1999;155:1811–5.
19. Potten CS, Watson RJ, Williams GT, et al. The effect of age and menstrual cycle upon proliferative activity of the normal human breast. *Br J Cancer* 1988;58:163–70.
20. Soderqvist G, Isaksson E, von Schoultz B, Carlstrom K, Tani E, Skoog L. Proliferation of breast epithelial cells in healthy women during the menstrual cycle. *Am J Obstet Gynecol* 1997;176:123–8.

Ki-67 Expression in Benign Breast Ductal Cells Obtained by Random Periareolar Fine Needle Aspiration

Qamar J. Khan, Bruce F. Kimler, Julie Clark, et al.

Cancer Epidemiol Biomarkers Prev 2005;14:786-789.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/14/4/786>

Cited articles This article cites 18 articles, 4 of which you can access for free at:
<http://cebp.aacrjournals.org/content/14/4/786.full#ref-list-1>

Citing articles This article has been cited by 12 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/14/4/786.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and
Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications
Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/14/4/786>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's
(CCC)
Rightslink site.