

# The *MTHFR* 1298A>C Polymorphism and Genomic DNA Methylation in Human Lymphocytes

Simonetta Friso,<sup>1</sup> Domenico Girelli,<sup>1</sup> Elisabetta Trabetti,<sup>2</sup> Oliviero Olivieri,<sup>1</sup> Patrizia Guarini,<sup>1</sup> Pier Franco Pignatti,<sup>2</sup> Roberto Corrocher,<sup>1</sup> and Sang-Woon Choi<sup>3</sup>

Departments of <sup>1</sup>Clinical and Experimental Medicine and <sup>2</sup>Biology and Genetics, University of Verona, Italy; and <sup>3</sup>Vitamin Carcinogenesis Laboratory, Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts

## Abstract

Methylenetetrahydrofolate reductase (*MTHFR*) balances the pool of folate coenzymes in one-carbon metabolism for DNA synthesis and methylation, both implicated in carcinogenesis. Epidemiologic studies have shown that two functional polymorphisms in *MTHFR* gene, 677C>T and 1298A>C, are related to increased cancer risk. We aimed to analyze lymphocyte DNA from 198 subjects to evaluate the *MTHFR* 1298A>C polymorphism and folate status affecting genomic DNA methylation as a possible mechanism underlying the relationship between *MTHFR* polymorphisms and cancer susceptibility. Carriers of the 1298AA wild-type genotype showed lower genomic DNA methylation compared with 1298AC or 1298CC genotypes [3.72 versus 8.59 or 6.79 ng 5-methyl-2'-deoxycytidine (5-mCyt)/ $\mu$ g DNA,  $P < 0.0001$  and  $P = 0.007$ , respectively]. When DNA methylation was evaluated according to plasma folate status, only 1298AA with low folate levels revealed diminished DNA methylation ( $P < 0.0001$ ). Moreover,

when the two *MTHFR* polymorphisms were concomitantly evaluated at the low folate status, DNA methylation was reduced only in 1298AA/677TT compared with 1298AA/677CC (3.11 versus 7.29 ng 5-mCyt/ $\mu$ g DNA,  $P = 0.001$ ) and 1298CC/677CC genotypes (3.11 versus 7.14 ng 5-mCyt/ $\mu$ g DNA,  $P = 0.004$ ). However, the high prevalence of 677TT mutants within the 1298AA group (79%) and the similar biochemical features of 1298AA/677CC and 1298CC/677CC combined genotypes suggest that the gene-nutrient interaction affecting DNA methylation in 1298AA is mainly due to the coexistence of the 677TT genotype and that the 1298A>C polymorphism may convey its protective effect not through this interaction but through another pathway in one-carbon metabolism. Further mechanistic studies are warranted to investigate how single polymorphisms as well as *MTHFR* combined genotypes exert their effect on cancer susceptibility. (Cancer Epidemiol Biomarkers Prev 2005;14(4):938–43)

## Introduction

The altered properties of lymphoid neoplastic cells have been traditionally attributed to changes in the gene sequence or structure such as point mutations, gene deletions/rearrangements, or double strand breaks with subsequent increased risk of chromosomal translocations. Epigenetic abnormalities, widely described in several solid tumor malignancies, have been similarly found in hematopoietic cancer cells, suggesting that epigenetic phenomena can play a substantial role also in the carcinogenic process affecting this particular tissue (1). DNA methylation is the most important epigenetic feature of eukaryotic DNA for the regulation of gene expression and the maintenance of genome integrity (2, 3). Preserving the patterns of DNA methylation is a major feature of DNA for the epigenetic control of genome stability and the regulation of gene transcription (4). An altered profile of DNA methylation is a frequent finding in cancer diseases, indicating that an aberrant regulation of the mechanisms of DNA methylation is crucial for the understanding of mechanisms of carcinogenesis (1, 5) and may constitute a

good target for future therapies (5, 6). Abnormal genomewide DNA methylation is a phenomenon known to lead to chromosome instability and damage (7), genetic abnormalities that are frequently described in malignancies, mostly those of the lymphoid tissue.

A crucial role for the control of DNA methylation as well as DNA synthesis and repair mechanisms is played by the provision of adequate pools of methyl groups through the functional activity of methylenetetrahydrofolate reductase (*MTHFR*) in one-carbon metabolism. The availability of methyl groups for DNA methylation reactions is mainly regulated by the activity of *MTHFR*, which is indeed a fundamental enzyme in one-carbon metabolism. Its substrate, 5,10-methylenetetrahydrofolate, is necessary for thymidylate and purine synthesis and its enzymatic product regulates the folate-dependent *de novo* synthesis of 5-methyltetrahydrofolate, the prominent circulatory form of folate and the precursor of *S*-adenosylmethionine, universal methyl donor for several biological methylation reactions, including that of DNA (8). We recently described that the different availability of methylated forms of folate in homozygous mutant carriers of the *MTHFR* 677C>T polymorphism (9) affects genomic DNA methylation through gene-nutrient interaction (10, 11). We observed that carriers of the 677TT genotype have a diminished level of DNA methylation compared with those with the 677CC wild-type genotype but only the 677TT subjects with low levels of folate accounted for the diminished DNA methylation, thus indicating that the *MTHFR* 677C>T polymorphism influences DNA methylation level through an interaction with folate status (11, 12).

A second common mutation in the *MTHFR* gene, 1298A>C, was reported. First described in association with ovarian carcinoma (13), it was subsequently investigated for its association with risk of neural tube defects, a group of diseases

Received 8/13/04; revised 12/21/04; accepted 1/3/05.

**Grant support:** National Funding of the Ministry of University, Scientific, and Technologic Research (O. Olivieri and R. Corrocher), and the Ministry of Health of Italy (R. Corrocher); the Fondazione Cassa di Risparmio di Verona Vicenza Belluno e Ancona, Verona, Italy (S. Friso); the Regione Veneto, Ricerca Finalizzata, Venezia, Italy (D. Girelli, O. Olivieri, and R. Corrocher); and National Cancer Institute grant RO3 CA96004-02 (S.W. Choi). This material is based on work supported by the U.S. Department of Agriculture, under agreement no. 581950-9-001. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Requests for reprints:** Simonetta Friso, Department of Clinical and Experimental Medicine, Policlinico Giambattista Rossi, Piazza Ludovico Antonio Scuro, 10, 37134 Verona, Italy. Phone: 39-045-580111; Fax: 39-045-580111. E-mail: simonetta.friso@univr.it

Copyright © 2005 American Association for Cancer Research.

closely linked to one-carbon metabolism functions (14). However, whether or not these associations are ultimately due to a reduced activity of the mutant *MTHFR 1298A>C* enzyme is yet controversial (14-18). Most recently, polymorphic sites in folate metabolizing genes, particularly those affecting *MTHFR* activity, have been associated with a reduced risk of adult (19-22) and childhood (23-25) acute lymphoblastic leukemia development as well as with modulation of the response to chemotherapy (26). Furthermore, epidemiologic studies have shown an association between *MTHFR* polymorphisms and susceptibility to cancer affecting also other tissues (27-30). Because of the fundamental role of folate in nucleic acid synthesis and repair systems (31) as well as DNA methylation (12), alterations in this metabolic pathway seem an elective candidate target for the understanding of folate-related carcinogenesis with potential diagnostic and therapeutic implications (25, 26). Furthermore, it is known that *MTHFR 677C>T* and folate, independently or interactively, are important determinants of genome stability (7), cell proliferation, and chromosomal breakage, particularly in lymphoid cells (32-35).

Recent reports indicate that the *MTHFR 1298A>C* polymorphism is associated with a decreased risk of acute lymphoblastic leukemia in adults and children carrying the mutant allele either in heterozygosity (*1298AC*) or homozygosity (*1298CC*; refs. 19, 24). Furthermore, it has been reported that individuals with the combined *1298AA/677TT* or *1298CC/677CC* genotype show respectively a 2.5- and 3.3-fold reduction of risk for childhood acute lymphoblastic leukemia (25). Although several hypotheses have been proposed, there is yet no clear evidence for a link between this polymorphism and DNA methylation status.

We designed the present study precisely with the aim of evaluating the interaction between the *MTHFR 1298A>C* mutation and folate in lymphocyte DNA methylation status as a potential mechanism by which this polymorphism exerts a protective effect, particularly on the risk of lymphoid cell malignancies, either alone or in combination with the *MTHFR 677C>T*.

## Materials and Methods

**Study Population.** We studied 198 unrelated, age- and sex-matched subjects from a single geographic area and recruited those referred to the Section of Internal Medicine of the Clinical and Experimental Medicine Department and to the Institute of Cardiology and Cardiovascular Surgery of the University of Verona (Verona, Italy), as previously described (36). Subjects for whom a complete set of biochemical data pertaining to folate-homocysteine metabolism (plasma folate levels, vitamin B12, and total plasma homocysteine concentrations) and a complete clinical history were available, were selected for the present study on gene-nutrient interaction between folate and *MTHFR* polymorphisms. Subjects were considered as a whole with no distinction referred to their pathologic status, and excluded from the study were those with conditions known to influence homocysteine levels such as those subjects taking folate or vitamin B<sub>12</sub> or any multivitamin supplementation, those with any major acute illness, and those using drugs interfering with folate metabolism (i.e., penicillamine, anticonvulsants, and methotrexate). Informed consent was obtained from all the subjects and the study received the approval of our Institutional Review Board.

**Biochemical Analyses.** Samples of venous blood were drawn from each subject after an overnight fast. DNA was extracted from peripheral lymphocytes using a phenol/chloroform/isoamyl alcohol protocol. All the laboratory

methods for biochemical analyses have been done as previously described (36). Levels of total homocysteine (tHcy) were measured by high-performance liquid chromatography with fluorescent detection according to Araki and Sako (37). Plasma folate was measured by an automated chemiluminescence method (Chiron Diagnostics, East Walpole, MA).

**Methylenetetrahydrofolate Reductase Genotyping.** For the *MTHFR 677C>T* single nucleotide polymorphism, we used primer sets, PCR-RFLP method, and nomenclature as previously described by Frosst et al. (38), after optimizing the conditions in our laboratory. The *1298A>C* mutation analysis was done as described by Weisberg et al. (ref. 15; i.e., by PCR followed by *Fnu4HI* endonuclease restriction). Positive and negative mutation controls were always included in each gel during the study for the genotyping of both the *677C>T* and *1298A>C* mutations (15).

**DNA Methylation Measurement.** To measure genomic DNA methylation, we used a recently developed liquid chromatography electrospray ionization mass spectrometry method that allows to define DNA methylation status as the absolute amount (in nanograms) of 5-methyl-2'-deoxycytidine (mCyt) per microgram of DNA (11, 39). Briefly, 1 µg of DNA was hydrolyzed by sequential digestion with three nucleases. DNA hydrolyzates were then injected onto an analytic column and subsequently separated by reversed-phase high-performance liquid chromatography in isocratic mode. The four major DNA bases and 5-methyl-2'-deoxycytidine were resolved and eluted in a short time run. The identification of 5-methyl-2'-deoxycytidine was obtained by spectral analysis of chromatographic peaks by isotope-labeled compounds as internal standard (39).

**Statistical Analysis.** All computations were done by using the Systat software 10.0 program for Windows (2000, SPSS, Inc., Chicago, IL). The distribution of continuous variables in groups is expressed as mean ± SD. Analysis was done with log-transformed data for all skewed variables including DNA methylation. Geometric means (antilogarithms of the transformed means) are presented, and 95% confidence intervals are calculated using the transformed values and displayed as the antilogarithms of the transformed data. Statistical significance for differences in continuous variables was tested by Student's unpaired *t* test or by AN(C)OVA with Tukey's post hoc comparison of the means when appropriate. Categorical variables were analyzed using a  $\chi^2$  test or linear regression analysis when appropriate. Statistical significance refers to a two-tailed analysis where *P* < 0.05.

## Results

Each end point was evaluated for the three *MTHFR 1298A>C* polymorphism genotypes (*1298AA*, *1298AC*, and *1298CC*; Table 1; Fig. 1) as well as for the three different combined genotype groups for *1298A>C* and *677C>T* polymorphisms (*1298AA/677TT*, *1298AA/677CC*, and *1298CC/677CC*; Table 2; Fig. 2).

### Frequency of Each *MTHFR* Polymorphism and Combined *MTHFR* Genotype Distribution

***MTHFR 1298A>C* Polymorphism.** The overall allele frequency of the *1298C* allele in the studied population was 0.33 and the genotype frequency was as follows: *1298AA*, 46% (*n* = 91); *1298AC*, 33% (*n* = 65); and *1298CC*, 21% (*n* = 42).

**Combination of *MTHFR 1298A>C* and *677C>T* Polymorphisms.** Among 91 *1298AA* subjects, 19 (21%) carried a *677CC* genotype and 72 (79%) carried a *677TT* genotype. The *677T* and *1298C* alleles only occurred in *trans*; no individual presented a *1298CC/677TT* genotype configuration.

**Table 1. Biochemical features of the study subjects according to MTHFR 1298A>C genotypes**

	1298AA, N = 91	1298AC, N = 65	1298CC, N = 42	P (AA vs AC)	P (AA vs CC)	P (AC vs CC)
Plasma folate (nmol/L)	10.90 (9.99-11.89)	12.19 (11.03-13.44)	12.40 (10.92-14.08)	0.094	0.097	0.828
Vitamin B <sub>12</sub> (pmol/L)	448 (404-495)	378 (346-413)	428 (380-482)	0.013	0.591	0.097
Vitamin B <sub>6</sub> (nmol//L)	28.16 (24.87-31.88)	28.47 (25.00-32.42)	29.10 (23.49-36.08)	0.900	0.772	0.840
tHcy (μmol/L)	18.71 (16.77-20.86)	13.97 (12.95-15.08)	14.45 (12.91-16.19)	<0.0001	0.001	0.603
Genomic DNA methylation (ng mCyt/μg DNA)	3.72 (2.73-5.07)	8.59 (6.83-10.77)	6.79 (4.96-9.28)	<0.0001	0.007	0.226

NOTE: Plasma folate, vitamin B<sub>12</sub>, vitamin B<sub>6</sub>, tHcy, and genomic DNA methylation are presented as geometric means (antilogarithms of the transformed means) and 95% confidence intervals are reported in parentheses with two-tailed *P* values. Statistical difference was evaluated by Student's *t* test except when differently indicated. Statistical difference was evaluated by  $\chi^2$  test.

### Plasma Folate Concentrations

**MTHFR 1298A>C Polymorphism.** As shown in Table 1, plasma folate concentrations did not differ among the three genotype groups. However, when the subjects carrying the 1298C allele either in hetero- or homozygosity were considered together, plasma folate concentration within the 1298AA group was significantly lower than that found in those carrying at least one 1298C allele (10.87 versus 12.22 nmol/L, *P* < 0.05).

**Combined MTHFR 1298A>C and 677C>T Polymorphisms.** Plasma folate concentration was lower in the 1298AA/677TT group compared with the 1298CC/677CC group (10.73 versus 12.62 nmol/L, *P* = 0.043; Table 2). Comparisons of 1298AA/677TT with 1298AA/677CC and 1298CC/677CC groups did not show any significant difference (Table 2).

### Total Plasma Homocysteine Concentrations


**MTHFR 1298A>C Polymorphism.** As shown in Table 1, the concentration of tHcy in the 1298AA group was significantly higher than that in either 1298AC or 1298CC group (18.71 versus 13.97 μmol/L and 18.71 versus 14.45 μmol/L, *P* < 0.0001 and *P* = 0.001, respectively). The difference was significant even when the carriers of the 1298C allele were considered

together (1298AA versus 1298AC/1298CC; 18.71 versus 14.16 μmol/L, *P* < 0.0001). When folate levels were taken into consideration, the 1298AA subjects with low folate concentrations (i.e., below the median value, 12 nmol/L) showed a significantly higher tHcy concentration compared with that of 1298AA individuals with adequate folate levels (i.e., above the median value, 12 nmol/L; 22.08 versus 13.18 μmol/L, *P* < 0.0001). Within the 1298CC genotype carrier group, those with low plasma folate levels also had higher tHcy concentrations compared with those with high folate concentrations (19.46 versus 13.93 μmol/L, *P* < 0.0001). However, there was a significant difference in tHcy concentrations between 1298AA and 1298CC genotypes (19.29 versus 15.89 μmol/L, *P* = 0.002) at the low level of folate, but no significant difference between 1298AA and 1298AC (19.29 versus 16.27, *P* = 0.11). Such difference was no longer present if the comparison was made between subjects with levels of folate above the median (13.63 versus 12.62 μmol/L, *P* = 0.321, and 13.63 versus 13.53 μmol/L, *P* = 0.891, for the comparison between 1298AA versus 1298CC and 1298AA versus 1298AC, respectively).

**Combination of MTHFR 1298A>C and 677C>T Polymorphisms.** Total plasma homocysteine concentration in the 1298AA/677TT group was significantly higher than that in the 1298AA/677CC and 1298CC/677CC group (19.57 versus 15.79 μmol/L, *P* = 0.045, and 19.57 versus 14.41 μmol/L, *P* = 0.001, respectively; Table 2). Even when the 1298AA/677TT was compared with the 1298AA/677CC and 1298CC/677CC groups taken together, the difference remained statistically significant (19.57 versus 14.86 μmol/L, *P* = 0.001). However, no difference was detected for the comparison of the 1298AA/677CC with the 1298CC/677CC group (15.79 versus 14.41 μmol/L, *P* = 0.358; Table 2). When we analyzed the data according to plasma folate concentrations, the 1298AA/677TT group showed a significant difference between low- and high-folate groups (23.87 versus 13.09 μmol/L, *P* < 0.0001). A difference according to low or high plasma folate levels was also detected within the 1298CC/677CC group (16.87 versus 12.55 μmol/L, *P* = 0.012). However, no differences were found within the 1298AA/677CC group (16.03 versus 13.49 μmol/L, *P* = 0.257).

### Genomic DNA Methylation

**MTHFR 1298A>C Polymorphism.** DNA methylation status within MTHFR 1298AA group was significantly decreased compared with that of either 1298AC (3.72 versus 8.59 ng mCyt/μg DNA, *P* < 0.0001) or 1298CC (3.72 versus 6.79 ng mCyt/μg DNA, *P* = 0.007; Table 1). DNA methylation was significantly decreased in the MTHFR 1298AA group compared with the 1298AC and 1298CC genotypes considered together (3.72 versus 7.82 ng mCyt/μg DNA, *P* < 0.0001). There was no significant difference between the 1298AC and 1298CC groups for DNA methylation status. However, when plasma folate levels were taken into account, the reduced DNA methylation in the 1298AA group was evident only for those subjects with low plasma folate concentrations (below the



**Figure 1.** Genomic DNA methylation according to MTHFR 1298A>C genotype and low or high plasma folate levels (i.e., below or above the median, 12 nmol/L). Statistical difference was evaluated by Student's *t* test. Within the 1298AA group, the levels of DNA methylation were significantly reduced in the low-folate group compared with the high-folate group (2.87 versus 8.59 ng mCyt/μg DNA, *P* = 0.001). No differences were detected within both the 1298AC and 1298CC genotypes according to folate levels (*P* = 0.771 and *P* = 0.135, respectively). Considering only the three genotype groups at low folate status, the levels of DNA methylation were significantly lower in 1298AA versus 1298AC and 1298AA versus 1298CC (*P* = 0.0001 and *P* = 0.021, respectively). Among high-folate groups, there were no statistically significant differences for 1298AA versus 1298AC (*P* = 0.98) and 1298AA versus 1298CC (*P* = 0.084).

**Table 2. Biochemical data according to the combined presence of *MTHFR* 1298A>C and 677C>T genotypes**

	Group 1 (1298AA/677TT), N = 72	Group 2 (1298AA/677CC), N = 19	Group 3 (1298CC/677CC), N = 42	P (group 1 vs group 2)	P (group 1 vs group 3)	P (group 2 vs group 3)
Plasma folate (nmol/L)	10.73 (9.73-11.85)	11.52 (9.31-14.25)	12.62 (11.15-14.30)	0.513	0.043	0.440
Vitamin B <sub>12</sub> (pmol/L)	437 (391-487)	487 (369-642)	433 (383-488)	0.450	0.900	0.419
Vitamin B <sub>6</sub> (nmol//L)	30.53 (26.62-35.02)	21.03 (16.05-27.52)	28.59 (23.01-35.51)	0.016	0.607	0.090
tHcy (μmol/L)	19.57 (17.18-22.28)	15.79 (13.34-18.70)	14.41 (12.83-16.18)	0.045	0.001	0.358
Genomic DNA methylation (ng mCyt/μg DNA)	3.11 (2.16-4.49)	7.29 (4.64-11.44)	7.14 (5.26-9.67)	0.004	0.001	0.936

NOTE: Plasma folate, vitamin B12, vitamin B6, tHcy, and genomic DNA methylation are presented as geometric means (antilogarithms of the transformed means) and 95% confidence intervals are reported in parentheses with two-tailed *P* values. Statistical difference was evaluated by Student's *t* test except when differently indicated. Statistical difference was evaluated by  $\chi^2$  test.

median value, i.e., 12 nmol/L), whereas under conditions of adequate plasma folate, the levels of genomic DNA methylation were similar to that of the other two *MTHFR* genotypes (Fig. 1).

To further evaluate the relationship between folate and *MTHFR* genotypes, we did additional analyses. After dividing the population sample into tertiles of plasma folate according to the three different 1298A>C genotypes, we observed that only among the individuals carrying the 1298AA genotype was there a graded effect of folate in determining levels of DNA methylation (2.39 ng mCyt/μg DNA for the lowest tertile, 5.33 ng mCyt/μg for the intermediate tertile, and 7.54 ng mCyt/μg DNA for the highest tertile, *P* for trend = 0.005). Among the carriers of the 1298AC or 1298CC genotype, however, DNA methylation status was not different across the increasing tertiles of plasma folate.


**Combination of *MTHFR* 1298A>C and 677C>T Polymorphisms.** Genomic DNA methylation status of the *MTHFR* 1298AA/677TT group was significantly decreased compared with that of 1298AA/677CC (3.11 versus 7.29 ng mCyt/μg DNA, *P* = 0.001) and 1298CC/677CC groups (3.11 versus 7.14 ng mCyt/μg DNA, *P* = 0.0001; Table 2). There was no significant difference between the 1298AA/677CC and 1298CC/677CC groups (Table 2). However, when considering plasma folate levels, the reduced DNA methylation was evident only for the

1298AA/677TT subjects with low plasma folate concentrations (1298AA/677TT, 2.44; 1298AA/677CC, 5.57; and 1298CC/677CC, 7.53 ng mCyt/μg DNA), whereas under conditions of adequate plasma folate, genomic DNA methylation status was similar to the other two *MTHFR* genotype groups (1298AA/677TT, 8.31 ng mCyt/μg DNA; 1298AA/677CC, 9.49 ng mCyt/μg DNA; and 1298CC/677CC, 6.69 ng mCyt/μg DNA; Fig. 2). No statistically significant difference for DNA methylation status was observed within the 1298AA/677CC and 1298CC/677CC groups between low and high levels of folate (5.57 versus 9.49 ng mCyt/μg DNA and 7.53 versus 6.69 ng mCyt/μg DNA, *P* = 0.67 and *P* = 0.69, respectively; Fig. 2).

## Discussion

Epidemiologic studies recently observed an association of polymorphic variants of the *MTHFR* gene with reduced risk of adult and childhood acute lymphoblastic leukemia (19, 22-25) and with the modulation of risk of other forms of cancer (27-30, 40, 41). The *MTHFR* 677TT homozygote mutant genotype has been consistently reported to have a protective effect against the risk of cancer of the colorectum (27, 42) and of the lymphoid tissue in several studies (19, 20, 22-25). The role of *MTHFR* polymorphisms in modulating cancer risk, however, is strongly associated with folate status, so that the protective effect of the 677TT genotype, under adequate folate conditions, turns to a situation of elevated risk of cancer of the colorectum among *MTHFR* 677TT carriers with low folate status or intake (27, 43). On the other hand, although some studies observed a protective effect also for the *MTHFR* 1298CC mutant genotype, results in this regard are conflicting (22, 23). Because the function of *MTHFR* is essential for providing methyl groups for DNA methylation (27), it is highly plausible that enzymatic variants due to functional polymorphisms may alter this major epigenetic feature of DNA, patterns of which greatly affect carcinogenesis (1, 3, 5). Different from 677C>T, the biochemical effect of the 1298A>C mutation on *MTHFR* function is still controversial (14-18), and its role in the risk of cancer is therefore more questionable (40, 41).

In the present study, we sought to evaluate the *MTHFR* 1298A>C polymorphism interaction with folate status and the subsequent effects on genomic methylation in human lymphocyte DNA as a potential mechanism underlying the association between the *MTHFR* 1298A>C polymorphism and the risk of cancer. We observed that lymphocyte DNA has reduced genomic DNA methylation compared with that of carriers of either the 1298AC or 1298CC genotype. This finding referred only to those subjects whose plasma folate concentrations are low with no difference in DNA methylation among the three 1298A>C genotypes when stratified for high levels of folate. A further evidence of such gene-nutrient interaction was the detection of a graded effect of



**Figure 2.** Genomic DNA methylation (ng mCyt/μg DNA) according to combined *MTHFR* 1298A>C and 677C>T genotypes and low or high plasma folate levels (i.e., below or above the median, 12 nmol/L). Statistical difference was evaluated by Student's *t* test. The reduced DNA methylation was evident only for 1298AA/677TT genotype carriers with low plasma folate, whereas among 1298AA/677TT subjects under adequate conditions of plasma folate, the levels of genomic DNA methylation were similar to that of the other two *MTHFR* genotype groups. The difference in DNA methylation between low- and high-folate subjects in the 1298AA/677TT group was statistically significant (2.44 versus 8.31 ng mCyt/μg DNA, *P* < 0.0001).

folate in determining DNA methylation, as shown by the increasing DNA methylation levels across increasing tertiles of plasma folate concentrations. This gene-nutrient interaction phenomenon affecting DNA methylation is supported also by the observation of a similar effect of the 1298AA genotype on tHcy levels, as shown by the higher tHcy in the 1298AA group compared with carriers of at least one 1298C allele, as similarly described for the 677TT genotypes (36). Because tHcy is a product of the intracellular hydrolysis of S-adenosylhomocysteine, which is known to be an inhibitor of DNA methylation, tHcy levels could be also considered as a marker of tissue DNA methylation status, as reported by Yi et al. (44).

The most crucial conflict in the present observation, however, is that the MTHFR 1298AA wild-type genotype for the MTHFR 1298A>C mutation shows a model of interaction with folate similar to that of the mutant MTHFR 677TT genotype. When considered by combined polymorphic genotypes, the 1298AA/677TT group showed decreased plasma folate concentration, increased tHcy, and decreased DNA methylation status compared with the 1298AA/677CC and 1298CC/677CC groups, and even when folate concentration was taken into account, only the 1298AA/677TT genotype group showed a decreased genomic DNA methylation when plasma folate levels are low. Furthermore, when we compared the 1298AA/677CC and 1298CC/677CC groups, to avoid any possible influence of the mutant 677TT genotype, there was no difference in plasma folate, tHcy, and lymphocyte genomic DNA methylation. These data suggest that the MTHFR 1298A>C polymorphism does not impair the enzyme function significantly enough to affect the pathway of homocysteine remethylation and biological methylation functions, thus substantiating previous findings showing no effect of the 1298A>C polymorphism *per se* in determining tHcy (14, 18), and leading to the understanding of a minor effect of the 1298A>C mutation on this pathway of one-carbon metabolism as a consequence of reduced MTHFR enzyme function.

Plausible explanations for the similar features observed for the 1298AA and 677TT arise from the analysis of the distribution of the 677C>T polymorphism among carriers of the 1298A>C polymorphism. Noteworthy, we observed that among 1298AA carriers, 79% had a 677TT genotype, and all of the 1298AC and 1298CC carriers had a 677CC genotype. Furthermore, the 1298C and 677T alleles only occurred in *trans* in our population, consistent with several previous reports on the very rare occurrence of the two polymorphisms on the same allele (14, 15, 18). This combined genotype setting, therefore, may explain the similar pattern of DNA methylation for the 1298AA and 677TT genotypes, and that of the 1298CC with the coexistence of the 677CC genotype. Indeed, such observations confirm as well the population specificity of these genotypes and could give reason for the controversial association and potentially ethnic-related association of MTHFR genotypes with acute lymphoblastic leukemia in some reports (22, 45).

Consistent with the results of the present study were the reports on the structural and biochemical characteristics of the two different enzyme variants, MTHFR 677C>T and 1298A>C (46, 47). The 677C>T polymorphism lies at the base of binding site for the MTHFR cofactor, flavin adenine dinucleotide, and has been shown to affect the MTHFR enzyme activity more significantly than the 1298A>C variant (14, 46). Individuals carrying the 677TT genotype show a 30% *in vitro* enzyme activity compared with the 677CC wild-type (38, 46). The MTHFR 1298A>C variant, instead, has been shown to have biochemical properties that are not markedly distinguishable from those of the 1298AA wild-type genotype (47). Further reports on the biochemical relevance of the 1298A>C mutation showed also that lymphocytes from individuals carrying the

mutant 1298CC genotype have ~60% specific wild-type MTHFR activity *in vitro* compared with the 1298AA wild-type (14), although such a decrease did not seem sufficient to affect tHcy levels (14, 15, 18). Thus, considering the strong effect of the MTHFR 677C>T variant on determining the availability of methylated forms of folate and on remethylation pathway (9, 11), we can speculate that the genomic DNA methylation pattern in 1298AA genotypes is mostly driven by the concomitant presence of the 677TT genotype. The effect of the 1298AA/677TT genotype also is likely due primarily to the high portion of 677TT in the combined genotype group, and the effect of both 1298AA/677CC and 1298CC/677CC combined genotypes is mainly related to the simultaneous presence of the 677CC genotype.

Therefore, it may be of crucial importance for future epidemiologic studies to evaluate both MTHFR single polymorphisms as well as combined genotype configurations in the MTHFR gene (48) and to simultaneously consider environmental factors such as folate status (49). Indeed, whereas a strong gene-nutrient interaction affects the MTHFR 677C>T polymorphism in determining DNA methylation, the mildly decreased activity due to MTHFR 1298A>C mutation may affect a different pathway in one-carbon metabolism, such as that of DNA synthesis, and therefore accounts for a different mechanism for the reduced risk of leukemia in carriers of the 1298C allele. Specific mechanistic studies are needed to clarify the molecular bases of such associations.

The evaluation of MTHFR common mutations and folate status should be considered in the view of an individually tailored assessment of cancer risk because it may as well lead to the ultimate goal of providing, in the near future, a better clinical and preventive approach (50).

## Acknowledgments

We are indebted to all subjects who consented to participate to this study. We are especially thankful to Diego Minguzzi and Patrizia Pattini from the Department of Clinical and Experimental Medicine at the University of Verona School of Medicine, Italy, for their excellent technical assistance and valuable contribution in data collection.

## References

- Melki JR, Baylin SJ. DNA methylation changes in leukaemia. *Semin Cancer Biol* 2002;12:347–57.
- Issa JP, Baylin SB. Epigenetics and human disease. *Nat Med* 1996 Mar;2:281–2.
- Laird PW. The power and the promise of DNA methylation markers. *Nat Rev Cancer* 2003;3:253–66.
- Wolffe AP, Matzke MA. Epigenetics: regulation through repression. *Science* 1999;286:481–6.
- Esteller M, Herman JG. Cancer as an epigenetic disease: DNA methylation and chromatin alterations in human tumours. *J Pathol* 2002;196:1–7.
- Egger G, Liang G, Aparicio A, Jones PA. Epigenetics in human disease and prospects for epigenetic therapy. *Nature* 2004;429:457–63.
- Eden A, Gaudet F, Waghmare A, Jaenisch R. Chromosomal instability and tumors promoted by DNA hypomethylation. *Science* 2003;300:455.
- Green R, Miller JW. Folate deficiency beyond megaloblastic anemia: hyperhomocysteinemia and other manifestations of dysfunctional folate status. *Semin Hematol* 1999;36:47–64.
- Bagley PJ, Selhub J. A common mutation in the methylenetetrahydrofolate reductase gene is associated with an accumulation of formylated tetrahydrofolates in red blood cells. *Proc Natl Acad Sci U S A* 1998;95:13217–20.
- Stern LL, Mason JB, Selhub J, Choi SW. Genomic DNA hypomethylation, a characteristic of most cancers, is present in peripheral leukocytes of individuals who are homozygous for the C677T polymorphism in the methylenetetrahydrofolate reductase gene. *Cancer Epidemiol Biomarkers Prev* 2000;9:849–53.
- Friso S, Choi SW, Girelli D, et al. A common mutation in the 5,10-methylenetetrahydrofolate reductase gene affects genomic DNA methylation through an interaction with folate status. *Proc Natl Acad Sci U S A* 2002;99:5606–11.
- Friso S, Choi SW. Gene-nutrient interactions and DNA methylation. *J Nutr* 2002;132:2382–7S.
- Viel A, Dall'Agnese L, Simone F, et al. Loss of heterozygosity at the 5,10-methylenetetrahydrofolate reductase locus in human ovarian carcinomas. *Br J Cancer* 1997;75:1105–10.

14. van der Put NMJ, Gabreels F, Stevens EMB, et al. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural tube defects? *Am J Hum Genet* 1998;62:1044–51.
15. Weisberg I, Tran P, Christensen B, Sibani S, Rozen R. A second genetic polymorphism in methylenetetrahydrofolate reductase (MTHFR) associated with decreased enzyme activity. *Mol Genet Metab* 1998;64:169–72.
16. Chango A, Boisson F, Barbé F, et al. The effect of 677C→T and 1298A→C mutations on plasma homocysteine and 5,10-methylenetetrahydrofolate reductase activity in healthy subjects. *Br J Nutr* 2000;83:593–6.
17. Friedman G, Goldschmidt N, Friedlander Y, et al. A common mutation A1298C in human methylenetetrahydrofolate reductase gene: association with plasma total homocysteine and folate concentrations. *J Nutr* 1999;129:1656–61.
18. Friso S, Girelli D, Trabetti E, et al. A1298C methylenetetrahydrofolate reductase mutation and coronary artery disease: relationships with C677T polymorphism and homocysteine/folate metabolism. *Clin Exp Med* 2002;2:7–12.
19. Skibola CF, Smith MT, Kane E, et al. Polymorphisms in the methylenetetrahydrofolate reductase gene are associated with susceptibility to acute leukemia in adults. *Proc Natl Acad Sci U S A* 1999;96:12810–5.
20. Matsuo K, Suzuki R, Hamajima N, et al. Association between polymorphisms of folate- and methionine-metabolizing enzymes and susceptibility to malignant lymphoma. *Blood* 2001;97:3205–9.
21. Skibola CF, Smith MT, Hubbard A, et al. Polymorphisms in the thymidylate synthase and serine hydroxymethyltransferase genes and risk of adult acute lymphocytic leukemia. *Blood* 2002;99:3786–91.
22. Gemmati D, Ongaro A, Scapoli GL, et al. Common gene polymorphisms in the metabolic folate and methylation pathway and the risk of acute lymphoblastic leukemia and non-Hodgkin's lymphoma in adults. *Cancer Epidemiol Biomarkers Prev* 2004;13:787–94.
23. Franco RF, Simoes BP, Tone LG, Gabellini SM, Zago MA, Falcao RP. The methylenetetrahydrofolate reductase C677T gene polymorphism decreases the risk of childhood acute lymphocytic leukaemia. *Br J Haematol* 2001;115:616–8.
24. Wiemels JL, Smith RN, Taylor GM, Eden OB, Alexander FE, Greaves MF; United Kingdom Childhood Cancer Study investigators. Methylenetetrahydrofolate reductase (MTHFR) polymorphisms and risk of molecularly defined subtypes of childhood acute leukemia. *Proc Natl Acad Sci U S A* 2001;98:4004–9.
25. Krajcinovic M, Lamothe S, Labuda D, et al. Role of MTHFR genetic polymorphisms in the susceptibility to childhood acute lymphoblastic leukemia. *Blood* 2004;103:252–7.
26. Ulrich CM, Yasui Y, Storb R, et al. Pharmacogenetics of methotrexate: toxicity among marrow transplantation patients varies with the methylenetetrahydrofolate reductase C677T polymorphism. *Blood* 2001;98:231–4.
27. Choi SW, Mason JB. Folate and carcinogenesis: an integrated scheme. *J Nutr* 2000;130:129–32.
28. Shen H, Spitz MR, Wang LE, Hong WK, Wei Q. Polymorphisms of methylenetetrahydrofolate reductase and risk of lung cancer: a case-control study. *Cancer Epidemiol Biomarkers Prev* 2001;10:397–401.
29. Song C, Xing D, Tan W, Wei Q, Lin D. Methylenetetrahydrofolate reductase polymorphisms increase risk of esophageal squamous cell carcinoma in a Chinese population. *Cancer Res* 2001;61:3272–5.
30. Lin J, Spitz MR, Wang Y, et al. Polymorphisms of folate metabolic genes and susceptibility to bladder cancer: a case-control study. *Carcinogenesis* 2004;25:1639–47.
31. Blount BC, Mack MM, Wehr CM, et al. Folate deficiency causes uracil misincorporation into human DNA and chromosome breakage: implications for cancer and neuronal damage. *Proc Natl Acad Sci U S A* 1997;94:3290–5.
32. Duthie SJ. Folic acid deficiency and cancer: mechanisms of DNA instability. *Br Med Bull* 1999;55:578–92.
33. Crott JW, Mashiyama ST, Ames BN, Fenech M. The effect of folic acid deficiency and MTHFR C677T polymorphism on chromosome damage in human lymphocytes *in vitro*. *Cancer Epidemiol Biomarkers Prev* 2001;10:1089–96.
34. Duthie SJ, Narayanan S, Brand GM, Pirie L, Grant G. Impact of folate deficiency on DNA stability. *J Nutr* 2002;132:2444–9S.
35. Kimura M, Umegaki K, Higuchi M, Thomas P, Fenech M. Methylenetetrahydrofolate reductase C677T polymorphism, folic acid and riboflavin are important determinants of genome stability in cultured human lymphocytes. *J Nutr* 2004 Jan;134:48–56.
36. Girelli D, Friso S, Trabetti E, et al. Methylenetetrahydrofolate reductase C677T mutation, plasma homocysteine, and folate in subjects from northern Italy with or without angiographically documented severe coronary atherosclerotic disease: evidence for an important genetic-environmental interaction. *Blood* 1998;91:4158–63.
37. Araki A, Sako Y. Determination of free and total homocysteine in human plasma by high-performance liquid chromatography with fluorescence detection. *J Chromatogr* 1987;422:43–52.
38. Frosst P, Blom HJ, Milos R, et al. Identification of a candidate genetic risk factor for vascular disease: A common mutation at the methylenetetrahydrofolate reductase locus. *Nat Genet* 1995;10:111–3.
39. Friso S, Choi SW, Dolnikowski GG, Selhub J. A method to assess genomic DNA methylation using high-performance liquid chromatography/electrospray ionization mass spectrometry. *Anal Chem* 2002;74:4526–31.
40. Shrubsole MJ, Gao YT, Cai Q, et al. MTHFR polymorphisms, dietary folate intake, and breast cancer risk: results from the Shanghai Breast Cancer Study. *Cancer Epidemiol Biomarkers Prev* 2004;13:190–6.
41. Curtin K, Bigler J, Slattery ML, Caan B, Potter JD, Ulrich CM. MTHFR C677T and A1298C polymorphisms: diet, estrogen, and risk of colon cancer. *Cancer Epidemiol Biomarkers Prev* 2004;13:285–92.
42. Chen J, Giovannucci EL, Hunter DJ. MTHFR polymorphism, methyl-replete diets and the risk of colorectal carcinoma and adenoma among U.S. men and women: an example of gene-environment interactions in colorectal tumorigenesis. *J Nutr* 1999;129:560–4S.
43. Potter JD. Methyl supply, methyl metabolizing enzymes and colorectal neoplasia. *J Nutr* 2002;132:2410–2S.
44. Yi P, Melnyk S, Pogribna M, Pogribny IP, Hine RJ, James SJ. Increase in plasma homocysteine associated with parallel increases in plasma S-adenosylhomocysteine and lymphocyte DNA hypomethylation. *J Biol Chem* 2000;275:29318–23.
45. Chiusolo P, Reddicono G, Cimino G, et al. Methylenetetrahydrofolate reductase genotypes do not play a role in acute lymphoblastic leukemia pathogenesis in the Italian population. *Haematologica* 2004;89:139–44.
46. Guenther BD, Sheppard CA, Tran P, Rozen R, Matthews RG, Ludwig ML. The structure and properties of methylenetetrahydrofolate reductase from *Escherichia coli* suggest how folate ameliorates human hyperhomocysteinemia. *Nat Struct Biol* 1999;6:359–65.
47. Yamada K, Chen Z, Rozen R, Matthews RG. Effects of common polymorphisms on the properties of recombinant human methylenetetrahydrofolate reductase. *Proc Natl Acad Sci U S A* 2001;98:14853–8.
48. Skibola CF, Forrest MS, Coppede F, et al. Polymorphisms and haplotypes in folate metabolizing genes and risk of non-Hodgkin lymphoma. *Blood* 2004 Oct 1; 104:2155–62 [Epub ahead of print, 2004 Jun 15].
49. Robien K, Ulrich CM. 5,10-Methylenetetrahydrofolate reductase polymorphisms and leukemia risk: a HuGE minireview. *Am J Epidemiol* 2003;157:571–82.
50. Ulrich CM, Robien K, McLeod HL. Cancer pharmacogenetics: polymorphisms, pathways and beyond. *Nat Rev Cancer* 2003;3:912–20.

# Cancer Epidemiology, Biomarkers & Prevention

**AACR** American Association  
for Cancer Research

## The *MTHFR* 1298A>C Polymorphism and Genomic DNA Methylation in Human Lymphocytes

Simonetta Friso, Domenico Girelli, Elisabetta Trabetti, et al.

*Cancer Epidemiol Biomarkers Prev* 2005;14:938-943.

**Updated version** Access the most recent version of this article at:  
<http://cebp.aacrjournals.org/content/14/4/938>

**Cited articles** This article cites 48 articles, 25 of which you can access for free at:  
<http://cebp.aacrjournals.org/content/14/4/938.full#ref-list-1>

**Citing articles** This article has been cited by 6 HighWire-hosted articles. Access the articles at:  
<http://cebp.aacrjournals.org/content/14/4/938.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://cebp.aacrjournals.org/content/14/4/938>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.