

## Short Communication

# Ethnic Disparity in the Relationship between Obesity and Plasma Insulin-Like Growth Factors: The Multiethnic Cohort

Katherine DeLellis Henderson,<sup>1</sup> Michael I. Goran,<sup>1</sup> Laurence N. Kolonel,<sup>2</sup> Brian E. Henderson,<sup>1</sup> and Loïc Le Marchand<sup>2</sup>

<sup>1</sup>Keck School of Medicine, University of Southern California, Los Angeles, California and <sup>2</sup>Cancer Research Center of Hawaii, University of Hawaii, Honolulu, Hawaii

## Abstract

Previous studies on the relationship between obesity and circulating insulin-like growth factor (IGF) hormones show inconsistent findings and have not considered the possibility of racial/ethnic-specific differences that may exist. We therefore examined the relationship between obesity status [as measured by body mass index (BMI)] and plasma levels of the IGF proteins, IGF-I, IGF-binding protein 3 (IGFBP-3), and the molar ratio of IGF-I/IGFBP-3 in Whites, African Americans, Latinos, Japanese Americans, and Native Hawaiians from the ongoing Hawaii and Los Angeles Multiethnic Cohort Study. We measured plasma IGF-I and IGFBP-3 by ELISA in a random sample of 811 Multiethnic Cohort participants (53% male, age range = 47-82 at blood draw). In a multivariate regression of IGF-I levels, we found a statistically significant interaction between race/ethnicity and obesity status ( $P = 0.005$ ).

Plasma IGF-I levels declined with increasing BMI most dramatically in Latinos and Japanese. This decline was attenuated in Whites and absent in African-American and Native Hawaiian subjects. In Japanese, the quadratic term (BMI<sup>2</sup>) was statistically significant in a multivariate model ( $P = 0.002$ ). In Latinos, the adjusted least-squares mean IGF-I levels in ng/mL for BMI < 25, 25 to 29.99, and  $\geq 30$  were 184.6, 147.7, and 132.7, respectively. No interaction between race/ethnicity and BMI explained the plasma IGFBP-3 levels in these data. These results may help to resolve the uncertainty in the relationship between circulating IGF levels and obesity and highlight the potential importance of racial/ethnic-specific effects among these factors in explaining ethnic disparities in obesity-related cancers. (Cancer Epidemiol Biomarkers Prev 2006;15(11):2298-302)

## Introduction

Investigation of the relationships between body fat and insulin-like growth factors (IGF) has clinical significance because this hormonal axis is a plausible mechanism linking obesity to cancer risk (1). Previous studies, however, have been inconsistent with respect to the relationships between adiposity and circulating levels of IGF proteins (2-11). Studies among healthy adults have reported a null association (2-4), a positive association (5), an inverse association (6-9), and a nonlinear association (10, 11) between body mass index (BMI) and IGF-I levels. We are unaware of any study that has looked at possible ethnic differences in the relationship between obesity and circulating IGF-I, and this may be important in addressing the metabolic basis for ethnic differences in cancer risk. Previous work has shown that ethnic differences in circulating IGF levels do exist (12, 13). We have previously reported that IGF-I levels are lower among Latinos, compared with four other major racial/ethnic groups (13). Building upon this previous work in the Multiethnic Cohort, we set out to investigate the relationships between plasma IGF protein

levels and obesity, using BMI as an indicator of adiposity, across five racial/ethnic groups (Whites, African Americans, Latinos, Japanese Americans, and Native Hawaiians).

## Materials and Methods

**Study Subjects.** Participants included in these analyses were selected from a large population-based cohort study, the Hawaii and Los Angeles Multiethnic Cohort study (14). The primary aim of this prospective study is to evaluate the dietary and other environmental contributions to the racial/ethnic variability in cancer risk. The Multiethnic Cohort consists of 215,251 men and women, mainly Japanese Americans, Whites, and Native Hawaiians in Hawaii and African Americans and Latinos in Los Angeles. Subjects were recruited between 1993 and 1996 primarily through driver's license files. All participants were between the ages of 47 and 82 at time of blood draw.

Baseline data were collected on cohort participants via a mailed questionnaire that contained five sections: (a) background, including medical history and family cancer history; (b) diet history; (c) medication use; (d) physical activity; and for women (e), female reproductive history, including use of hormones. Height and weight were self-reported by respondents on the baseline questionnaire, and Quetelet's index (kg/m<sup>2</sup>), or BMI, was calculated as a measure of obesity.

For this analysis, blood was collected on a subcohort of about 5,000 randomly selected participants. The draw was completed in the morning, typically at the person's home, after informed consent was obtained. The participation rate for providing a blood sample was 66%. Details of the study have

Received 4/27/06; revised 7/20/06; accepted 8/17/06.

**Grant support:** National Cancer Institute grants CA54281 and CA63464.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Requests for reprints:** Katherine DeLellis Henderson, Department of Preventive Medicine, Norris Comprehensive Cancer Center, University of Southern California, Topping Tower Room 3429A, 1441 Eastlake Avenue, Los Angeles, CA 90033-0800. Phone: 323-865-0312; Fax: 1-323-865-0127. E-mail: kahender@usc.edu

Copyright © 2006 American Association for Cancer Research.

doi:10.1158/1055-9965.EPI-06-0344

been published previously (14). Written informed consent from every subject was obtained. This investigation was approved by the institutional review boards of the University of Southern California and the University of Hawaii.

We measured plasma IGF-I and IGF-binding protein 3 (IGFBP-3) in a random sample of 1,000 of these Multiethnic Cohort participants [100 for each of 10 sex/ethnic groups with equal representation of each 5-year age group at blood draw (>45 years for men and >55 years for women)] who donated a blood sample. Nine hundred sixty subjects had complete IGF-I and IGFBP-3 measurements. One hundred forty-nine subjects were excluded for incomplete questionnaire data for BMI, having prevalent breast, prostate, or colon cancer or use of menopausal hormone therapy at the time of blood draw, leaving 811 subjects for the analysis. The baseline characteristics of these subjects (Table 1) did not differ from those which were described previously (13).

**IGF-I and IGFBP-3 Measurements in Plasma.** Samples were analyzed blind as to the ethnicity and sex of the participants. To reduce the effect of laboratory variability, each analytic batch included equal numbers of subjects from each ethnic/sex group. IGF-I and IGFBP-3 were measured by ELISA from Diagnostic System Laboratories (Webster, TX). IGF-I assays included an acid-ethanol precipitation of IGF-I binding proteins, to avoid interference of IGFBPs with the IGF-I assay. The overall intra-batch coefficients of variation were <10% for both IGF-related proteins. The average inter-batch coefficients of variation were 13.94% and 10.35% for IGF-I and IGFBP-3, respectively.

**Statistical Analysis.** Building on previous gender-stratified work on these proteins in the Multiethnic Cohort (13), we used stepwise and best subsets ( $R^2$  and  $C_p$  criteria) selection to find the set of variables that best explained the variation in IGF-I, IGFBP-3, and IGF-I/IGFBP-3 molar ratio (molar ratio) levels. Variables tested in the previous sex-specific analyses were allowed to compete in these regressions (13), in addition to several new continuous variables. New variables included gender, dietary fiber intake density, percentage of daily calories from carbohydrates, added sugar intake (teaspoons added sugar per day), total fruit and vegetable intake density, regular soda intake density, diet soda intake density, metabolic equivalents of physical activity per day, and energy density of the diet (total calories/total grams food and caloric beverages/day). In the regression analyses, a continuous BMI variable was used. The interaction terms for race/ethnicity  $\times$  BMI, also as a continuous variable, were allowed to compete in the multivariate regression for each protein. To perform these stepwise regressions, the distributions of the continuous protein values were assessed, and subsequently, the Box-Cox procedure was used to find the optimal power transformations, which produced the best approximate normality. Thus, IGF-I, IGFBP-3, and the molar ratios were transformed by raising values to the powers 0.4, 0.5, and 0.15, respectively.

After assessment of the regression models, adjusted mean plasma IGF-I, IGFBP-3, and molar ratios were calculated. These values were adjusted for the covariates found to be important in the regression models and calculated for each category of BMI, defined according to the WHO guidelines as follows: normal, < 25; overweight, 25.01 to 29.99; and obese,  $\geq 30$ , among each racial/ethnic group. Means presented are least-squares means, which were calculated using the GLM procedure. Adjusted mean values were transformed back to their original scale for the purpose of presentation. Predicted values of transformed plasma IGF-I at the average age and gender effect were plotted against BMI by race/ethnicity to visually examine the adjusted relationships between these factors. A similar plot was created for molar ratio, adjusted only for gender. All analyses were done in SAS v9 (SAS Institute, Cary, NC).

## Results

Table 1 shows characteristics for the 811 healthy subjects selected for this analysis by racial/ethnic group. In our sample, African Americans had the lowest proportion of males. Native Hawaiians had the greatest proportion of males. Japanese Americans had the highest proportion of subjects falling into the greatest age category. African Americans had the highest proportion of subjects in the youngest age category. As previously published (13), body size characteristics (weight and height) also differed significantly across racial/ethnic groups in our sample. African-American, Native Hawaiian, and Latino subjects had a higher proportion of subjects with BMI > 25 than Whites or Japanese Americans.

In a multivariate regression, with both genders combined, of the transformed IGF-I variable [root(IGF-I)], the most parsimonious model for IGF-I levels included gender, age, race/ethnicity, BMI, and the interaction between race/ethnicity and BMI. The interaction term for race/ethnicity  $\times$  BMI was highly statistically significant ( $P = 0.005$ ). IGF-I levels differed by BMI category in a race-specific manner. Table 2 provides the least-squares mean plasma IGF-I levels (ng/mL) by racial/ethnic group and BMI category. The least-squares mean IGF-I level declined as BMI increased among Latinos ( $P_{\text{trend}} = 0.0002$ ), Whites ( $P_{\text{trend}} = 0.009$ ), and Japanese ( $P_{\text{trend}} = 0.04$ ). Least-squares mean plasma IGF-I remained stable across BMI categories in African Americans ( $P_{\text{trend}} = 0.97$ ) and Native Hawaiians ( $P_{\text{trend}} = 0.35$ ). We did not find a significant three-way interaction for sex, race/ethnicity, and BMI, and we found that the relationship between IGF-I and BMI was similar for males and females of the same racial/ethnic group (data not shown).

In a multivariate regression, with both genders combined, of the transformed IGFBP-3 variable [root(IGFBP-3)], the most parsimonious model for IGFBP-3 levels included age, race/ethnicity, fat intake from meat, and BMI. In further testing, addition of the interaction between race/ethnicity and BMI in the adjusted model was not significant ( $P = 0.22$ ). Table 2

**Table 1. Characteristics of 811 Multiethnic Cohort participants**

	Latino	White	Japanese American	African American	Native Hawaiian
Total N (%)	171 (21.1)	165 (20.4)	169 (20.8)	147 (18.1)	159 (19.6)
Gender					
Male (%)	52.1	54.6	52.1	47.6	59.8
Age (%)					
<55	28.6	27.9	23.7	32.6	28.3
55-59	18.7	18.2	17.8	21.1	17.6
60-64	20.5	18.8	17.8	10.9	23.9
$\geq 65$	32.2	35.1	40.8	35.4	30.2
BMI (%)					
<25	26.9	41.8	63.3	28.6	28.3
25-29	49.1	38.8	31.4	41.5	37.7
$\geq 30$	24.0	19.4	5.3	29.9	34.0

**Table 2. Least-squares mean (95% confidence limits) plasma IGF-I, IGFBP-3, and IGF-I/IGFBP-3 molar ratio levels (ng/mL) by BMI category (by race/ethnicity)**

	BMI (kg/m <sup>2</sup> )			<i>P</i> <sub>trend</sub>
	<25	25-29.99	≥30	
<b>IGF-I*</b>				
Latino	184.6 (164.7-205.7)	147.7 (134.8-161.4)	132.7 (115.2-151.7)	0.0002
White	184.9 (168.4-202.3)	155.2 (139.9-171.4)	151.3 (130.1-174.5)	0.009
Japanese American	173.7 (161.8-186)	166.5 (150.2-183.9)	122.9 (91.3-160.2)	0.04
African American	172.2 (149.5-196.9)	172.3 (153.3-192.5)	172.9 (150.6-197.1)	0.97
Native Hawaiian	166.0 (147.8-185.4)	184.6 (167.9-202.2)	179.4 (162-197.8)	0.35
<b>IGFBP-3†</b>				
Latino	2,498 (2,252-2,756)	2,488 (2,306-2,676)	2,279 (2,031-2,541)	0.25
White	2,847 (2,635-3,067)	2,881 (2,664-3,106)	2,938 (2,620-3,275)	0.66
Japanese American	2,860 (2,687-3,039)	2,783 (2,542-3,034)	3,112 (2,487-3,806)	0.92
African American	2,609 (2,369-2,861)	2,549 (2,355-2,752)	2,718 (2,476-2,971)	0.55
Native Hawaiian	2,744 (2,488-3,013)	2,971 (2,746-3,205)	2,807 (2,570-3,054)	0.79
<b>IGF molar ratio‡</b>				
Latino	0.28 (0.25-0.30)	0.22 (0.21-0.24)	0.21 (0.19-0.24)	0.002
White	0.24 (0.22-0.26)	0.20 (0.18-0.22)	0.19 (0.17-0.22)	0.002
Japanese American	0.22 (0.21-0.24)	0.22 (0.20-0.24)	0.17 (0.13-0.21)	0.09
African American	0.24 (0.21-0.27)	0.25 (0.23-0.27)	0.24 (0.22-0.27)	0.80
Native Hawaiian	0.22 (0.20-0.25)	0.23 (0.21-0.25)	0.24 (0.22-0.27)	0.22

\*IGF-I is adjusted for covariates found to be significant in a multivariate analysis (gender, age, race/ethnicity, BMI, and the interaction between race/ethnicity and BMI).

†IGFBP-3 is adjusted for covariates found to be significant in a multivariate analysis (age, race/ethnicity, fat intake from meat, and BMI).


‡IGF molar ratio is adjusted for covariates found to be significant in a multivariate analysis (race/ethnicity, gender, BMI, and the interaction between race/ethnicity and BMI).

provides the least-squares mean plasma IGFBP-3 levels (ng/mL) by BMI category and racial/ethnic group. In no individual racial/ethnic group was the difference across BMI categories statistically significant.


Race/ethnicity, gender, BMI, and the interaction between race/ethnicity and BMI were statistically significantly associated with the molar ratio of IGF-I/IGFBP-3 in a multivariate model. The interaction between race/ethnicity and BMI

category was statistically significant for the molar ratio of IGF-I to IGFBP-3 ( $P = 0.003$ ). Table 2 illustrates the pattern of least-squares mean IGF-I/IGFBP-3 molar ratio levels by BMI category and racial/ethnic group. These differences were driven by differences in IGF-I and were statistically significant only in Latinos ( $P_{\text{trend}} = 0.002$ ) and Whites ( $P_{\text{trend}} = 0.002$ ).

Figure 1 is a plot of the predicted IGF-I values at the average value of age and gender plotted against BMI by race/ethnicity.



**Figure 1.** Predicted values for root(IGF-I) for BMI, averaged over gender and age, by race/ethnicity.



**Figure 2.** Predicted values for root(MR) for BMI, averaged over gender, by race/ethnicity.

Visual examination of this plot illustrates the difference across racial/ethnic groups in the adjusted relationship between IGF-I and BMI. IGF-I levels declined in a linear manner with increasing BMI most dramatically in Latinos. This decline was less clear in the White subjects and absent or slightly reversed in the African-American and Native Hawaiian subjects. Among Japanese subjects, IGF-I seemed to have an inverse quadratic relationship with BMI. The quadratic term was statistically significant when tested in the multivariate model ( $P = 0.002$ ).

Figure 2 is a plot of the predicted IGF molar ratios at the average value of gender plotted against BMI by race/ethnicity. Visual examination of this plot illustrates the difference across racial/ethnic groups in the adjusted relationship between the molar ratio and BMI. IGF molar ratio levels declined in a linear manner with increasing BMI most dramatically in Latinos, Whites, and Japanese. In contrast, the relationship tended to be in the opposite direction in African-American and Native Hawaiian subjects, with the molar ratio increasing with increasing BMI.

## Discussion

This study examined the relationship between circulating levels of two primary proteins in the IGF pathway and obesity, among five racial/ethnic groups, using BMI as an indicator of adiposity. IGF-I levels varied not only by racial/ethnic group but also by BMI category in a race-specific fashion. IGF-I levels declined with increasing obesity status most dramatically in Latinos and Japanese. This decline was attenuated in Whites and absent in African-American and Native Hawaiian subjects. In Japanese, the quadratic term ( $BMI^2$ ) was statistically significant in a multivariate model ( $P = 0.002$ ). The interaction between race/ethnicity and BMI was not significant for plasma IGFBP-3. With respect to the interaction between

race/ethnicity and BMI, differences in the molar ratio reflect those in IGF-I.

Previous studies have reported inconsistent findings with respect to the association between BMI and circulating IGF-I levels. Studies among healthy adults have reported a null association (2-4), a positive association (5), an inverse association (6-9), and a nonlinear association (10, 11) between BMI and IGF-I levels. The nonlinear associations were similar to what we observed in Japanese Americans. A recent study among European women reported a relationship between BMI and serum levels of IGF-I similar to that which we saw in Japanese (15). Studies in children, which are important because they allow examination of potentially underlying biological differences across subgroups in the absence of the potential confounding effects of factors, such as smoking, alcohol, aging, menopausal status, and hormone therapy, have previously reported positive correlations between adiposity and IGF-I concentrations (16, 17); however, the relationships could be physiologically different in children in light of data indicating that IGF-I does begin to decrease after puberty (18).

A possible biological mechanism mediating the association between obesity and IGF-I may be through the effect of growth hormone. In individuals with high BMI, IGFBP1 and IGFBP2 are depressed, leading to an increased negative feedback by free IGF-I on pituitary GH secretion and a decreased IGF-I synthesis (ref. 19; for review, see ref. 20). The current data suggest that the GH-IGF axis may be regulated differently in different racial/ethnic groups. Racial/ethnic differences in the distribution of polymorphisms in genes involved in IGF regulation may be one mechanism mediating such an effect.

One possible explanation for the IGF-I result may be if the age-related decline in IGF-I differed between ethnic groups, and there were also differences in the age distribution of obesity within some groups. In these data, obesity uniformly increased with age, and the age-related decline in IGF-I



seemed to be significant only among African Americans. Therefore, these factors could not account for the findings. However, an alternate explanation that cannot be ruled out is that these differences are due to unmeasured confounding factors.

To ensure that the regressions would not be affected by the choice of cut points, we used a continuous BMI variable in the regressions and when assessing the possible significance of the interaction between race and BMI. However, in the models used to calculate the least-squares means, we used a categorical BMI variable, defined according to WHO guidelines, to facilitate comparison with other populations. These categories may not be appropriate to the same extent for all ethnic/racial groups to characterize obesity, as the correlation of adiposity with BMI may vary, and the complications of obesity may seem at different levels of BMI, across ethnic/racial groups.

One possible limitation of the current study is the reliance on a single time point measurement of plasma IGF-I and IGFBP-3. Although data from the Rancho Bernardo Study reported that the intra-individual variation in plasma IGF-I is minimal (21), suggesting that a single measurement may be adequate to predict long-term circulating levels, it should be noted that these samples were collected only 8 to 54 days apart. Other studies have shown measurable inter-individual variation in IGF-I and IGFBP-3 (3, 22-24).

The present results illustrate the importance of race/ethnicity in the relationship between circulating IGF hormones and BMI, particularly for IGF-I. In addition, these findings, if reproduced, have potentially important implications for understanding the link between obesity and cancer risk. In order for a definitive link between circulating IGF-I levels and cancer risk (for review, see ref. 25) to be established, it may be that this apparent interaction between race/ethnicity and BMI in determination of circulating IGF-I levels must be taken into account. This interaction may aid in explaining certain apparent anomalies in the relationship between obesity and cancer. Despite a similar pattern of obesity, degree of insulin resistance, and high risk for type 2 diabetes across some ethnic groups, such as African Americans and Latinos, there are marked differences in cancer incidence across these racial/ethnic groups. In particular, the Latino population in the Multiethnic Cohort has been shown to have a low risk for postmenopausal breast cancer relative to African Americans despite similarly high BMI values. These results may help to resolve the uncertainty in the relationship between the IGF proteins and obesity and highlight the potential importance of racial/ethnic-specific effects among these factors in explaining ethnic disparities in obesity-related cancers.

## Acknowledgments

We thank the participants in the Multiethnic Cohort, whose generosity makes this research possible; Jennifer Yamamoto for analytic support; and Drs. Rudolf Kaaks and Sabina Rinaldi for IGF hormones analyses.

## References

- Giovannucci E. Nutrition, insulin, insulin-like growth factors and cancer. *Horm Metab Res* 2003;35:694-704.
- Lukanova A, Toniolo P, Akhmedkhanov A, et al. A cross-sectional study of IGF-I determinants in women. *Eur J Cancer Prev* 2001;10:443-52.
- Jernstrom H, Deal C, Wilkin F, et al. Genetic and nongenetic factors associated with variation of plasma levels of insulin-like growth factor-I and insulin-like growth factor-binding protein-3 in healthy premenopausal women. *Cancer Epidemiol Biomarkers Prev* 2001;10:377-84.
- Schoen RE, Schragin J, Weissfeld JL, et al. Lack of association between adipose tissue distribution and IGF-1 and IGFBP-3 in men and women [see comment]. *Cancer Epidemiol Biomarkers Prev* 2002;11:581-6.
- Teramukai S, Rohan T, Eguchi H, Oda T, Shinchi K, Kono S. Anthropometric and behavioral correlates of insulin-like growth factor I and insulin-like growth factor binding protein 3 in middle-aged Japanese men. *Am J Epidemiol* 2002;156:344-8.
- Copeland KC, Colletti RB, Devlin JT, McAuliffe TL. The relationship between insulin-like growth factor-I, adiposity, and aging. *Metabolism* 1990;39:584-7.
- Landin-Wilhelmsen K, Wilhelmsen L, Lappas G, et al. Serum insulin-like growth factor I in a random population sample of men and women: relation to age, sex, smoking habits, coffee consumption and physical activity, blood pressure and concentrations of plasma lipids, fibrinogen, parathyroid hormone and osteocalcin. *Clin Endocrinol* 1994;41:351-7.
- Chang S, Wu X, Yu H, Spitz MR. Plasma concentrations of insulin-like growth factors among healthy adult men and postmenopausal women: associations with body composition, lifestyle, and reproductive factors. *Cancer Epidemiol Biomarkers Prev* 2002;11:758-66.
- Morimoto LM, Newcomb PA, White E, Bigler J, Potter JD. Variation in plasma insulin-like growth factor-1 and insulin-like growth factor binding protein-3: personal and lifestyle factors (United States). *Cancer Causes Control* 2005;16:917-27.
- Lukanova A, Soderberg S, Stattin P, et al. Nonlinear relationship of insulin-like growth factor (IGF-I) and IGF-I/IGF-binding protein-3 ratio with indices of adiposity and plasma insulin concentrations (Sweden). *Cancer Causes Control* 2002;13:509-16.
- Lukanova A, Lundin A, Zeleniuch-Jacquotte A, et al. Body mass index, circulating levels of sex-steroid hormones, IGF-I and IGF-binding protein-3: a cross-sectional study in healthy women. *Eur J Endocrinol* 2004;150:161-71.
- Yanovski JA, Sovik KN, Nguyen TT, Sebring NG. Insulin-like growth factors and bone mineral density in African American and White girls. *J Pediatr* 2000;137:826-32.
- DeLellis K, Rinaldi S, Kaaks RJ, Kolonel LN, Henderson B, Le Marchand L. Dietary and lifestyle correlates of plasma insulin-like growth factor-I (IGF-I) and IGF binding protein-3 (IGFBP-3): the multiethnic cohort [erratum appears in *Cancer Epidemiol Biomarkers Prev*. 2004 Nov;13(11 Pt 1):1825]. *Cancer Epidemiol Biomarkers Prev* 2004;13:1444-51.
- Kolonel LN, Henderson BE, Hankin JH, et al. A multiethnic cohort in Hawaii and Los Angeles: baseline characteristics. *Am J Epidemiol* 2000;151:346-57.
- Gram I, Norat T, Rinaldi S, et al. Body mass index, waist circumference and waist-hip ratio and serum levels of IGF-I and IGFBP-3 in European women. *Int J Obes (London)* 2006. Epub ahead of print.
- Garnett SP, Hogler W, Blades B, et al. Relation between hormones and body composition, including bone, in prepubertal children. *Am J Clin Nutr* 2004;80:966-72.
- Ong K, Kratzsch J, Kiess W, Dunger D, Team AS. Circulating IGF-I levels in childhood are related to both current body composition and early postnatal growth rate. *J Clin Endocrinol Metab* 2002;87:1041-4.
- Clemmons DR, Van Wyk JJ. Factors controlling blood concentration of somatomedin C. *J Clin Endocrinol Metab* 1984;13:113-43.
- Attia N, Tamborlane WV, Heptulla R, et al. The metabolic syndrome and insulin-like growth factor I regulation in adolescent obesity. *J Clin Endocrinol Metab* 1998;83:1467-71.
- Calle EE, Kaaks R. Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer* 2004;4:579-91.
- Goodman-Gruen D, Barrett-Connor E. Epidemiology of insulin-like growth factor-I in elderly men and women. *Am J of Clin Epidemiol* 1997;145:970-6.
- Goodman-Gruen D, Barrett-Connor E. Epidemiology of insulin-like growth factor-I in elderly men and women. The Rancho Bernardo Study [see comment][erratum appears in *Am J Epidemiol* 1997 Aug 15;146(4):357]. *Am J Epidemiol* 1997;145:970-6.
- Juul A, Bang P, Hertel NT, et al. Serum insulin-like growth factor-I in 1030 healthy children, adolescents, and adults: relation to age, sex, stage of puberty, testicular size, and body mass index. *J Clin Endocrinol Metab* 1994;78:744-52.
- Juul A, Main K, Blum WF, Lindholm J, Ranke MB, Skakkebaek NE. The ratio between serum levels of insulin-like growth factor (IGF)-I, the IGF binding proteins (IGFBP-1, 2 and 3) decreases with age in healthy adults and is increased in acromegalic patients. *Clin Endocrinol (Oxf)* 1994;41:85-93.
- LeRoith D, Roberts CT, Jr. The insulin-like growth factor system and cancer. *Cancer Lett* 2003;195:127-37.

## Ethnic Disparity in the Relationship between Obesity and Plasma Insulin-Like Growth Factors: The Multiethnic Cohort

Katherine DeLellis Henderson, Michael I. Goran, Laurence N. Kolonel, et al.

*Cancer Epidemiol Biomarkers Prev* 2006;15:2298-2302.

**Updated version** Access the most recent version of this article at:  
<http://cebp.aacrjournals.org/content/15/11/2298>

**Cited articles** This article cites 23 articles, 6 of which you can access for free at:  
<http://cebp.aacrjournals.org/content/15/11/2298.full#ref-list-1>

**Citing articles** This article has been cited by 11 HighWire-hosted articles. Access the articles at:  
<http://cebp.aacrjournals.org/content/15/11/2298.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://cebp.aacrjournals.org/content/15/11/2298>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.