

The Association of a Tobacco-Specific Biomarker and Cigarette Consumption and Its Dependence on Host Characteristics

Jay H. Lubin,¹ Neil Caporaso,² Dorothy K. Hatsukami,³
Anne M. Joseph,³ and Stephen S. Hecht³

¹Biostatistics Branch and ²Genetic Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, Maryland and ³University of Minnesota Transdisciplinary Tobacco Use Research Center and Cancer Center, Minneapolis, Minnesota

Abstract

The tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) is a potent carcinogen, which can be characterized by urinary concentrations of the metabolite 4-(methylnitrosamino)-1-(3-pyridyl)-butanol (NNAL) and its glucuronide. Using baseline data in current smokers from four clinical trials, we examine the associations of urinary cotinine with CPD and of total NNAL with cotinine and the modification of these associations by several host factors. There was a linear relationship between $\ln(\text{cotinine})$ and $\ln(\text{CPD})$ within categories of the Fagerstrom Test of Nicotine Dependence and of age. The increasing trend was significantly smaller for subjects with high and very

high nicotine addiction and for older subjects and larger in females than males. The regression of $\ln(\text{total NNAL}/\text{cotinine})$ on $\ln(\text{cotinine})$ declined linearly, suggesting reduced NNK uptake per unit cotinine with increasing cotinine. The decline in trend was greater in subjects with increased CPD, with greater nicotine addiction, and at older ages and was smaller in females, although gender differences were small. Variations in the ratio with host characteristics were generally similar to a recent epidemiologic analysis of effect modification of the association between lung cancer and cigarette smoking. (Cancer Epidemiol Biomarkers Prev 2007;16(9):1852–7)

Introduction

The tobacco-specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) is a potent carcinogen in rodents and, together with the related compound *N'*-nitrosornicotine, is considered to be a potent human carcinogen (1-3). Uptake of NNK can be characterized by measuring urinary concentrations of the metabolite 4-(methylnitrosamino)-1-(3-pyridyl)-butanol (NNAL) and its glucuronide (termed "total NNAL").

An analysis in current smokers found an increasing, but curvilinear, relationship for total NNAL and cigarettes per day (CPD), with the rate of increase in total NNAL declining with increasing CPD (4). This relationship is suggestive of the attenuation of odds ratios (OR) with increasing CPD observed in epidemiologic studies of lung and bladder cancers (5, 6) and later quantified for several smoking-related cancers (7-9). If attenuation of ORs by CPD derives from biological effects, then we would expect comparable patterns for biomarkers of exposure and effect. It is well known that age, frequency,

and depth of inhalation and sex can modify ORs by cigarette exposure (10). Thus, we would anticipate that patterns for biomarkers may also vary by host factors, which may provide insights into mechanisms of action.

In this article, we use data from Joseph et al. (4) to evaluate the relationships between urinary cotinine and CPD and between total NNAL and cotinine and modification of the relationships by various host factors. This represents the first detailed examination of the regression of total NNAL on cotinine and factors influencing the relationship. Joseph et al. also analyzed 1-hydroxypyrene, which is a metabolite of pyrene, a noncarcinogenic polycyclic aromatic hydrocarbon. However, 1-hydroxypyrene is not tobacco specific, and we do not consider it further.

Materials and Methods

Data. We combine baseline data for current smokers from four clinical trials (4). (a) Tobacco Reduction Intervention Program: the study was a clinical trial of cigarette smokers ages 18 to 70 years, who wanted to reduce, but not quit, smoking. All subjects smoked 15 to 45 CPD for at least 1 year prior, were apparently in good health, had no contraindications to nicotine replacement treatments, had no history of schizophrenia or unstable depressive disorder, did not use other tobacco products, and were not pregnant or nursing. CPD was calculated from a daily diary. Biomarker data were determined at two points, 1 week apart, and averaged. The study included 69 males and 82 females. (b) Reduction of

Received 1/10/07; revised 5/2/07; accepted 6/20/07.

Grant support: Intramural Research Program of the National Cancer Institute, NIH, and NIH grant P50 DA013333.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Jay H. Lubin, Biostatistics Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, 6120 Executive Boulevard, Rockville, MD 20852. Phone: 301-496-3357; Fax: 301-402-0081. E-mail: lubinj@mail.nih.gov

Copyright © 2007 American Association for Cancer Research.

doi:10.1158/1055-9965.EPI-07-0018

Smoking in Cardiac Patients Study: the study was a clinical trial of cigarette smokers ages 18 to 80 years with at least one diagnosis of heart disease who wanted to reduce, but not quit, smoking. All subjects smoked ≥ 15 CPD, did not use other tobacco products, were diagnosed with at least one cardiac disease (coronary artery disease, arrhythmia, congestive heart failure, or peripheral vascular disease) or had a history of cerebrovascular disease, had no unstable angina in the prior 2 weeks, had no unstable mental health or substance use diagnoses, and had no contraindications to nicotine replacement therapy. CPD was based on recall for the prior week. Biomarkers were measured once at baseline. The study included 135 males and 17 females. (c) Adult Cross-Sectional Study: the study recruited subjects from the Minneapolis-Saint Paul metropolitan area under the Tobacco Reduction Intervention Program eligibility criteria. The study included 86 subjects (34 males, 46 females, and 6 with unknown gender). (d) Low Level Smoking Study: the study recruited subjects from the Minneapolis Veterans Affairs Medical Center under the Reduction of Smoking in Cardiac Patients eligibility criteria, with the additional restriction that smokers consumed <15 CPD for 1 or more years. The study included 11 subjects (8 males and 3 with unknown gender).

The pooled data set includes 400 subjects. Nine subjects have unknown gender, 48 are missing total NNAL or cotinine measurements, and 3 are missing CPD, leaving 340 subjects. We omit two smokers with extreme cotinine values (0.17 nmol/mg creatinine, one third the next higher value; and 159.4 nmol/mg creatinine, over two times the next lower value) and one smoker who consumed 100 CPD (40 CPD more than the next lower value). The final data set includes 337 subjects and currently represents the largest data set of smokers with total NNAL measurements. Means (SDs) for CPD for males and females are the following: Tobacco Reduction Intervention Program, 24.3 (6.2) and 22.8 (5.2); Reduction of Smoking in Cardiac Patients, 27.0 (10.0) and 28.6 (11.0); Adult Cross-sectional Study, 11.3 (10.0) and 8.2 (6.2); and Low Level Smoking Study, 7.1 (3.6).

Statistical Analysis. We apply standard linear regression methods with either cotinine or total NNAL as dependent variable. Within categories of sex and CPD, cotinine and total NNAL are consistent with log-normal distributions, and we therefore apply logarithmic transformations. For response Y and continuous variable X , we fit the linear regression

$$\ln(Y) = \beta_0 + \beta_1 \ln(X) + \beta_2 Z \quad (A)$$

where β_1 is the slope parameter (i.e., the exponent in a power relationship between Y and X) and Z is a vector of adjustment variables. We evaluate departures from linearity by including $[\ln(X)]^2$ and testing the null hypothesis that its coefficient equals zero.

Depending on the analysis, adjustment factors (Z) include age (four levels: <40 , 40-49, 50-59, ≥ 60 years), the Fagerstrom Test of Nicotine Dependence (FTND) score (11, 12) grouped into five levels of addiction (0-2, very low; 3-4, low; 5, medium; 6-7, high; 8-10, very high), and sex. We calculate a likelihood ratio test of homogeneity of

the effect of $\ln(X)$ over an adjustment variable by adding to the model the product of $\ln(X)$ and the indicator variables for the adjustment variable.

We first regress $\ln(\text{cotinine})$ on $\ln(\text{CPD})$ and evaluate the shape of the regression and its modification by age, FTND, and sex. We then regress $\ln(\text{total NNAL}/\text{cotinine})$ on $\ln(\text{cotinine})$. We use the ratio because it approximates a measure of smoking potency (i.e., the uptake of NNK per unit cotinine). In addition, a horizontal line represents a proportional relationship between total NNAL and cotinine so that deviations from proportionality are easily identified. Because the half-life of total NNAL is 3 to 4 days (13) and of cotinine is 15 to 19 h (14, 15), the ratio is temporally appropriate for current smokers.

There is homogeneity across studies for the associations between cotinine and CPD ($P = 0.75$ for males and $P = 0.40$ for females) and between total NNAL and cotinine ($P = 0.58$ for males and $P = 0.33$ for females after adjusting for CPD). Therefore, we do not further consider study effects.

Results

Females smoke fewer CPD than males, with means (SDs) of 18.1 (10.2) and 23.3 (10.9) CPD, respectively. Means for cotinine and total NNAL increase with CPD, with variability generally higher at lower CPD (Table 1). Adjusting for CPD, concentrations do not differ by sex for cotinine ($P = 0.24$) or total NNAL ($P = 0.82$).

Cotinine and CPD Association. Cotinine increases with CPD but at a diminishing rate at higher CPD (Fig. 1, top left). Compared with males, cotinine levels in females are similar at lower CPD but higher at higher CPD. In females, cotinine increases across CPD, whereas in males cotinine increases to 34.0 CPD [maximum for fitted line at $\exp(\beta_1/-2\beta_2)$] and then declines.

Figure 1 also shows the regression of $\ln(\text{cotinine})$ on $\ln(\text{CPD})$ within FTND categories. Cotinine increases

Table 1. Statistics on urinary cotinine (nmol/mg creatinine) and total NNAL (pmol/mg creatinine) for categories of CPD, including number of subjects, arithmetic mean, geometric mean, and geometric SD

	CPD	n*	Cotinine			Total NNAL		
			AM	GM	GSD	AM	GM	GSD
Males	Overall	216	21.4	17.0	2.23	2.13	1.75	1.96
	10	28	9.6	6.8	2.66	0.99	0.74	2.30
	11-20	80	21.5	17.7	2.13	2.07	1.79	1.73
	21-30	71	25.5	22.5	1.70	2.34	2.04	1.71
	31+	37	22.0	18.2	2.02	2.72	2.35	1.73
Females	Overall	121	23.2	15.8	2.85	2.09	1.48	2.62
	10	32	7.8	5.0	2.82	0.77	0.52	2.55
	11-20	40	25.0	19.9	2.18	2.15	1.76	2.02
	21-30	39	30.6	26.7	1.77	2.60	2.36	1.62
	31+	10	36.6	32.7	1.78	4.11	3.50	1.94

NOTE: Data from four clinical trials of smokers (see text).

Abbreviations: AM, arithmetic mean; GM, geometric mean; GSD, geometric SD.

*Omits two males with extreme cotinine measurement of 0.17 and 159.4 nmol/mg creatinine, one subject who smoked 100 CPD, and subjects with missing data.




Figure 1. Scatter plot of urinary cotinine concentrations in nmol/mg creatinine by CPD and fitted model A for females and males, overall and within scores of the FTND, grouped into levels of addiction (0-2, very low; 3-4, low; 5, medium; 6-7, high; 8-10, very high).

with CPD in subjects with very low, low, and medium addiction scores but is unrelated to CPD in subjects with high and very high addiction scores. Homogeneity of slopes by FTND category is rejected ($P < 0.001$). The female differential is significant ($P = 0.01$) but similar across FTND category ($P = 0.35$). The FTND index measures nicotine dependency and is the sum of categorical scores for CPD and five other variables. Thus, the inclusion of CPD in the FTND score may have influenced these results. Adjusting for sex and each component, regression patterns of $\ln(\text{cotinine})$ on $\ln(\text{CPD})$ within each component are consistent with overall FTND results (i.e., increased trends in lower addictive categories and diminished trends in higher addictive categories). Homogeneity of slopes across categories, however, is not always rejected. Tests of homogeneity for FTND components include time to first cigarette on awaking ($P < 0.01$), difficulty in refraining from smoking ($P = 0.14$), difficulty giving up first morning cigarette ($P = 0.02$), increased smoking during the morning ($P = 0.31$), and smoke when ill ($P < 0.01$).

The regression of $\ln(\text{cotinine})$ on $\ln(\text{CPD})$ varies by age group ($P = 0.01$) and sex ($P = 0.03$), with a greater slope at ages under 50 and for females (Fig. 2). In the oldest age group, three subjects who consumed one CPD are highly influential. Omitting these subjects, slopes decline monotonically by age group [$P < 0.001$, estimates of β_1 for males are 0.93, 0.64, 0.44, and 0.28 (dot line), respectively]. The female differential is homogeneous across age groups ($P = 0.90$).

The regressions remain significantly modified by sex ($P = 0.02$), age ($P = 0.05$), and FTND ($P < 0.01$) when these factors are jointly analyzed.

Total NNAL and Cotinine Association. The ratio of total NNAL to cotinine decreases with increasing cotinine, indicating a decrease in NNK uptake per unit cotinine (Fig. 3, top left). Linearity on log-log scales is not rejected ($P = 0.12$), although curvilinearity is suggested in males ($P = 0.05$; dot line).

Trends decline within each CPD category (Fig. 3) and are consistent with linearity ($P = 0.72$). Homogeneity of slopes is rejected ($P = 0.01$), with β_1 estimates for males of -0.39 , -0.63 , -0.75 , and -0.69 , respectively. The female differential effect is significant ($P = 0.01$). The power parameter is increased for females by 0.20 with 95% confidence interval of 0.03 to 0.44.

The regression of $\ln(\text{total NNAL}/\text{cotinine})$ on $\ln(\text{cotinine})$ across FTND scores varies significantly ($P < 0.01$), due primarily to the very low addiction group, and




Figure 2. Scatter plot of urinary cotinine concentrations in nmol/mg creatinine by CPD and fitted model A for females and males within categories of age. Dot line is fitted model A for males with one cigarette per day smokers omitted.




Figure 3. Scatter plot of the ratio of total NNAL (in pmol/mg creatinine) to cotinine (in pmol/mg creatinine) by cotinine and fitted model A for females and males, overall and within categories of smoking intensity. Dot line is fitted model A with linear and quadratic terms for males.

differs by sex ($P = 0.01$; Fig. 4). A post hoc test of homogeneity of slopes for the very low addiction category compared with higher addiction categories is rejected ($P < 0.01$). Results are consistent within each component variable of the FTND. There is a greater decline in the higher “addictive” categories, including CPD ($P = 0.01$), time to first cigarette on awaking ($P = 0.07$), difficulty in refraining from smoking ($P = 0.55$), difficulty giving up first morning cigarette ($P = 0.10$), increased

frequency during the morning ($P = 0.36$), and smoke when ill ($P < 0.01$).

$\ln(\text{total NNAL}/\text{cotinine})$ declines with $\ln(\text{cotinine})$ within age groups, with slopes homogeneous ($P = 0.10$; Fig. 5). Patterns differ by sex ($P < 0.01$). Estimates of β_1 for males, -0.46 , -0.38 , -0.43 , and -0.62 , respectively, suggest an increased rate of decline in the oldest age group. A post hoc test of homogeneity of slopes for subjects over age 60 is rejected ($P = 0.02$).




Figure 4. Scatter plot of the ratio of total NNAL (in pmol/mg creatinine) to cotinine (in pmol/mg creatinine) by cotinine and fitted model A for females and males within score of the FTND, grouped into levels of addiction (0-2, very low; 3-4, low; 5, medium; 6-7, high; 8-10, very high).

Discussion

Previous reports suggest a linear relationship between cotinine and CPD under 20 to 25 CPD (4, 16-27). At higher CPD, the relationship has been observed to increase without diminution (19, 23, 25, 26), increase but at a diminished rate (17, 19-21, 24, 27), or increase to a plateau and possibly even decline (4, 16, 19, 22). In our data, urinary cotinine concentration diminished at higher CPD. Using log-log scales, the relationship was linear but varied within categories of FTND score and age. Trends were greater in females, and were significantly reduced in subjects with high and very high addiction levels, and in older subjects. Confounding by these or other host factors may help explain the diverse relationships observed between CPD and cotinine in the various studies.

Our study is the first detailed evaluation of the ratio of NNAL to cotinine and its relationship with cotinine. The ratio declined with increasing cotinine, with the rate of decline greater in subjects with greater cigarette consumption, with increased nicotine addiction, at older ages, and in males. One possible explanation for the general pattern is that enzymes involved in other pathways of NNK metabolism are induced at higher levels of smoking intensity, thus decreasing the flux of NNK metabolites through the NNAL pathway. We have previously observed decreased levels of total NNAL in the urine of smokers who were exposed to indole-3-carbinol presumably due to induction of hepatic cytochrome *P450 1A2* (28, 29). Explanations for the variations in the regression of $\ln(\text{total NNAL}/\text{cotinine})$ on $\ln(\text{cotinine})$ by the various host factors need further development.

We standardized total NNAL by cotinine, rather than CPD, because cotinine better reflects "internal" tobacco exposure (23). This minimizes bias due to misclassification of smoking rate and minimizes the possible effect of a decline in "nicotine boost" (i.e., an increase in blood plasma nicotine per cigarette) with increasing CPD (30). Our results therefore do not likely reflect misclassification of tobacco exposure by CPD or nicotine satiation with increasing CPD but likely reflect host factor-dependent biological effects linked to NNK metabolism.

The effects of host factors on the regression of $\ln(\text{total NNAL}/\text{cotinine})$ on $\ln(\text{cotinine})$ (Figs. 3-5) are broadly similar to analyses of cigarette smoking risks and effect modifications in two lung cancer case-control studies (31). The analysis was based on a linear relationship for the excess OR with pack-years and included a function for the modification of the excess OR/pack-year by CPD. Results indicated that for a fixed total cigarette consumption there was a decreasing exposure rate effect above 15 CPD (i.e., for equal total exposure risk decreased with CPD). In addition, CPD effects were greater for older subjects, which was similar to our finding that the declining trend of $\ln(\text{total NNAL}/\text{cotinine})$ on $\ln(\text{cotinine})$ was greatest at ages >60 years. In the epidemiologic analysis, CPD had less effect on risk in subjects with putatively "lower risk" smoking behavior (i.e., subjects who inhaled less frequently or into the mouth and throat compared with subjects who inhaled frequently or moderately or deeply into the chest). Although comparable inhalation information was




Figure 5. Scatter plot of the ratio of total NNAL (in pmol/mg creatinine) to cotinine (in pmol/mg creatinine) by cotinine and fitted model A for females and males within categories of age.

not available, we found lower rates of decline in the regression trends with low CPD and with low FTND scores. Finally, in the lung cancer analysis, although not statistically significant, CPD had a smaller effect on risk in females compared with males, which was comparable with our results of smaller declines in regression trends in females.

In summary, we find a linear relationship between $\ln(\text{cotinine})$ and $\ln(\text{intensity})$ within FTND addiction score and within age group. The increasing trend was significantly smaller for subjects with high nicotine dependence, for older subjects, and for males compared with females. The regression of $\ln(\text{total NNAL}/\text{cotinine})$ on $\ln(\text{cotinine})$ showed a declining trend, suggesting reduced uptake of NNK per unit cotinine with increasing cotinine. The declining trend was greater with increased CPD, with high nicotine dependence and at older ages, and in males. Variations in the ratio of total NNAL to cotinine by host characteristics were generally consistent with an epidemiologic analysis of effect modification of lung cancer and cigarette smoking (31).

References

1. Hecht SS. Human urinary carcinogen metabolites: biomarkers for investigating tobacco and cancer. *Carcinog* 2002;23:907-22.
2. Hecht SS. Biochemistry, biology, and carcinogenicity of tobacco-specific *N*-nitrosamines. *Chem Res Toxicol* 1998;11:559-603.
3. International Agency for Research on Cancer. IARC monographs on the evaluation of carcinogenic risks to humans. Lyon (France): IARC; 2006.
4. Joseph AM, Hecht SS, Murphy SE, et al. Relationships between cigarette consumption and biomarkers of tobacco toxin exposure. *Cancer Epidemiol Biomarkers Prev* 2005;14:2963-8.
5. Vineis P, Alavanja M, Garte S. Dose-response relationship in tobacco-related cancers of bladder and lung: a biochemical interpretation. *Int J Cancer* 2004;108:2-7.
6. Vineis P, Kogevinas M, Simonato L, Brennan P, Boffetta P. Levelling-off of the risk of lung and bladder cancer in heavy smokers: an analysis based on multicentric case-control studies and a metabolic interpretation. *Mutat Res Rev in Mutat Res* 2000;463:103-10.

7. Lubin JH, Caporaso N. Cigarette smoking and lung cancer: modeling total exposure and intensity. *Cancer Epidemiol Biomarkers Prev* 2006;15:517–23.
8. Lubin JH, Alavanja MCR, Caporaso N, et al. Cigarette smoking and cancer: modeling total exposure and intensity. *Am J Epidemiol* 2007;166:479–89.
9. Lubin JH, Kogevinas M, Silverman DT, et al. Evidence for an intensity dependent interaction of NAT2 acetylation genotype and cigarette smoking in the Spanish Bladder Cancer Study. *Int J Epidemiol* 2007;36:236–41.
10. U.S. Department of Health and Human Services. The health consequences of smoking: a report of the Surgeon General. Washington (DC): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, Superintendent of Documents, U.S. Government Printing Office; 2004.
11. Fagerstrom KO, Schneider NG. Measuring nicotine dependence—a review of the Fagerstrom tolerance questionnaire. *J Behav Med* 1989;12:159–82.
12. Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerstrom test for nicotine dependence—a revision of the Fagerstrom tolerance questionnaire. *Br J Addict* 1991;86:1119–27.
13. Hecht SS, Carmella SG, Chen ML, et al. Quantitation of urinary metabolites of a tobacco-specific lung carcinogen after smoking cessation. *Cancer Res* 1999;59:590–6.
14. Benowitz NL, Perez-Stable EJ, Fong I, Modin G, Herrera B, Jacob P. Ethnic differences in N-glucuronidation of nicotine and cotinine. *J Pharmacol Exp Ther* 1999;291:1196–203.
15. Perez-Stable EJ, Herrera B, Jacob P, Benowitz NL. Nicotine metabolism and intake in black and white smokers. *JAMA* 1998;280:152–6.
16. Etter JF, Perneger TV. Measurement of self reported active exposure to cigarette smoke. *J Epidemiol Community Health* 2001;55:674–80.
17. Law MR, Morris JK, Watt HC, Wald NJ. The dose-response relationship between cigarette consumption, biochemical markers and risk of lung cancer. *Br J Cancer* 1997;75:1690–3.
18. Abrams DB, Follick MJ, Biener L, Carey KB, Hitti J. Saliva cotinine as a measure of smoking status in field settings. *Am J Public Health* 1987;77:846–8.
19. Blackford AL, Yang G, Hernandez-Avila M, et al. Cotinine concentration in smokers from different countries: relationship with amount smoked and cigarette type. *Cancer Epidemiol Biomarkers Prev* 2006;15:1799–804.
20. Campuzano JC, Hernandez-Avila M, Jaakkola MS, et al. Determinants of salivary cotinine levels among current smokers in Mexico. *Nicotine Tob Res* 2004;6:997–1008.
21. Jaakkola MS, Ma JM, Yang GH, et al. Determinants of salivary cotinine concentrations in Chinese male smokers. *Prev Med* 2003;36:282–90.
22. Heinrich J, Holscher B, Seiwert M, Carty CL, Merkel G, Schulz C. Nicotine and cotinine in adults' urine: The German Environmental Survey 1998. *Expo Anal Environ Epidemiol* 2005;15:74–80.
23. Lewis SJ, Cherry NM, Niven RM, Barber PV, Wilde K, Povey AC. Cotinine levels and self-reported smoking status in patients attending a bronchoscopy clinic. *Biomarkers* 2003;8:218–28.
24. O'Connor RJ, Giovino GA, Kozlowski LT, et al. Changes in nicotine intake and cigarette use over time in two nationally representative cross-sectional samples of smokers. *Am J Epidemiol* 2006;164:750–9.
25. Olivieri M, Poli A, Zuccaro P, Ferrari M, et al. Tobacco smoke exposure and serum cotinine in a random sample of adults living in Verona, Italy. *Arch Environ Health* 2002;57:355–9.
26. Richie JP, Carmella SG, Muscat JE, Scott DG, Akerkar SA, Hecht SS. Differences in the urinary metabolites of the tobacco-specific lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in black and white smokers. *Cancer Epidemiol Biomarkers Prev* 1997;6:783–90.
27. Hill P, Haley NJ, Wynder EL. Cigarette-smoking: carboxyhemoglobin, plasma nicotine, cotinine and thiocyanate vs self-reported smoking data and cardiovascular disease. *J Chronic Dis* 1983;36:439–49.
28. Hecht SS, Carmella SG, Kenney PMJ, Low SH, Arakawa K, Yu MC. Effects of cruciferous vegetable consumption on urinary metabolites of the tobacco-specific lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in Singapore Chinese. *Cancer Epidemiol Biomarkers Prev* 2004;13:997–1004.
29. Taioli E, Garbers S, Bradlow HL, Carmella SG, Akerkar S, Hecht SS. Effects of indole-3-carbinol on the metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in smokers. *Cancer Epidemiol Biomarkers Prev* 1997;6:517–22.
30. Patterson F, Benowitz N, Shields P, et al. Individual differences in nicotine intake per cigarette. *Cancer Epidemiol Biomarkers Prev* 2003;12:468–71.
31. Lubin JH, Caporaso N, Wichmann HE, Schaffrath-Rosario A, Alavanja MCR. Cigarette smoking and lung cancer: modeling effect modification of total exposure and intensity. *Epidemiol* 2007;18:639–48.

The Association of a Tobacco-Specific Biomarker and Cigarette Consumption and Its Dependence on Host Characteristics

Jay H. Lubin, Neil Caporaso, Dorothy K. Hatsukami, et al.

Cancer Epidemiol Biomarkers Prev 2007;16:1852-1857.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/16/9/1852>

Cited articles This article cites 29 articles, 10 of which you can access for free at:
<http://cebp.aacrjournals.org/content/16/9/1852.full#ref-list-1>

Citing articles This article has been cited by 5 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/16/9/1852.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/16/9/1852>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.