

Review

Reconciling Human Smoking Behavior and Machine Smoking Patterns: Implications for Understanding Smoking Behavior and the Impact on Laboratory Studies

Catalin Marian,¹ Richard J. O'Connor,² Mirjana V. Djordjevic,³ Vaughan W. Rees,⁴ Dorothy K. Hatsukami,⁵ and Peter G. Shields¹

¹Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, District of Columbia; ²Department of Health Behavior, Roswell Park Cancer Institute, Buffalo, New York; ³Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, Maryland; ⁴Division of Public Health Practice, Harvard School of Public Health, Boston, Massachusetts; and ⁵University of Minnesota Transdisciplinary Tobacco Use Research Center, Minneapolis, Minnesota

Abstract

Background: Recent Food and Drug Administration legislation enables the mandating of product performance standards for cigarette smoke and the evaluation of manufacturers' health claims for modified tobacco products. Laboratory studies used for these evaluations and also for understanding tobacco smoke toxicology use machines to generate smoke. The goal of this review is to critically evaluate methods to assess human smoking behavior and replicate this in the laboratory. **Methods:** Smoking behavior and smoking machine studies were identified using PubMed and publicly available databases for internal tobacco company documents. **Results:** The smoking machine was developed to generate smoke to allow for comparing cigarette tar and nicotine yields. The intent was to infer relative human disease risk, but this concept was flawed because humans tailor their smoking to the product, and chemical

yields and toxicologic effects change with different smoking profiles. Although smoking machines also allow for mechanistic assessments of smoking-related diseases, the interpretations also are limited. However, available methods to assess how humans puff could be used to provide better laboratory assessments, but these need to be validated. Separately, the contribution of smoke mouth-holding and inhalation to dose need to be assessed, because these parts of smoking are not captured by the smoking machine. Better comparisons of cigarettes might be done by tailoring human puff profiles to the product based on human studies and comparing results across regimens.

Conclusions: There are major research gaps that limit the use of smoking machine studies for informing tobacco control regulation and mechanistic studies. (Cancer Epidemiol Biomarkers Prev 2009;18(12):3305–20)

Introduction

In June 2009, the Food and Drug Administration (FDA) received regulatory authority over tobacco products. The FDA is now empowered to develop product performance standards and evaluate manufacturers' health claims for modified tobacco products. Tobacco manufacturers have again publicly focused efforts on lowering cigarette smoke emissions, and may be able to make health claims following the FDA review of their scientific data. The World Health Organization (WHO) Study Group on Tobacco Product Regulation and others also have recognized potential benefits and pitfalls for tobacco harm reduction strategies (1–10). The Institute of Medicine

furthered this harm reduction concept by concluding that harm reduction through smoke exposure reduction was feasible (11, 12). The Institute of Medicine coined an overarching term, PREPs, for potential reduced exposure products (a comprehensive list of existing PREPs can be found at Tobaccoproducts.org).⁶ As FDA performance standards to reduce exposure are developed and implemented, and the manufacturers develop new product designs proposed to reduce human tobacco toxicant exposure, reliable and validated methods are needed to assess changes in cigarette smoke chemical yields and toxicologic effects. Critical to the laboratory evaluation of these products is the generation of cigarette smoke by smoking machines, for example, as have been historically used to estimate tar and nicotine yields. However, prior uses of the smoking machine have been invalidated in the context of understanding human risks and comparing different types of cigarettes, because smoking machine protocols do not replicate human exposure. Thus, current

Received 9/28/09; revised 10/19/09; accepted 10/19/09; published online 12/3/09.

Grant support: National Cancer Institute N01-PC-64402-Laboratory Assessment of Tobacco Use Behavior and Exposure to Toxins Among Users of New Tobacco Products Promoted to Reduce Harm.

Requests for reprints: Peter G. Shields, Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, 3800 Reservoir Road Northwest, LL (S) Level, Room 150, Washington, DC 20057-1465. Phone: 202-687-0003; Fax: 202-687-0003. E-mail: pgs2@georgetown.edu

Copyright © 2009 American Association for Cancer Research.
doi:10.1158/1055-9965.EPI-09-1014

⁶ <http://www.tobaccoproducts.org>

methods preclude an estimation of human exposure and toxicologic effects, challenging new regulatory processes.

The best example of the flawed use of the smoking machine relates to the earlier assumptions that reduced tar and nicotine yield cigarettes (as ranked based on data obtained by standard FTC/ISO machine-smoking method), the so-called "lights," were less harmful than higher yield cigarettes (13). Almost 3 decades ago, the public health community advocated that for smokers who could not or would not quit to switch to lower tar yield cigarettes, for example as recommended by the Surgeon General (14-16). Advertising and marketing by the tobacco industry reinforced the perceptions that lower tar was less harmful. We now know that smoking machine yields were misunderstood in relation to human exposure and tobacco companies intentionally misrepresented the impact of lowering tar yields on smokers' health (13, 17, 18). Development of smoking behavior measurements revealed that compensatory mechanisms for adjusting to the reduced nicotine yields of reduced yield cigarettes led smokers to increase their nicotine exposure by increasing cigarette puffing intensity and smoking more cigarettes per day (19-23). Moreover, human biomarker studies have shown that smokers' exposures were not different when smoking cigarettes with different tar yields (24-26). Separately, although early epidemiology data supported the hypothesis for reduced risk in relation to tar yields, a recent reanalysis of the data established that the early interpretations were wrong (13, 27). As the realization for the limitations of smoking machine studies became clear, and how the uses for public health recommendations were based on flawed interpretations, the Federal Trade Commission (FTC) in November 2008 officially rescinded its widely used guidance for reporting smoking machine determined tar and nicotine yields.⁷ Thus, today, there are no recommended smoking machine protocols in the United States that the FDA can use to inform their decision making processes regarding performance standards and health claims, although the WHO has made recommendations (see below; refs.10, 28).

To develop and validate new smoking machine methods, a better understanding of how to assess human smoking behavior is needed. Currently, smoking behavior is assessed by smoking topography devices that record puff profiles (e.g., puff volume, interpuff interval, puff duration, and air flow) and methods to assess inhalation. However, there are limitations to these methods for estimating human exposure, and very few studies combine these research tools to cover all the components of smoking. Conceptually, these methods could be validated by human biomarker studies, and some studies have been done.

The goal of this review is to critically evaluate methods to assess human smoking behavior and how best to replicate this on smoking machines. Although there will always be limitations to such studies, certain limitations can be mitigated, and the context for other limitations can be better understood. This review will summarize the state of the art in smoking machine protocols and human smoking behavior measurement, identifying what is and

is not captured by smoking machine replication of human smoking. These data will be synthesized to identify research gaps related to laboratory research on cigarette smoke and regulation of tobacco products. This review is organized into three major sections, followed by a discussion. The first section provides a review of the technical aspects of machine smoking and the early development of the standardized smoking machine. This will set the stage to contrast this early work with what we know about human smoking behavior and how well we measure that. The third section reviews methods where researchers have tried to apply what we know about human smoking behavior for smoking machine studies. Last, the discussion provides an overall summary of the most important points and identifies the research gaps that lead from earlier work.

Materials and Methods

Smoking behavior and smoking machine studies were identified using PubMed search strategies. The search keyword strings included "human smoking behavior, smoking topography, human puff profiles, smoking machines, smoke exposure, and PREPs," and combinations of these. All identified studies were reviewed that have been published since 1980, and citation lists were cross-referenced to ensure that the most complete list of publications was identified. Articles published before 1980 with high relevance to the study of PREPs or low-yield cigarettes also were identified and reviewed. Separately, internal tobacco company documents were reviewed, as identified by searches using TobaccoDocuments.org⁸ and the Legacy Tobacco Documents Library.⁹

Studies were identified that investigated methodologic, descriptive, validation, and application aspects related to the assessment of human smoking behavior, human puff and respiration patterns, biomarkers of acute smoke exposure, and smoking machine regimens and yields as they relate to exposure. Research publications were compiled to examine the following: (a) the goals of the study; (b) the methods for assessing human smoking behavior or machine smoking protocols; (c) the experimental designs that were used; and (d) the effects of smoking behavior in relation to the effects of smoking machine protocols on smoke yields. The information was synthesized to provide usefulness for the study of cigarettes and identify research gaps. Although others have reviewed the origins and limitations of smoking machine yield testing (29-31), the focus of this article is to identify how to better replicate human smoking in the laboratory through understanding both the design of smoking machine and human behavior studies, and identify the research gaps associated with this.

Results

Technical Aspects of Machine Smoking. Smoking machines are intended to generate smoke in a systematic

⁷ <http://www.ftc.gov/opa/2008/11/cigarettestesting.shtm>

⁸ <http://www.tobaccodocuments.org>

⁹ <http://legacy.library.ucsf.edu/>

Table 1. Overview of smoking regimens (from ref. 149)

Regimen	FTC	Massachusetts	Canadian	ISO	ISO A	ISO B	ISO C
Puff volume	35 ± 0.5 mL	45 ± 0.5 mL	55 ± 0.5 mL	35 ± 0.3 mL	55	60	45
Puff duration	2 ± 0.05 s	2 ± 0.05 s	2 ± 0.05 s	2 ± 0.05 s	2	2	2
Puff frequency	60 ± 0.5 s	30 ± 0.5 s	30 ± 0.5 s	60 ± 0.5 s	30	30	30
Ventilation holes	Open	50% blocked	100% blocked	Open	50%	50%	100%
Conditioning atmosphere	60% RH ± 2% RH 23.9°C ± 1.1°C min 1, max 14 d	60% RH ± 2% RH 23.9°C ± 1.1°C min 1, max 14 d	60% RH ± 3% RH 22°C ± 1°C min 2, max 10 d	60% RH ± 3% RH 22°C ± 1°C min 2, max 10 d	blocked	blocked	blocked
Smoking environment	60% RH ± 3% RH 23.9°C 4 ± 2°C	60% RH ± 3% RH 23.9°C ± 2°C	60% RH ± 5% RH 22°C ± 2°C	60% RH ± 5% RH 22°C ± 2°C			
Air flow							
linear ind.			200 ± 50 mL/min	200 ± 50 mL/min			
port linear			200 ± 30 mL/min	200 ± 30 mL/min			
avg. rotary							
Butt length (whichever is the highest value)	Tipping + 3 or 23 mm from butt	Tipping + 3 or 23 mm from butt	Tipping + 3 mm or filter + 8 or 23 mm from butt	Tipping + 3 mm or filter + 8 or 23 mm from butt			

NOTE: FTC and Massachusetts protocols have air flow sufficient to exhaust the smoke, about 20 mL/min. Abbreviation: RH, relative humidity.

fashion for laboratory testing, and they have been used to compare cigarette smoke toxicant yields by puffing cigarettes according to specified settings. Cigarette smoke is a suspension of particles in a gaseous vapor, and so it can be collected and analyzed in various ways. A recent review comprehensively describes how smoke is collected for toxicology studies (32). Particles in smoke can be collected on a Cambridge filter pad, which is composed of glass fibers. The change in weight of the pad defines the total particulate matter or wet total particulate matter. Tar is a mathematically derived value defined as total particulate matter minus water and nicotine. The gas and vapor phase passes through the Cambridge filter pad and can be collected or tested directly. Alternatively, smoke can be collected as a condensate usually in a liquid trap (termed cigarette smoke condensate) or directly assayed as whole smoke. Total particulate matter and cigarette smoke condensate are typically used in studies assessing the toxicology of tobacco smoke *in vitro* and for animal skin painting studies, and for assessing the chemical constituents. Whole smoke is used to determine the smoke constituents and in inhalational animal studies, although it is sometimes used for *in vitro* toxicology studies. Although smoking machines have several variables that can be adjusted, typically, the programmable parameters are puff volume, puff frequency, puff duration, the length of cigarette smoked (butt length), and more recently, puff shape.

The first smoking machines that had some accuracy and reproducibility were developed by Pfy and Bradford et al. in the 1930s (33, 34). Today, commercially available analytic smoking machines, which have flexibility for controlling puffing parameters, are manufactured by various companies (e.g., Borgwaldt GmbH¹⁰ and Cerulean¹¹). The analytic cigarette smoking machines of today vary in the number of ports, how many cigarettes they hold, whether they have ports that are in-line or rotary, and

by their ability to capture mainstream or sidestream smoke. Different smoking machine designs are suitable for different tasks. Rotary machines are ideally suited for smoking a large number of cigarettes quickly (usually the same type or brand) and the smoke is funneled into a single smoke trapping system. One major drawback of the rotary machine is that it cannot easily accommodate modification of the puff interval. Linear smoking machines, on the other hand, are ideally suited for smoking a number of replicates (same or different types) onto individual smoke trapping systems and have more flexibility for altering puff profiles.

Most smoking machines use electric lighters to ignite test cigarettes for machine smoking. However, Adam and coworkers (35) found different yields from the first puff of a cigarette as it is lit, depending on the lighting device. Comparing an electric lighter, a propane/butane gas lighter, a match, a candle, and the burning zone of another cigarette, they found that the three open flame sources produced mainly unsaturated hydrocarbons, whereas the electric lighting device produced oxygen-containing compounds. Therefore, they suggest that the use of electric lighters in smoking machines be reconsidered, because human smokers generally use open flame lighters. Some smoking machines have sensors to determine if the cigarette is lit and they are programmed to stop smoking once the cigarette is smoke down to a specified distance from the end of the filter (e.g., by using a laser detector). Less sophisticated machines rely on a string to mark the stopping point — when the cigarette burns through the string, the puffing mechanism is deactivated.

The Early Development of Standardized Smoking Regimens. The development of smoking machine regimens has been extensively reviewed elsewhere (36-38). In 1936, Bradford et al. (34), who worked for the American Tobacco Company, described the need for standardized smoking parameters that would aid in the characterization and reproducibility of cigarette smoke experiments in the laboratory. However, machine-measured emissions were not widely publicized until the early 1950s (39, 40),

¹⁰ <http://www.borgwaldt.de/cms>

¹¹ <http://www.cerulean.com>

when studies became available linking smoking and lung cancer, and as cigarette manufacturers were racing to produce lower smoking machine tar yield products (commonly called a "tar derby"), making a multitude of inconsistent, noncomparable claims about tar yields to consumers (31). The tar derby ended in 1960 with a voluntary agreement by the FTC and the manufacturers to end tar and nicotine yield claims.¹² The FTC later reversed this agreement and decided to develop a standardized testing method. The initial protocol was largely based on the work of U.S. Department of Agriculture chemist C.L. Ogg in 1964 (31, 41). It seems, however, that this protocol was based on one person's observations about how people smoked, was not determined with some systematic method, and it actually was very similar to the 1936 method of Bradford and coworkers (34). It should be noted that the protocol was not intended to represent the typical smoker, but rather to offer a common basis for a comparison among brands. It should not be inferred that the FTC protocol represents how any person might smoke.

The FTC puffing protocol prescribes drawing a 35-mL puff of 2-second duration every minute until the length of the cigarette is no less than 23 mm for nonfiltered cigarettes or filter overwrap plus 3 mm for filtered cigarettes. Table 1 describes this protocol and others that have been developed over time. The original protocol developed by Ogg et al. (42), also consisted of conditioning of cigarettes at 23.9°C and 60% relative humidity for 24 hours. At the outset, the FTC method was intended only to compare tar and nicotine yields across brands, although carbon monoxide (CO) was added to the protocol in 1980. The analysis of other smoke constituents has never been specified by the FTC, but the FTC protocol has been widely adopted in analyses of other constituents for product testing and research. It also has been widely used for toxicology studies.

Following the work of the FTC, virtually identical standardized smoking regimens were developed by the Cooperation Centre for Scientific Research Relative to Tobacco (CORESTA), and later the International Organization for Standardization (ISO). CORESTA's initial standardized smoking method was published in 1969 (43). The ISO protocol uses the same puffing regimen as the FTC method, except that it specifies an air flow of 200 mL/minute. Additionally, CORESTA and ISO stipulate standards for physical components of the machine: the cigarette holders, smoke traps, ports, channels, and ashtray specifications (43). It should be noted that the tobacco companies heavily influenced CORESTA to motivate ISO to set standards and generate research results in an attempt to preempt regulations (37). Although ISO and CORESTA were seemingly independent, ISO essentially adopted CORESTA's recommended methods, as the ISO committees overseeing standards development for tobacco products have been composed mostly of persons affiliated with the tobacco industry (37, 44).

Human Smoking Behavior

Physical Processes Involved in Smoking. To understand the limitations and misuse of the smoking machine mea-





Figure 1. Schematic representation of the puff and inhalation/exhalation pattern. Reproduced from a BATCo document,¹³ simultaneous measurements were assessed for inhalation/exhalation and smoking topography. The following parameters are defined from this figure: puff volume (mL), integration of puff pressure curve from A to C; lit draw resistance (cm H₂O/mL), the ratio of integrated pressure to puff volume; puff duration, time from A to C; inhalation delay time (s), the time from completion of the puff to the start of inhalation from C to E; inspiratory time (s), the duration of time from E to F; breath hold time (s), the delay from the end of active inhalation to start of exhalation from F to G; expiratory time (s), the time for exhalation from G to H; inhalation volume (mL), the volume difference from E1 to F1; exhalation volume (mL), the volume difference from G1 to H1; volume change before inhalation (mL), volume shift in the lungs (usually exhalation) that occurs during the puff and inhalation delay period from D to E1; volume change after puff (mL), volume change after the puff but before the inhalation, from C1 to E1.

surements, it is important to understand how smokers smoke their cigarettes. The physical process of smoking a cigarette is continuous, but can be divided into three phases: puffing, mouth-holding, and inhalation. The smoking cycle is shown in a diagram reproduced from the British American Tobacco Company (BATCo) research in 1986 (Fig. 1).¹³ This figure defines different parameters that can be measured during smoking. Puffing refers to the act of drawing smoke from the cigarette into the mouth. The act of puffing draws air through the burning rod that causes an increase in temperature that in turn burns some amount of tobacco and the cigarette paper wrap. During puffing, the tongue contracts down creating a negative pressure to aid the puffing process and the soft palate contracts, essentially blocking airflow into the nasopharynx and lungs. Puffing is then followed by a period of mouth-holding before air moves into the lungs, as typically smoke is not directly inhaled from the cigarette through the mouth into the lungs.¹³ Following puffing, as reported via the BATCo documents, the smoke is either

¹² <http://www.time.com/time/magazine/article/0,9171,871506,00.html>

¹³ http://tobaccodocuments.org/bat_cdc/8652.html

Figure 2. Puffing and inhalation patterns showing interindividual variation during the interval between puffing and inhalation.¹⁴ Puffing begins during exhalation from points *a* to *b*. For some smokers, the smoke is immediately inhaled from the mouth into the pharynx and lungs and completed at point *c* (A); for others, there is a mouth-holding period where point *c* marks the beginning of the inhalation and completed at *d* (B); for others, there is an immediate inhalation until point *c* then an exhalation followed by a larger inhalation at point *d* (C); others have an immediate exhalation followed by an inhalation from points *c* to *d* (D).



immediately inhaled via nose inhalation into the lungs, paused in the mouth before nose inhalation (perhaps to enhance the sensation and taste), or paused in the mouth with some exhalation of smoke before nose inhalation. According to BATCo, nose inhalation allows the soft palate to relax, providing an easy path for the smoke to be drawn into the pharynx and nasopharynx.¹⁴ The mouth is closed so that the air pressure sucking the smoke into the lungs is the same as the pressure from air moving from the nose into the lungs. Following nose inhalation, exhalation occurs after some period of time. Puffing resumes after an interpuff interval, and in at least 80% of smokers, this takes place during the exhalation phase of a breath, which can occur at any point during exhalation, e.g., at the onset, in the middle, or at the end of exhalation (45). An example of the various parameters for puffing and inhalation is shown in Fig. 2. Thus, it is the combination of puffing, mouth-holding, nose inhalation, and inhalation time that determines a smoker's internal dose of smoke toxicants and nicotine. These studies only had a few subjects ($n = 8$), measurement was confined to one setting, and this study has not been replicated; a systematic study might show different or more patterns of inhalation, e.g., mouth inhalation in addition to nose inhalation.

Measuring Puff Topography. Puff profiles are measured by assessing smoking topography, namely puff volume, puff duration, interpuff interval, flow rate (sometimes al-

so termed puff velocity), the number of puffs per cigarette, and total puff volume (46-55). These correspond to parameters that can be programmed on a smoking machine. Most commonly, puff topography is measured by having the cigarette smoked through a small tube that can measure air flow via a transducer, and the analog signal is converted to a digital signal for recording and interpretation. Initially, various research groups used their own puff profile recording devices, such as the "tobacco smoke inhalation testing system" originally developed by Puustinen and coworkers in 1986, and then manufactured by the College of Engineering at the University of Kentucky (56-58). Other early techniques included flow meters (59, 60), pneumotachographs (61), pressure transducers and Grass polygraphs (62), and puff analyzers (63). Tobacco industry scientists developed devices in the late 60s and early 70s, such as the cigarette-holder flow meter described by Adams and Creighton (64-66). Portable devices that can be used for at-home monitoring also have been developed (49, 66-69).

Although custom-built apparatuses are still used (70), commercially available topography devices, such as the CRESS units from Plowshare Technologies, Inc.¹⁵ and the SODIM SPA/D and SPA/M smoking topography units¹⁶ have largely superseded them. To our knowledge, there are no published studies using the SODIM topography units and the majority of studies assessing human smoking topography have used the Plowshare

¹⁴ <http://legacy.library.ucsf.edu/tid/zll13f00>

¹⁵ <http://www.plowshare.com/products/index.html>

¹⁶ <http://www.sodim.com/English.htm>

CRess and CRessmicro units (25, 53, 71-80). CRess desktop topography units are capable of real-time recording of individual puffs, including shape and flow rates, for later replication of human puff profiles on smoking machines (portable devices at the present time cannot do this and only provide statistical means of the parameters). The desktop units also contain the ability to cue the smoker for controlled smoking conditions, for example cueing them when to puff, puff duration, and puff volume. These systems also can integrate subjective, performance, and physiologic measures.

Validation of Puffing Topography Recording Devices. Although there are numerous studies about puffing topography, there are few studies that have validated the available methods. Validation would be done in several ways, namely by assessing intraindividual, intralaboratory, and interlaboratory variation, as well as by comparing different methods to assess topography. Published validation studies for intraindividual and intralaboratory methods are limited to the CRess devices. These studies have conducted repeated measures on different days, which assesses both intralaboratory and intraindividual variation measured in the laboratory, and these have generally shown good reproducibility (53, 76, 81). For example, Lee and coworkers (53) found good reliability by intraclass correlation coefficients computed for puff volume (0.66), puff duration (0.75), and maximum puff velocity (0.68). Hammond et al. (82) investigated the smoking topography characteristics with the portable CRessmicro device on 59 subjects smoking an average of 19 cigarettes per day, reporting similar measures of smoking topography for the same subject over time. For a biomarker assessment, in a study of 180 smokers measuring CO and nicotine boosts for two cigarettes 1 hour apart, the use of the topography device did not affect the CO or nicotine levels, because these were measured with and without the topography device in place, although the correlation coefficient with and without the device for CO was only 0.31 ($P < 0.001$).¹⁷

Similar results have been reported in a small study by Lee and coworkers (53). Blank and colleagues (83) recently reported head-to-head comparisons of the desktop and portable CRess devices versus observed smoking using a video tape. The authors found that measured puff duration and interpuff interval on both devices, as well as the video were well correlated (r 's > 0.70), although there were quantitative differences among the devices for puff volume and duration, indicating that comparisons of data across devices might not be reliable. Separately, it has been shown that topography assessments in the laboratory provide similar assessment in the naturalistic environment, e.g., at home (79, 84). Thus, these studies support the reliability and validity of the CRess devices for topography measurement.

What Is Known about Human Puffing Patterns? There is wide interindividual variation for smoking behavior, but a low intraindividual variability because smokers in general show a stable smoking pattern over time, or at least a short period of time (23, 54, 55, 82, 85-88). A clear and consistent finding is that human smoking behavior differs

substantially from the commonly used FTC and ISO parameters, a fact recognized early by the tobacco companies (82, 89-91).¹⁸ Several factors have been documented to influence smoking behavior, such as gender, race, psychological factors, and genetic background. In general, men smoke more cigarettes per day than women and have higher serum cotinine levels (92-96). Although the data are less consistent for smoking topography, men tend to have larger puffs of longer duration, but women may smoke more puffs per cigarettes (52, 70, 71, 97, 98). Differences in smoking topography have been observed between Whites and African-Americans: the latter group generally reflecting greater exposure to smoke toxicants (60, 70, 96, 99). Psychological factors, concurrent use of psychoactive drugs, time of the day and place where a cigarette is smoked also can have an effect on the smoking topography (80, 88, 100, 101, 102).

Generally, there is a high correlation for various puff parameters, e.g., interpuff interval, puff duration, and puff volume; all of these directly impact total puff volume per cigarette (85, 97, 103, 104). However, these parameters are not sufficient proxies for each other and so all need to be recoded when measuring smoking topography (97). Other studies indicate that topography results are not kept constant during the course of smoking a cigarette, where puff volume decreases and interpuff interval initially increases and then decreases (58, 85, 105).

The number of cigarettes smoked per day generally do not relate to puffing topography, or sometimes only is positively correlated with longer interpuff intervals (49, 62, 82, 104, 106). Published studies also are inconsistent for relating puff topography to various biomarkers such as CO and nicotine/cotinine levels, where different parameters affect these biomarkers differently (61, 104, 107-117). For example, puff number and to a lesser extent the puff volume and duration affect nicotine levels, whereas CO levels are mostly influenced by puff volume and less by puff number (112). Zacny and coworkers (61) reported that both nicotine and CO increase proportionally with an increase in puff volume. In a study of 180 subjects, there was a statistically significant correlation for CO boost and puff volume, but not the interpuff interval.¹⁷

Product Design Effects on Smoking Topography. Cigarette design characteristics affect puffing topography when smokers first switch, for example by changes in the draw resistance, sensation, and taste (23, 30, 82, 89). Numerous studies indicate that switching from higher to lower yield cigarettes increase topography parameters such as puff volume and puffs per cigarette (61, 82, 86, 110, 118-126), whereas a decrease in puffing intensity or longer time spent on smoking a cigarette takes place when smokers switch to a higher overall yield cigarette, or cigarettes with constant tar but increased nicotine content (56, 110, 127-132). In a 1986 British American Tobacco study,¹³ 19 subjects who were "low" tar (< 10 mg tar yields) and

¹⁷ Shields, Lerman, Pickworth, and Loffredo, unpublished data.

¹⁸ <http://legacy.library.ucsf.edu/tid/xpt60f00>; <http://legacy.library.ucsf.edu/tid/czv24f00>; <http://legacy.library.ucsf.edu/tid/yci66a99/pdf>; <http://legacy.library.ucsf.edu/tid/lsn86a99/pdf>; <http://www.library.ucsf.edu/tobacco/batco/html/13200/13274/index.html>; <http://legacy.library.ucsf.edu/tid/dss00f00>; <http://legacy.library.ucsf.edu/tid/syj51f00>; <http://www.library.ucsf.edu/tobacco/batco/html/6900/6922/index.htm>

Table 2. Smoking parameters, 1986 BAT study (means \pm SD)

Cigarette type	Puff number	Puff volume (mL)	Mouth hold (s)	Inhaled volume (mL)	Exhaled volume (mL)	Inhalation time (s)	Exhalation time (s)	Breath hold (s)
>10 mg tar yield ($n = 11$)	9.4 \pm 2.9	44.9 \pm 12.3	0.49 \pm 0.27	702 \pm 437	577 \pm 329	1.19 \pm 0.29	2.01 \pm 0.76	0.45 \pm 0.48
<10 mg tar yield ($n = 8$)	12.1 \pm 5.6	44.5 \pm 10.9	0.65 \pm 0.39	636 \pm 138	655 \pm 195	1.22 \pm 0.37	2.89 \pm 0.72	0.45 \pm 0.57

NOTE: http://tobaccodocuments.org/bat_cdc/8652.html. Similar smoking parameters were observed in subjects grouped according to cigarette tar yields with a cutoff of 10-mg tar.

“middle” tar (>10 mg tar yields) smokers had similar puff topography and inhalation parameters, as shown in Table 2. The investigators found that puff volume increased when their subjects who were middle tar yield smokers were switched to a low-tar cigarette, but decreased for the opposite test scenario. The investigators concluded that the increased puff volume was due to decreased draw resistance. In this study, inhalation parameters did not change with switching. Studies by Benowitz et al. (133) suggest that during short-term switching studies, smokers that switch to lower yield cigarettes tend to compensate by changing their behavior by smoking more vigorously or by increasing cigarettes per day (13).

An important design feature of lower yield cigarettes is ventilation via holes punched on the filter paper that allow smoke to be diluted with air during puffing. However, some smokers block these ventilation holes by their fingers or lips, which would then result in yields different than predicted by a smoking machine. In a study of smokers who were trained to uniformly smoke with a particular puff profile that restricted the puffs per cigarette and puff frequency, Strasser and coworkers (117) showed that hole blocking resulted in an increase of CO boost, implying an increase of other tobacco smoke constituents. Puff volumes decreased for both cigarettes with 50% hole blocking. Other switching studies reported similar results but differed in the magnitude of the CO response depending on the cigarette type that was smoked, namely the effects are greatest for ultralight smokers (116, 134). One explanation for the difference in results might be the lack of controlling for puff number and puff interval; in the latter two studies, there were many more puffs per cigarettes that might have obscured a difference. Regardless, it is clear that smoking machine studies that compare cigarettes with different physical design characteristics using the same puffing profile fail to accommodate for what happens to smokers who switch or naturally adopt one product versus another.

Filter efficiency is affected by puffing. Increasing smoke flow through the filter, such as with greater puff volumes and decreasing filter ventilation, but not so much decreasing puff frequency, will tend to decrease filter efficiency, leading to a narrower range of yields across brands.¹⁹ For example, Marlboro UltraSmooth (MUS) with a novel carbon filter is much less effective in reducing toxic smoke constituents when smoked under the HC regimen compared with the FTC method (135).

For many PREPs, design features are varied and switching studies show that smoking behavior changes (summarized in Table 3). For example, smokers who

switched to the Advance cigarette that has a modified filter took fewer puffs and had higher nicotine levels, whereas the rest of puffing characteristics remained unchanged (136, 137). Two studies investigating the Accord electronic smoking system found that subjects had shorter puff intervals and fewer puffs per cigarette, because this is electronically controlled, and higher puff volume and duration compared with smoking own brand cigarettes (72, 74). Eclipse smokers, which is a product designed to heat tobacco rather than burn it, substantially increased their puff volumes, and decreased the interpuff interval (138-140). For Eclipse, CO levels also increased, and for some smokers, the levels were quite high (141). Acrolein also is increased. For Quest cigarettes that vary in nicotine yields, there is compensatory smoking with an increase in the total puff volume and CO boost (115). Another study reported that switching to Omni cigarettes that had a modified filter resulted in fewer puffs compared with the usual brand, but there also was an increase in CO boost and not a significant decrease in carcinogen exposure when compared with conventional cigarettes (75). When comparing MUS that had a modified filter with charcoal particles embedded in cellulose acetate, with two conventional cigarettes (Marlboro Lights and Ultralights), investigators observed a decrease in number of puffs, but higher puff volumes (79). The overall conclusion of the study was that there is no significant change in smoking topography between the MUS and conventional cigarettes; therefore, there will be no reduced exposure among smokers that switch from a conventional brand. Thus, smoking machine studies that compare PREPs to conventional products using the same puffing profile could be misleading in terms of relative effects.

Measuring Inhalation and Exhalation. Smoking behavior also involves not only assessing puffing behavior, but also inhalation, which more closely relates to biological dose. Several techniques have been developed for measuring times and volumes for inhalation and exhalation. Some early methods were reviewed in a report from Imperial Tobacco Ltd.²⁰ These techniques are summarized in Table 4. The main conclusion was that these devices were accurate in measuring the physical mechanics of inhalation and exhalation, but they did not permit studies in the naturalistic setting and they imposed restrictions on free smoking behavior. Tobin and coworkers (142) used chest plethysmography, to assess the pattern of inhalation in smokers and then compared this with the smokers' subjective reports for inhalation. They found that smokers inaccurately perceived their inhalation patterns. In another

¹⁹ <http://legacy.library.ucsf.edu/tid/cbi31d00> and <http://legacy.library.ucsf.edu/tid/htu61e00>

²⁰ <http://legacy.library.ucsf.edu/tid/kjn70f00>

Table 3. Selected smoking topography characteristics among PREP studies

Author/year	Products	Participants (n)	IPI (s)	Puff numbers	Puff volume (mL)	Total puff volume (mL)
Breland AB 2003 (136)	Advance Own	12 (8F, 4M)		9.6 (2.8) 11.7 (4.2)		
Breland AB 2002 (137)	Advance Own Sham	20 (10F, 10M)	34.5 (21.9) 33.9 (23.6) 17.2 (14.2)	51.6 (9.4) 56.5 (11.2) 66.5 (43.7)		
Buchhalter R 2000 (72)	Accord Own	10 (7F, 3M)	24.0 (12.1) 35.0 (17.9)	7.8 (0.7) 10.3 (2.1)	55.4 (17.0) 38.4 (11.7)	432.12 395.52
Breland AB 2006 (138)	Eclipse Own	35 (8F, 27M)	21.38 30.74	17.03 10.03	65.01 50.97	1,107.12 511.22
Slade J 2002 (139)	Eclipse Own		19.7		67	1371 640
Breland AB 2002 (74)	Eclipse Accord Own	20 (10F, 10M)			53.3 (4.3) 61.8 (4.8) 49.8 (3.3)	
Strasser AA 2007 (115)	Quest 0.05 Quest 0.3 Quest 0.6 Own	50	18.6 19.6 21.6 21.6	10 9.9 9.8 14.3	59.4 55.9 58.1 60.5	570.5 (156.9) 518.1 (145.6) 540.3 (144.9) 832
Rees VW 2008 (79)	ML MUS MUL ML MUS MUL	32 (21F, 11M) Tampa Salt Lake City	32.1 (11.9) 28.4 (8.4) 33.0 (14.9) 24.8 (11.45) 23.9 (9.2) 21.8 (10.4)	11.4 (3) 10.2 (2.9) 11.3 (2.6) 13.1 (4.8) 12.5 (5) 13.9 (5.6)	50.7 (19.6) 54.2 (19.4) 51.4 (19) 47.4 (16.9) 56.7 (15.2) 50.1 (15.6)	578 552.84 580.82 620.94 708.75 696.39
Lee EM 2004 (140)	Eclipse Own	10		16.1 (2.1) 11.5 (0.7)	89.3 (10.8) 60.1 (4.0)	1,437.73 691.15
Hughes JR 2004 (75)	Omni Own	34		11.6 (0.5) 12.7 (0.7)	49 (2) 50 (2)	547 (25) 612 (34)

NOTE: Data presented as mean (SD) or mean only as available in the original paper. Empty cells mean no value exists in the paper for that parameter. Calculated values are presented in *italics*.
Abbreviations: ML, Marlboro Light; MUL, Marlboro Ultra Light; MUS, Marlboro UltraSmooth; IPI, Interpuff interval; F, Females; M, Males.

study, Tobin and Sackner (128) used the same system to assess switching from high to low-tar cigarettes, showing that there was no change in the inhalation characteristics.

The most widely used device by the tobacco industry to assess smoke inhalation by inductive respiratory plethysmography has been the RespiTrace, developed for assessing respiratory function and disease (NonInvasive Monitoring Systems, Inc.; ref. 143). The system consists of insulated coils enclosed in elastic bands applied on the rib cage and abdomen of the subject, registering the changes in respiratory movements that alter the self-inductance of the coils. The device must be calibrated for tidal volume with the use of a spirometer (144). BATCo used the RespiTrace system in the studies discussed above to discern the physical process of smoking.²¹ Research has been conducted to assess whether smoking machine tar and nicotine yields affect inhalation in two studies, but one reported no effect and the other found a positive relationship (122, 145).

The effects of inhalation on dose measured via biomarkers has received little attention. Zacny and coworkers trained smokers to smoke their cigarettes according to a controlled smoking regimen for inhalation depth and time (61). They measured CO and nicotine boosts, and showed that postpuff inhalation volume and duration under *ad libitum* and controlled smoking conditions had no effect on the CO and nicotine levels. Similarly, Herning and coworkers (103) found that nicotine blood levels were not related to inhalation. In a third study, nicotine retention was almost complete even at low inhalation volumes

(146). These studies indicate that nicotine absorption is very quick and so unrelated to inhalation, but it may be that other tobacco smoke constituents would be affected by inhalation. This has received even less attention, but one study has reported that the retention of solanesol was related to inhalation volume (146). In a study by Philip Morris scientists, a novel method was used to measure the estimated intake into the lungs by having smokers exhale through a Cambridge filter pad (147). The difference between the estimated chemical yield, as measured by a smoking machine, and the amount of the chemical constituent on the pad was considered retained in the smoker. Under controlled smoking conditions where the smokers varied their depth of inhalation, they found similar results as above for no relation of inhalation to nicotine retention (61), but that the retention of tobacco-specific nitrosamines was greater with deeper breaths. For the gas vapor phase, however, depth of inhalation had little effect on retention. Thus, inhalation can be an important parameter for some smoke constituents such as tobacco-specific nitrosamines.

In summary, smoking behavior is complex and many of the individual components covary, so that affecting one might affect each other. These are directly affected by cigarette designs. However, the various aspects of smoking also affect smoking machine yields and smoke toxicant effects, as indicated below. Some parts of human smoking are not captured at all by the smoking machine, whereas some variables such as puff velocity and puff shape are usually not considered. Smokers vary their puffing behavior during the course of their cigarette, by day, and by who they are. These added variables make it impossible to replicate a typical smoker using one smoking machine regimen.

²¹ <http://legacy.library.ucsf.edu/tid/zll13f00>; http://tobaccodocuments.org/bat_cdc/8652.html

Smoking Machine Profiles and Mimicking Human Smoking Behavior

As evidence accumulated that smokers' behaviors and exposures were distinct from machine-measured yields, increased interest was placed on altering machine smoking methods to better reflect smoker practices. The 1981 Surgeon General Report, for example, acknowledged that the FTC testing method needed to account for compensatory smoking (via larger and more frequent puffs) and ventilation hole blocking (16). A National Cancer Institute *ad hoc* expert committee convened in 1994 came to similar conclusions (88). Research on alternative testing regimens was ongoing in the public health/regulatory community. For example, Rickert and coworkers (148) tested smoke yields under ISO conditions and two more intensive conditions and reported that the yields of tar, nicotine, and CO more than doubled when cigarettes were smoked under the intensive regimens compared with the standard one. Djordjevic et al. (109) determined the actual human puff profiles of

133 smokers and replicated the profiles of a randomly chosen subset of 72 on the smoking machine. The investigators found that the yields of tobacco-specific *N*'-nitrosamines and benzo(*a*)pyrene (BaP) increased by 2-fold, whereas the nicotine and tar levels increased >2-fold compared with the FTC measures yields.

In 1996, the Massachusetts Department of Public Health (MDPH) Tobacco Control Program began a research project to establish a machine smoking regimen that more resemble human smoking. Initially, two sets of smoking regimens were chosen, derived from 32 studies on *ad libitum* smoking topography presented in the 1988 Surgeon General's report (149). One was termed the "average smoker" protocol and the other a more intense "heavy smoker" protocol. The former had a 45-mL puff volume every 30 seconds, with a puff duration of 2 seconds and taping closed 50% of the ventilation holes. The MDPH 50% hole blocking in particular was recommended in the context that smokers will block ventilation holes when they smoke, for example with their fingers or

Table 4. Summary of inhalation/exhalation monitoring methods used for assessing human smoking behavior (excerpted from internal company documents)

Author/year	Method	Variables measured	Limitations
Cinkotai F.F. 1967*	Partial body plethysmography Puff volumes and duration determined with a modified cigarette holder as a flow meter	Volume of the puff Duration of the puff Holding time of the puff in the mouth Lung volume at the beginning of the puff Time of inhalation Volume of air inhaled with the puff Volume of exhaled air Time of exhalation.	Discomfort leading to high puff by puff variation observed in the breathing patterns of individual smokers and abnormal tidal breathing caused by stress
Creighton D.E. 1978 (66)	Impedance pneumography Puff profiles and puff volumes measured with a special cigarette holder and a pressure transducer	Puff profiles and puff volumes Semiquantitative estimates of breathing patterns	Needs calibration against a partial body plethysmograph before each use. Nonlinear response and day-to-day variations for individual and variation between subjects.
Guillerm R. and Radziszewski E. 1975 (170)	The Guillerm and Radziszewski Method A flow meter constructed from a classic cigarette holder with a bead placed between the two snap-in-parts of the holder connected by flexible polyvinyl tubing to a differential transducer. A special IR pyrometer used to measure the temperature variations of the combustion cone of the cigarette.	Puff volume and duration No. of and intervals between puffs Volume of air taken between puffs Volume of air inhaled immediately after the puff Location of the puff in the ventilatory cycle The breathing pattern was measured at the same time as the puff analysis	The puff volume recorded did not always correspond to the true inhaled puff volume and the technique imposed some physical restrictions on the subject, particularly concerning the cigarette holder.
Rawbone R.G. 1978 (171)	Mercury strain gauge chest pneumography The puff parameters were obtained from measurements of the pressure drop across a small resistance inserted between the cigarette and the smoker. The depth of inhalation was measured by recording movements of the chest wall with a mercury strain gauge chest pneumogram.	Puff volumes, Puff duration Interpuff interval Semiquantitative estimates of breathing patterns	Calibration was required before each study.
Sackner M.A. 1980 (172) Tobin M.J. 1982 (128, 142)	Respiratory inductive plethysmography Consisting in two coils of Teflon-insulated wire, which were sown into elastic bands encircling the rib cage and the abdomen and connected to an oscillator module. Tidal volume measured by spirometry.	No. of puffs Puff duration, Puff volumes, Integrated puff pressure. Accurate estimation of breathing patterns	Accuracy of the results depended on the initial calibration and the stability of the calibration during changes in body positions and lung volumes.

NOTE: <http://legacy.library.ucsf.edu/tid/zll13f00>; <http://legacy.library.ucsf.edu/tid/kjn70f00>.

*<http://legacy.library.ucsf.edu/tid/hos00f00>

Table 5. Average smoking topography parameters values of 2,432 subjects compared with the ISO/FTC parameters (modified from the reviewed by the WG 9 of the ISO TC 126) ref. 154

		ISO/FTC	HPPs grouped by tar yield (mg)			
			≥14	8-14	3-8	<3
Puff characteristics*	Puff volume (mL)	35	48.1 (10.7)	47.8 (6.3)	54.7 (9.7)	57.2 (8.9)
	Puff duration (s)	2	1.9 (0.4)	1.8 (0.3)	2 (0.3)	1.9 (0.1)
	Puff interval (s)	60	26.1 (8.8)	27.3 (8.7)	22.6 (7.1)	18.9 (0.7)

NOTE: The HPP puff volumes are higher and puffs are drawn at less than half the interval of the ISO/FTC parameters. Also, the HPP puff volumes increase and the puff intervals decrease corresponding to the decrease in tar yields.

*Mean (SD) values for HPPs.

lips (20, 21, 91, 150, 151). The initial proposal also included an “intense,” or “heavy,” smoking condition (60-mL puff every 26 seconds, 100% vent blocking), but this was dropped from the final plans. From 1997, cigarette manufacturers have been required to report results to the MDPH under the “average” protocol, along with levels of filter ventilation, tobacco nicotine content, and smoke “pH” (152). It should be noted that derivation of average and intense smoking for this protocol reflected topographical data available before 1988 and not necessarily reflective of today’s products’ design and smokers’ behavior.

In the same year (1996) as the MDPH protocol was being adopted, Health Canada (HC) began work on amending its tobacco regulatory authority and convened an Expert Committee on cigarette modifications. Discussions on reducing the harmfulness of cigarettes led to a formal exploration of alternative smoking conditions (153). In this report, Rickert (153) noted that puff volume and inter-puff interval are the key variables to consider in a new machine smoking regimen. This resulted in the proposal of an HC protocol with a 56-mL puff volume of 2-second duration and a 26-second inter-puff interval; the ventilation holes would be fully blocked. Other elements of the ISO protocol (conditioning, duration, butt length) were retained. The 100% hole blocking was adopted to directly compare the performance of cigarettes removing the strongest predictor of tar and nicotine yields. The report concluded that testing under two conditions (ISO and HC) would be sufficient to capture the range of deliveries that might be experienced by smokers [and later adopted by the WHO Study Group on Tobacco Product Regulation (ref. 10)]. In June 1998, the Health Protection Branch of HC outlined proposed reporting requirements of 40 constituents in mainstream smoke based on the standard and extreme regimens. The Tobacco Act of 2000 in Canada made the new regimen official. During the regulatory process, the parameters were changed to a 55-mL puff volume of 2-second duration and a 30-second puff frequency (149).

In 2004, an ISO Working Group (ISO/TC126/WG9) was convened to craft an alternative smoking regimen that more closely hewed to human smoking behaviors (149). The ISO was faced with the overwhelming evidence that the ISO/FTC regimen inadequately characterized modern cigarette exposures, that there were emerging test methods in different jurisdictions, and the prospect of impending regulations under Articles 9 and 10 of the Framework Convention on Tobacco Control.²² The group, which included members affiliated with the tobacco in-

dustry, reviewed published literature on smoking topography from 1956 to 2004, and used 100 data sets comprising 2,432 subjects (154). They derived summary statistics for puff volume, duration, interval, number of puffs per cigarette, and how these vary with cigarette tar yield as determined by the ISO/FTC smoking regimen. Significant differences were noted between the experimentally derived average human puffing profiles (HPP) and the ISO/FTC parameters, as summarized in Table 5. Ultimately, the Working Group proposed three different smoking machine protocols for testing, as shown in Table 1. These were determined by grouping the HPPs according to machine-smoked ISO/FTC cigarette “tar” yield ranges. The activities of Working Group 9 were set aside in May 2006 and Working Group 10 was established. The activities of Working Group 10 are ongoing as a forum for exchange of information between WHO (the public health sector in general) and the tobacco industry scientists. For this, and given that laboratories affiliated with tobacco companies are precluded from participating in the validation work of TobLabNet, the Working Group 10 is an important forum. No tangible products have yet come out of the Working Group 10 because the purpose to date is for information exchange.

In 1997, the FTC announced plans to revise its cigarette testing method with a public comment period (Federal Register 62/177, 9/12/97). In addition to the standard method, a more intense method was being considered (a 2-second, 55-mL puff every 30 seconds). However, no action was taken at that time. Later, in 2008, the FTC proposed rescinding in its entirety their 40-year guidance for smoking machine testing, rather than recommending a second and more intense puffing regimen. The Agency stated: “Today, however, the scientific consensus is that machine-based measurements of tar and nicotine yields based on the Cambridge Filter Method do not provide meaningful information on the amounts of tar and nicotine smokers receive from cigarettes or on the relative amounts of tar and nicotine they are likely to receive from smoking different brands of cigarettes. The primary reason for this is smoker compensation—that is, smokers alter their smoking behavior to obtain the necessary nicotine dosage.”²³ After a 60-day public comment period, the FTC followed-through and rescinded its guidance, drawing the era of “FTC” yields to a close.

Changes in Yields by Smoking Regimen. Changing specific parameters of the puff profile independently can directly

²² <http://www.who.int/ftc/en>

²³ <http://www2.ftc.gov/opa/2008/07/cigarettefyi.shtml>

affect smoke yields. For example, decreasing puff volume, increasing puff frequency (decrease interpuff interval), and increasing filter ventilation decrease tar and chemical yields on a per cigarette basis (155).²⁴ In smokers, though, using higher ventilated cigarettes generally results in larger puff volumes. Toxicology studies also show the influence of puff volume, ventilation, and ventilation hole blocking (156).²⁵ The ISO/FTC, MDPH, and HC methods use different puff volumes, puff frequency, and ventilation hole blocking, and increases in these variables result in increased tar, nicotine, and other constituent yields on a per cigarette basis (109, 148, 157, 158).

The data indicate that the relative rankings of different products, on a per cigarette basis, will generally be preserved across regimens although the gap in toxicant emissions with more intense protocol is reduced. Counts and colleagues from Philip Morris (157) published a large survey of emissions from international brands tested under ISO, MDPH, and HC conditions, showing that the ratios of constituents to total tar were dependent on the puffing profile, and mostly driven by filter ventilation. For example, when cigarettes were grouped broadly by filter ventilation, the yields of individual constituents relative to tar changed differently as the different profiles were compared. This effect was greater for vapor phase compared with particulate phase constituents. However, the effect was least for the cigarettes with lower ventilation and higher tar yields. Separately, Hammond and O'Connor (155) examined the relationships between yields under the ISO and HC regimens for the 2004 Canadian market and showed that the increased intensity of the HC system changes the absolute concentrations of constituents, but also their concentrations relative to nicotine (157).

Both Philip Morris and RJ Reynolds Tobacco companies, as early as 1974, developed the capability to capture human topography data and mimic this on a smoking machine, and it was shown that the yields predicted for different smokers substantially varied among them, and higher than the FTC predicted yields.²⁶ In a 1982 report by RJ Reynolds' scientists, an analysis indicated that using five variables within the puff for flow velocity at different times of the puff and the time to reach V_{max} , six types of profiles could be described.²⁷ Although, each smoker would vary their shape within a cigarette, it was reported that 12 patterns would characterize all 550 smokers.

Whether changing the shape of the puff affects yields is unclear, and there are no recently published studies. A 1968 report from Brown and Williamson showed that when air flow peaked (early versus late), different yields were obtained.²⁸ The parameters that affect the shape of the puff or the variability for the puff-by-puff profile are unknown, but it seems that filter ventilation does not af-

fect the latter.²⁹ None of the above studies, however, measured specific chemical constituents. New commercial topography devices have the capability to record puff-by-puff data, including the change of airflow within a puff. Today, smoking machines also can be programmed with the use of specialized pumps and software to better replicate the human-type puff on a puff-by-puff basis. However, whether this new technology affects the smoke yields and provides for a better replication of human smoking remains to be determined.

A more meaningful comparison might come from an assessment using different puff profiles tailored to the product as it might be used by smokers. For example, a method has been proposed based on nicotine yields by Kozlowski and O'Connor (91). They proposed a two-step system where the first step would use the traditional ISO/FTC yield on a per cigarette basis, whereas a second step would use puff parameters adjusted to yield the same nicotine levels, for example by adjusting the puff volume. Later, Hammond and colleagues revised this recommendation to propose a system whereby puffing profiles would be iteratively adjusted so that all brands yielded a specific nicotine level. In both cases, the goal would be to better simulate compensatory smoking by humans within the limitations of machines. However, little work has been done to operationalize these methods. It should be noted that the above methods adjust smoking machine parameters based on total cigarette yields and not on a per milligram of tar basis and assumes that the chemical composition of tars are similar; however, this is known not to be true (157-159).

Hammond and coworkers (30) compared the smoke yields produced under ISO, MDPH, HC, and the two-stage compensatory regimen described above to the average of actual topography measures for 51 smokers using their usual brand and 21 who switched to ultralights (human mimic profiles). Ventilation hole blocking was 50% for the MDPH and compensatory and human mimic profiles, whereas it was 100% for the HC method. None of the yields for the four smoking regimens replicated the human mimic profiles. Tar, nicotine, and CO yields obtained for the regular tar smokers under the mimic protocol were double of those obtained with the ISO and compensatory regimens, but lower than the HC regimen. For the ultralight switchers, the human mimic yields were three to four times greater than the ISO and MDPH regimens, but slightly lower than the HC regimen and similar to the compensatory regimen. Importantly, none of the standardized machine determined nicotine yields predicted levels of salivary cotinine, except for the human mimic regimen. Thus, it is likely that no single smoking regimen can adequately characterize smoking.

Philip Morris has proposed another method for comparing products, which is to characterize human smoking behavior on a smoking machine based on several regimens statistically modeled based on topography data and urinary nicotine metabolites (160). The method uses the determined 10th percentile, mean, and the 90th percentile of the puff volumes, and the other parameters

²⁴ <http://legacy.library.ucsf.edu/tid/aob34c00>; <http://legacy.library.ucsf.edu/tid/cbi31d00>; <http://legacy.library.ucsf.edu/tid/qtp03f00>; <http://legacy.library.ucsf.edu/tid/rto73d00>; <http://legacy.library.ucsf.edu/tid/gkb11d00>

²⁵ <http://legacy.library.ucsf.edu/tid/mxa35d00>; <http://tobacodocuments.org/rjr/508352445-2461.html>

²⁶ <http://legacy.library.ucsf.edu/tid/fgo46b00>; <http://legacy.library.ucsf.edu/tid/pg81b00>

²⁷ <http://legacy.library.ucsf.edu/tid/irp93a00>²⁸ <http://legacy.library.ucsf.edu/tid/rwu69d00>; <http://legacy.library.ucsf.edu/tid/bmp84a99>

²⁹ <http://legacy.library.ucsf.edu/tid/aob34c00>

were modeled. Thus, they proposed testing cigarettes with a low (25-mL puff volume, 0.8-second puff duration, 2.4/minute puff frequency), a medium (48-mL puff volume, 1.3-second puff duration, 1.8/minute puff frequency), and a high (65-mL puff volume, 1.6-second puff duration, 1.9/minute puff frequency) puffing profile. However, we are unaware of any actual implementation of this proposal.

Data on comparative emissions for PREPs are rare. A specific example of a PREP for the utility of testing under multiple smoking machine methods is the Eclipse cigarette, which is claimed to heat rather than burn tobacco under the FTC conditions. When smoked on a machine in a way more similar to what smokers do, the tobacco becomes significantly charred and the smoke chemistry differences compared with conventional cigarettes become much less.³⁰

Discussion

The need for validated laboratory methods to assess tobacco smoke for chemical constituents and toxic effects has recently been underscored by the new FDA authority to enact product performance standards and evaluate manufacturer health claims for modified tobacco products. Prior uses of smoking machine results led to misinterpretations and misunderstandings about cigarette comparisons and their relationship to human health (13, 27). As a result, smoking machine data are regarded as poor indicators of health risk, leading the FTC to rescind its imprimatur from the method.⁷ However, smoking machines will continue to be used for laboratory screening of product design changes and the assessment of performance standards, and so better methods need to be developed (10, 28). Critical to the development of new smoking machine methods is a better understanding of human smoking behavior, including how interindividual variation in puffing, mouth-holding, and inhalation affect exposure. These studies can then inform the use of cross-regimen comparisons, for example as previously described (30, 91, 160), which may better reflect the differences among human exposure for specific product comparisons about product design. This would lead to tailoring puff profiles to particular products as used by smokers. Thus, there are several research gaps that need to be addressed to maximally apply and interpret smoking machine studies.

Currently, almost all methods for assessing human puffing that can be extrapolated to smoking machine protocols is through commercial topography devices. Data from such studies suggest that topography may differ by gender and race (52, 60, 70, 71, 92-99). However, there are many other likely determinants, such as age, comorbidities, prior smoking history, nicotine metabolism, genetics, and psychological factors that have been studied even less in the context of topography and application to smoking machine studies (88, 100, 101, 161-169). Other variables include smoking environment at time of measurement (naturalistic versus laboratory), time of day,

and circadian rhythms (80, 102). Without a better understanding of how much these variables affect the range of human exposures, it will be difficult to know if future smoking machine regimens are sufficiently mimicking human exposure.

Although there are some data demonstrating the replicability of smoking behavior using these devices (53, 76, 81, 82), additional studies are needed to compare different commercial units and to validate them. It is unknown if these devices are measuring accurately air flow and volumes, and so a major limitation for validating topography measurements is the comparison to some "gold standard." But none exists. Validation of topography as an indicator of exposure requires statistically significant and consistent correlations with biomarkers that have been validated for smoking (141), but the data thus far for comparing topography to biomarkers have produced conflicting results. Biomarkers of exposure reflect not only puff topography but also mouth-holding and inhalation, and so it may be that a biomarker comparison is not valid, assuming that varying mouth-holding and inhalation affect the dose to smokers. Thus, additional studies are needed to assess mouth-holding and inhalation to determine how much, if any, these components of smoking affect exposure. However, methods to assess mouth-holding and inhalation are poorly developed, and so better technologies are needed that can be applied to human studies. Once developed, controlled smoking and cross-sectional studies can be conducted with biomarkers to determine how much puffing, mouth-holding, and inhalation contribute to variance in human smoke exposure.

The current designs for smoking machine puffing profiles have been developed considering each parameter as independent effects, but changing one actually influences the others (58, 85, 105). So a better understanding of the impact of changing one parameter on others is needed, both for topography and for smoking machine studies. It is known that many of the various topography parameters covary (85, 97, 103, 104), but a systematic study has not been done to identify the extent of this. Another parameter that is only partially characterized is blocking ventilation holes (54, 116, 117, 134). To determine how people block holes, how much, and how often has been insufficiently studied. However, filter ventilation affects smoke dilution and puff volume, and so is critical for determining smoke yields. Thus, better technologies are needed to determine ventilation hole blocking and incorporate them into human studies that assess topography.

There is sufficient data to know that different machine puff profiles cause cigarettes to burn differently and have different chemical yields and biological activity, and that this would also result in different exposures in humans (157, 158).³¹ Thus, smoking machines need to better mimic human smoking, including methods to replicate puff-by-puff parameters, and studies need to be done to determine if the shape of the puff significantly affects yields. How to model the diversity of human smoking behavior needs to be developed. Then, better methods to compare cigarettes

³⁰ <http://legacy.library.ucsf.edu/tid/kqm60a99>; <http://legacy.library.ucsf.edu/tid/pkd56a00>

³¹ <http://tobaccodocuments.org/rjr/508352445-2461.html>; <http://legacy.library.ucsf.edu/tid/cbi31d00>; <http://legacy.library.ucsf.edu/tid/aob34c00>; <http://legacy.library.ucsf.edu/tid/qtp03f00>; <http://legacy.library.ucsf.edu/tid/rto73d00>; <http://legacy.library.ucsf.edu/tid/gkb11d00>; <http://legacy.library.ucsf.edu/tid/mxa35d00>

and product design changes through cross-regimen comparisons are needed. Whether this is done using topography data or by standardizing for nicotine yields needs to be developed, and there is sufficient rationale to indicate that both methods might have utility.

Since the passing of the FDA legislation and the Institute of Medicine report conclusion that risk reduction through PREPs is a feasible approach (11), a comprehensive framework for studying tobacco products, including PREPs, is needed. This would include studies ranging from premarket assessments using laboratory studies to population surveillance. It would use integrative approaches by examining individual smoking behavior for new products and establish their relationship with actual delivered dosages of nicotine and a select panel of toxic and carcinogenic agents. An iterative process would therefore be used, where product design changes are tested first in the laboratory for increases in smoking yields and toxicity, followed by human use in short-term studies, and then replication of human use in the laboratory to confirm the yield and toxicity changes. Central to this process is the understanding of human smoking behavior and how to replicate this in the laboratory, but current knowledge and methods are insufficient to do this. Additional research, however, can fill in the research gaps to improve tobacco product assessment. Having validated methods for assessing tobacco products in the laboratory is vital for the fulfillment of the promise of regulatory oversight to protect the public health.

Disclosure of Potential Conflicts of Interest

P.G. Shields serves as an expert witness in tobacco litigation cases on behalf of plaintiffs. D.K. Hatsukami has received grants from NabiBiopharmaceuticals to undertake clinical trials on the nicotine vaccine.

Acknowledgments

We thank Dr. William Farone (Applied Power Concepts, Inc., Anaheim, CA) for his insightful comments on this manuscript.

References

- Warner KE. Tobacco harm reduction: promise and perils. *Nicotine Tob Res* 2002;4 Suppl 2:S61-71.
- Britton J. Smokeless tobacco: friend or foe? *Addiction* 2003;98:1197-207.
- Ramstrom L. Snus: part of the problem or part of the solution? *Addiction* 2003;98:1198-9.
- Fagerstrom KO, Schildt EB. Should the European Union lift the ban on snus? Evidence from the Swedish experience. *Addiction* 2003;98:1191-5.
- Pierce JP. Harm reduction or harm maintenance. *Nicotine Tob Res* 2002;4:S53-4.
- Fox BJ, Cohen JE. Tobacco harm reduction: a call to address the ethical dilemmas. *Nicotine Tob Res* 2002;4:S81-7.
- Hatsukami DK, Slade J, Benowitz NL, et al. Reducing tobacco harm: research challenges and issues. *Nicotine Tob Res* 2002;4 Suppl 2: S89-101.
- Gartner CE, Hall WD, Chapman S, Freeman B. Should the health community promote smokeless tobacco (snus) as a harm reduction measure? *PLoS Med* 2007;4:e185.
- Foulds J, Kozlowski L. Snus-what should the public-health response be? *Lancet* 2007;369:1976-8.
- WHO Study Group on Tobacco Product Regulation Report on the Scientific Basis of Tobacco Product Regulation. WHO Technical Report Series, no. 951; 2008, Report No.: 951.
- Institute of Medicine Committee to Assess the Science Base for Tobacco Harm Reduction and Board on Health Promotion and Disease Prevention. *Clearing the Smoke: Assessing the Science Base for Tobacco Harm Reduction*. Washington, DC: National Academy Press; 2001.
- Stratton K, Shetty P, Wallace R, Bondurant S. Products for tobacco exposure reduction. *Clearing the SMOKE: Assessing the Science Base for Tobacco Harm Reduction*. pp. 82-92: National Academy Press (Institute of Medicine); 2001.
- National Cancer Institute. Risks Associated with Smoking Cigarettes with Low Machine-Measured Yields of Tar and Nicotine. Smoking and Tobacco Control Monograph No. 13 2001. U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute, NIH Pub No.02-5074. Ref Type: Serial (Book Monograph).
- Cohen JB. Smokers' knowledge and understanding of advertised tar numbers: health policy implications. *Am J Public Health* 1996;86:18-24.
- US. Department of Health and Human Services. Reducing the Health Consequences of Smoking: 25 Years of Progress. 1989, Report No.: DHHS Publication No. (CDC) 89-8411.
- US. Department of Health and Human Services Public Health Service Office on Smoking and Health. *The Health Consequences of Smoking: the Changing Cigarette*, a report of the Surgeon General. 1981.
- United States District Court for the District of Columbia. Final Opinion on U.S. Department of Justice v. Philip Morris Inc, et al. Civil Action No. 99-2496. 2006.
- United States Court of Appeals. Unanimous Opinion on U.S. Department of Justice v. Philip Morris Inc. et al., No. 99cv02496. 2009.
- Stellman SD, Muscat JE, Hoffmann D, Wynder EL. Impact of filter cigarette smoking on lung cancer histology. *Prev Med* 1997;26:451-6.
- Kozlowski LT, Mehta NY, Sweeney CT, et al. Filter ventilation and nicotine content of tobacco in cigarettes from Canada, the United Kingdom, and the United States. *Tob Control* 1998;7:369-75.
- Kozlowski LT, Rickert WS, Robinson JC, Grunberg NE. Have tar and nicotine yields of cigarettes changed? *Science* 1980;209:1550-1.
- Benowitz NL, Hall SM, Herning RI, et al. Smokers of low-yield cigarettes do not consume less nicotine. *N Engl J Med* 1983;309:139-42.
- Benowitz NL. Compensatory Smoking of Low-Yield Cigarettes. In: US. Department of Health and Human Services NNCI, editor. Smoking and tobacco control monograph no. 13. Risks associated with smoking cigarettes with low machine-measured yields of tar and nicotine. Bethesda, MD: US Department of Health and Human Services NIH, National Cancer Institute; 2001, p. 39-63, Ref Type: Serial (Book Monograph).
- Scherer G, Engl J, Urban M, et al. Relationship between machine-derived smoke yields and biomarkers in cigarette smokers in Germany. *Regul Toxicol Pharmacol* 2007;47:171-83.
- Mendes P, Kapur S, Wang J, Feng S, Roethig H. A randomized, controlled exposure study in adult smokers of full flavor Marlboro cigarettes switching to Marlboro Lights or Marlboro Ultra Lights cigarettes. *Regul Toxicol Pharmacol* 2008;51:295-305.
- Hecht SS, Murphy SE, Carmella SG, et al. Similar uptake of lung carcinogens by smokers of regular, light, and ultralight cigarettes. *Cancer Epidemiol Biomarkers Prev* 2005;14:693-8.
- Thun MJ, Burns DM. Health impact of "reduced yield" cigarettes: a critical assessment of the epidemiological evidence. *Tob Control* 2001;10 Suppl 1:i4-11.
- Burns DM, Dybing E, Gray N, et al. Mandated lowering of toxicants in cigarette smoke: a description of the World Health Organization TobReg proposal. *Tob Control* 2008;17:132-41.
- Hammond D, Wiebel F, Kozlowski LT, et al. Revising the machine smoking regime for cigarette emissions: implications for tobacco control policy. *Tob Control* 2007;16:8-14.
- Hammond D, Fong G, Cummings KM, et al. Cigarette yields and human exposure: a comparison of alternative testing regimens. *Cancer Epidemiol Biomark Prev* 2006;15:1495-501.
- National Cancer Institute. The FTC Cigarette Test Method for Determining Tar, Nicotine, and Carbon Monoxide Yields of U.S. Cigarettes. 1996, US Department of Health and Human Services, National Institutes of Health, National Cancer Institute, NIH Pub No.96-4028. Ref Type: Serial (Book Monograph).
- Johnson MD, Schilz J, Djordjevic MV, Rice JR, Shields PG. Evaluation of *in vitro* assays for assessing the toxicity of cigarette smoke and smokeless tobacco. *Cancer Epidemiol Biomarkers Prev* 2009;18:3263-304.
- Pfyl B. The determination of nicotine in tobacco smoke II. *Z Untere Lebensm* 1933;6:501-9.
- Bradford JA, Harlan WR, Hanmer HR. Nature of cigarette smoke. *Technic of experimental smoking*. Industrial and Engineering Chemistry 1936;28:836-9.
- Adam T, Baker RR, Zimmermann R. Investigation, by single photon ionisation (SPI)-time-of-flight mass spectrometry (TOFMS), of

- the effect of different cigarette-lighting devices on the chemical composition of the first cigarette puff. *Anal Bioanal Chem* 2007; 387:575-84.
36. Peeler CL, Butters GR. Re: It's time for a change: cigarette smokers deserve meaningful information about their cigarettes. *J Natl Cancer Inst* 2000;92:842-3.
 37. Bialous SA, Yach D. Whose standard is it, anyway? How the tobacco industry determines the International Organization for Standardization (ISO) standards for tobacco and tobacco products. *Tob Control* 2001;10:96-104.
 38. Baker RR, Massey ED, Smith G. An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food Chem Toxicol* 2004; 42 Suppl:S53-83.
 39. Wynder EL, Graham EA. Tobacco smoking as a possible etiologic factor in bronchogenic carcinoma. *JAMA* 1950;143:329-36.
 40. Doll R, Hill AB. Smoking and carcinoma of the lung: preliminary report. *Br Med J* 1950;2:739-48.
 41. Federal Trade Commission. News Release. FTC to Begin Cigarette Testing. Legacy Tobacco Documents Library; 1967.
 42. Ogg CL. Determination of particulate matter and alkaloids (as nicotine) in cigarette smoke. *Journal of the Association of Official Agricultural Chemists* 1964;47:356-62.
 43. CORESTA Recommended Method N° 22. Routine Analytical Cigarette-Smoking Machine Specifications, Definitions and Standard Conditions. pp. 124-40: CORESTA Information Bulletin; 1991.
 44. International Organisation for Standardisation. Routine Analytical Cigarette Smoking Machine. Specification for the Machine and Auxiliary Equipment; ISO 3308. First Edition; 1977.
 45. Guillemin R, Radziszewski E. Analysis of smoking pattern including intake of carbon monoxide and influences of changes in cigarette design. In: Thornton RE, editor. *Smoking behaviour*. Edinburgh, London and New York: Churchill Livingstone; 1978.
 46. International Agency for Research on Cancer. IARC Monographs on the Evaluation of the Carcinogenic Risks of Chemicals to Humans: Tobacco Smoking and Involuntary Smoking. [83]; 2004, Ref Type: Serial (Book Monograph).
 47. Frederiksen LW, Frazier M. Temporal distribution of smoking. *Addict Behav* 1977;2:187-94.
 48. Frederiksen LW, Miller PM, Peterson GL. Topographical components of smoking behavior. *Addict Behav* 1977;2:55-61.
 49. Hatsukami D, Morgan SF, Pickens RW, Hughes JR. Smoking topography in a nonlaboratory environment. *Int J Addict* 1987;22: 719-25.
 50. Hatsukami DK, Pickens RW, Svikis DS, Hughes JR. Smoking topography and nicotine blood levels. *Addict Behav* 1988;13:91-5.
 51. Nemeth-Coslett R, Griffiths RR. Determinants of puff duration in cigarette smokers: I. *Pharmacol Biochem Behav* 1984;20:965-71.
 52. Battig K, Buzzi R, Nil R. Smoke yield of cigarettes and puffing behavior in men and women. *Psychopharmacology (Berl)* 1982;76:139-48.
 53. Lee E, Malson J, Waters A, Moolchan E, Pickworth W. Smoking topography: reliability and validity in dependent smokers. *Nicotine Tob Res* 2003;5:673-9.
 54. Djordjevic MV, Stellman SD, Zang E. Doses of nicotine and lung carcinogens delivered to cigarette smokers [see comments]. *J Natl Cancer Inst* 2000;19:106-11.
 55. Melikian AA, Djordjevic MV, Chen S, Richie J, Jr., Stellman SD. Effect of delivered dosage of cigarette smoke toxins on the levels of urinary biomarkers of exposure. *Cancer Epidemiol Biomarkers Prev* 2007;16: 1408-15.
 56. Puustinen P, Olkkonen H, Kolonen S, Tuomisto J. Microcomputer-aided measurement of puff parameters during smoking of low- and medium-tar cigarettes. *Scand J Clin Lab Invest* 1987;47:655-60.
 57. Puustinen P, Olkkonen H, Kolonen S, Tuomisto J. Microcomputer assisted measurement of inhalation parameters during smoking. *Arch Toxicol Suppl* 1986;9:111-4.
 58. Djordjevic MV, Fan J, Ferguson S, Hoffmann D. Self-regulation of smoking intensity. Smoke yields of the low-nicotine, low-'tar' cigarettes. *Carcinogenesis* 1995;16:2015-21.
 59. Hofer I, Nil R, Battig K. Nicotine yield as determinant of smoke exposure indicators and puffing behavior. *Pharmacol Biochem Behav* 1991; 40:139-49.
 60. Ahijevych K, Gillespie J, Demirci M, Jagadeesh J. Menthol and non-menthol cigarettes and smoke exposure in black and white women. *Pharmacol Biochem Behav* 1996;53:355-60.
 61. Zacny JP, Stitzer ML, Brown FJ, Yingling JE, Griffiths RR. Human cigarette smoking: effects of puff and inhalation parameters on smoke exposure. *J Pharmacol Exp Ther* 1987;240:554-64.
 62. Ossip-Klein DJ, Martin JE, Lomax BD, Prue DM, Davis CJ. Assessment of smoking topography generalization across laboratory, clinical, and naturalistic settings. *Addict Behav* 1983;8:11-7.
 63. Sutton SR, Russell MA, Iyer R, Feyerabend C, Saloojee Y. Relationship between cigarette yields, puffing patterns, and smoke intake: evidence for tar compensation? *Br Med J (Clin Res Ed)* 1982;285:600-3.
 64. Adams PI. Measurements on Puffs Taken by Human Smokers [Abstract]. 1966.
 65. Adams PI. Changes in Personal Smoking Habits brought about by changes in cigarette smoke yields. pp. 102-8 Tokyo: Japan Tobacco and Salt Public Corporation; 1976.
 66. Creighton DE, Noble MJ, Whewell RT. Instruments to measure, record, and duplicate human smoking patterns. In: Thornton RE, editor. *Smoking Behaviour - Physiological and Psychological Influences*. Edinburgh, London and New York: Churchill Livingstone; 1978, p. 277-88.
 67. O'Connor RJ, Ashare RL, Cummings KM, Hawk LW, Jr. Comparing smoking behaviors and exposures from flavored and unflavored cigarettes. *Addict Behav* 2007;32:869-74.
 68. Pickens R, Gust S, Catchings P, Svikis D. Measurement of some topographical aspects of smoking in the natural environment. In: Grabowski J, Bell C, editors. *Monograph 48: Measurement in the Analysis and Treatment of Smoking Behavior*. Washington, DC: National Institute on Drug Abuse Research; 1983, p. 62-73, Ref Type: Serial (Book Monograph).
 69. Henningfield JE, Yingling J, Griffiths RR, Pickens R. An inexpensive portable device for measuring puffing behavior by cigarette smokers. *Pharmacol Biochem Behav* 1980;12:811-3.
 70. Melikian AA, Djordjevic MV, Hosey J, et al. Gender differences relative to smoking behavior and emissions of toxins from mainstream cigarette smoke. *Nicotine Tob Res* 2007;9:377-87.
 71. Eissenberg T, Adams C, Riggins EC, Likness M, III. Smokers' sex and the effects of tobacco cigarettes: subject-rated and physiological measures. *Nicotine Tob Res* 1999;1:317-24.
 72. Buchhalter AR, Eissenberg T. Preliminary evaluation of a novel smoking system: effects on subjective and physiological measures and on smoking behavior. *Nicotine Tob Res* 2000;2:39-43.
 73. Buchhalter AR, Schriener L, Eissenberg T. Withdrawal-suppressing effects of a novel smoking system: comparison with own brand, not own brand, and de-nicotinized cigarettes. *Nicotine Tob Res* 2001;3: 111-8.
 74. Breland AB, Buchhalter AR, Evans SE, Eissenberg T. Evaluating acute effects of potential reduced-exposure products for smokers: clinical laboratory methodology. *Nicotine Tob Res* 2002;4 Suppl 2:S131-40.
 75. Hughes JR, Hecht SS, Carmella SG, Murphy SE, Callas P. Smoking behaviour and toxin exposure during six weeks use of a potential reduced exposure product: Omni. *Tob Control* 2004;13:175-9.
 76. Tidey JW, Rohsenow DJ, Kaplan GB, Swift RM. Cigarette smoking topography in smokers with schizophrenia and matched non-psychiatric controls. *Drug Alcohol Depend* 2005;80:259-65.
 77. Franken FH, Pickworth WB, Epstein DH, Moolchan ET. Smoking rates and topography predict adolescent smoking cessation following treatment with nicotine replacement therapy. *Cancer Epidemiol Biomarkers Prev* 2006;15:154-7.
 78. Strasser AA, Malaiyandi V, Hoffmann E, Tyndale RF, Lerman C. An association of CYP2A6 genotype and smoking topography. *Nicotine Tob Res* 2007;9:511-8.
 79. Rees VW, Wayne GF, Connolly GN. Puffing style and human exposure minimally altered by switching to a carbon-filtered cigarette. *Cancer Epidemiol Biomarkers Prev* 2008;17:2995-3003.
 80. Grainge M, Shahab L, Hammond D, O'Connor R, McNeill A. First cigarette on waking and time of day as predictors of puffing behaviour in UK adult smokers. *Drug Alcohol Depend* 2009;PM:19264427.
 81. Djordjevic MV, Hoffmann D, Hoffmann I. Nicotine regulates smoking patterns. *Prev Med* 1997;26:435-40.
 82. Hammond D, Fong GT, Cummings KM, Hyland A. Smoking topography, brand switching, and nicotine delivery: results from an *in vivo* study. *Cancer Epidemiol Biomarkers Prev* 2005;14:1370-5.
 83. Blank MD, Disharoon S, Eissenberg T. Comparison of methods for measurement of smoking behavior: mouthpiece-based computerized devices versus direct observation. *Nicotine Tob Res* 2009;11:896-903.
 84. Evans SE, Buchhalter AR, Breland A, Kleykamp BA, Eissenberg T. Ambulatory puff topography measurement: A validation study. New Orleans, LA; 2003.
 85. Gust SW, Pickens RW, Pechacek TF. Relation of puff volume to other topographical measures of smoking. *Addict Behav* 1983;8:115-9.
 86. Guyatt AR, Kirkham AJ, Baldry AG, Dixon M, Cumming G. How does puffing behavior alter during the smoking of a single cigarette? *Pharmacol Biochem Behav* 1989;33:189-95.
 87. Ahijevych K, Gillespie J. Nicotine dependence and smoking topography among black and white women. *Res Nurs Health* 1997;20:505-14.
 88. National Cancer Institute Expert Committee. The FTC cigarette test method for determining, tar, nicotine and carbon monoxide yields of U.S. cigarettes. [Monograph No. 7]. U.S. Department of Health and

- Human Services National Cancer Institute; 1996, Ref Type: Serial (Book Monograph).
89. Hammond D, Collishaw NE, Callard C. Secret science: tobacco industry research on smoking behaviour and cigarette toxicity. *Lancet* 2006;367:781-7.
 90. Kozlowski LT, O'Connor RJ. Cigarette filter ventilation is a defective design because of misleading taste, bigger puffs, and blocked vents. *Tob Control* 2002;11 Suppl 1:140-50.
 91. Kozlowski LT, O'Connor RJ. Official cigarette tar tests are misleading: use a two-stage, compensating test. *Lancet* 2000;355:2159-61.
 92. Assaf AR, Parker D, Lapane KL, Carleton McKenney JL. Are there gender differences in self-reported smoking practices? Correlation with thiocyanate and cotinine levels in smokers and nonsmokers from the Pawtucket Heart Health Program. *J Womens Health (Larchmt)* 2002;11:899-906.
 93. Thun MJ, Heath CW, Jr. Changes in mortality from smoking in two American Cancer Society prospective studies since 1959. *Prev Med* 1997;26:422-6.
 94. Kandel DB, Chen K. Extent of smoking and nicotine dependence in the United States: 1991-1993. *Nicotine Tob Res* 2000;2:263-74.
 95. Etter JF, Vu DT, Perneger TV. Saliva cotinine levels in smokers and nonsmokers. *Am J Epidemiol* 2000;151:251-8.
 96. Mustonen TK, Spencer SM, Hoskinson RA, Sachs DP, Garvey AJ. The influence of gender, race, and menthol content on tobacco exposure measures. *Nicotine Tob Res* 2005;7:581-90.
 97. Epstein LH, Dickson BE, Ossip DJ, et al. Relationships among measures of smoking topography. *Addict Behav* 1982;7:307-10.
 98. Wood T, Wewers ME, Groner J, Ahijevych K. Smoke constituent exposure and smoking topography of adolescent daily cigarette smokers. *Nicotine Tob Res* 2004;6:853-62.
 99. Ahijevych K, Parsley LA. Smoke constituent exposure and stage of change in black and white women cigarette smokers. *Addict Behav* 1999;24:115-20.
 100. Lombardo T, Carreno L. Relationship of type A behavior pattern in smokers to carbon monoxide exposure and smoking topography. *Health Psychol* 1987;6:445-52.
 101. Hatsukami DK, Morgan SF, Pickens RW, Champagne SE. Situational factors in cigarette smoking. *Addict Behav* 1990;15:1-12.
 102. Morgan SF, Gust SW, Pickens RW, Champagne SE, Hughes JR. Temporal patterns of smoking topography in the natural environment. *Int J Addict* 1985;20:613-21.
 103. Herning RI, Jones RT, Benowitz NL, Mines AH. How a cigarette is smoked determines blood nicotine levels. *Clin Pharmacol Ther* 1983;33:84-90.
 104. Bridges RB, Combs JG, Humble JW, et al. Puffing topography as a determinant of smoke exposure. *Pharmacol Biochem Behav* 1990;37:29-39.
 105. Kolonen S, Tuomisto J, Puustinen P, Airaksinen MM. Puffing behavior during the smoking of a single cigarette in a naturalistic environment. *Pharmacol Biochem Behav* 1992;41:701-6.
 106. Lichtenstein E, Antonuccio DO. Dimensions of smoking behavior. *Addict Behav* 1981;6:365-7.
 107. Burling TA, Stitzer ML, Bigelow GE, Mead AM. Smoking topography and carbon monoxide levels in smokers. *Addict Behav* 1985;10:319-23.
 108. Humble JW, Turbek JA, Rehm SR. Smoking history, cigarette yield and smoking behavior as determinants of smoke exposure. *Eur J Respir Dis Suppl* 1986;146:129-37.
 109. Djordjevic MV, Stellman SD, Zang E. Doses of nicotine and lung carcinogens delivered to cigarette smokers. *J Natl Cancer Inst* 2000;92:106-11.
 110. Scherer G. Smoking behaviour and compensation: a review of the literature. *Psychopharmacology (Berl)* 1999;145:1-20.
 111. Buzzi R, Nil R, Battig K. Development of puffing behavior along burning time of a cigarette-no relation to alveolar inhalation or nicotine delivery of the cigarettes? *Psychopharmacology (Berl)* 1985;86:102-7.
 112. Rieben FW. Smoking behaviour and increase in nicotine and carboxyhaemoglobin in venous blood. *Clin Invest* 1992;70:335-42.
 113. Zaczny JP, Stitzer ML, Yingling JE. Cigarette filter vent blocking: effects on smoking topography and carbon monoxide exposure. *Pharmacol Biochem Behav* 1986;25:1245-52.
 114. Weinhold LL, Stitzer ML. Effects of puff number and puff spacing on carbon monoxide exposure from commercial brand cigarettes. *Pharmacol Biochem Behav* 1989;33:853-8.
 115. Strasser AA, Lerman C, Sanborn PM, Pickworth WB, Feldman EA. New lower nicotine cigarettes can produce compensatory smoking and increased carbon monoxide exposure. *Drug Alcohol Depend* 2007;86:294-300.
 116. Sweeney CT, Kozlowski LT. Blocking filter vents increases carbon monoxide levels from ultralight, but not light cigarettes. *Pharmacol Biochem Behav* 1998;59:767-73.
 117. Strasser AA, Ashare RL, Kozlowski LT, Pickworth WB. The effect of filter vent blocking and smoking topography on carbon monoxide levels in smokers. *Pharmacol Biochem Behav* 2005;82:320-9.
 118. Kolonen S, Tuomisto J, Puustinen P, Airaksinen MM. Smoking behavior in low-yield cigarette smokers and switchers in the natural environment. *Pharmacol Biochem Behav* 1991;40:177-80.
 119. Stepney R. Would a medium-nicotine, low-tar cigarette be less hazardous to health? *Br Med J (Clin Res Ed)* 1981;283:1292-6.
 120. Russell MA, Sutton SR, Iyer R, Fejerabend C, Vesey CJ. Long-term switching to low-tar low-nicotine cigarettes. *Br J Addict* 1982;77:145-58.
 121. Ossip-Klein DJ, Epstein LH, Winter MK, et al. Does switching to low tar/nicotine/carbon monoxide-yield cigarettes decrease alveolar carbon monoxide measures? A randomized controlled trial. *J Consult Clin Psychol* 1983;51:234-41.
 122. Nil R, Buzzi R, Battig K. Effects of different cigarette smoke yields on puffing and inhalation: is the measurement of inhalation volumes relevant for smoke absorption? *Pharmacol Biochem Behav* 1986;24:587-95.
 123. Armitage AK, Alexander J, Hopkins R, Ward C. Evaluation of a low to middle tar/medium nicotine cigarette designed to maintain nicotine delivery to the smoker. *Psychopharmacology (Berl)* 1988;96:447-53.
 124. Zaczny JP, Stitzer ML. Cigarette brand-switching: effects on smoke exposure and smoking behavior. *J Pharmacol Exp Ther* 1988;246:619-27.
 125. Nil R, Battig K. Separate effects of cigarette smoke yield and smoke taste on smoking behavior. *Psychopharmacology (Berl)* 1989;99:54-9.
 126. Baldinger B, Hasenfratz M, Battig K. Switching to ultralight nicotine cigarettes: effects of different tar yields and blocking of olfactory cues. *Pharmacol Biochem Behav* 1995;50:233-9.
 127. Herning RI, Jones RT, Bachman J, Mines AH. Puff volume increases when low-nicotine cigarettes are smoked. *Br Med J (Clin Res Ed)* 1981;283:187-9.
 128. Tobin MJ, Sackner MA. Monitoring smoking patterns of low and high tar cigarettes with inductive plethysmography. *Am Rev Respir Dis* 1982;126:258-64.
 129. McBride MJ, Guyatt AR, Kirkham AJ, Cumming G. Assessment of smoking behaviour and ventilation with cigarettes of differing nicotine yields. *Clin Sci (Lond)* 1984;67:619-31.
 130. Gust SW, Pickens RW. Does cigarette nicotine yield affect puff volume? *Clin Pharmacol Ther* 1982;32:418-22.
 131. Woodman G, Newman SP, Pavia D, Clarke SW. The separate effects of tar and nicotine on the cigarette smoking manoeuvre. *Eur J Respir Dis* 1987;70:316-21.
 132. Woodman G, Newman SP, Pavia D, Clarke SW. Inhaled smoke volume and puff indices with cigarettes of different tar and nicotine levels. *Eur J Respir Dis* 1987;70:187-92.
 133. Benowitz NL, Jacob P, III, Bernert JT, et al. Carcinogen exposure during short-term switching from regular to "light" cigarettes. *Cancer Epidemiol Biomarkers Prev* 2005;14:1376-83.
 134. Sweeney CT, Kozlowski LT, Parsa P. Effect of filter vent blocking on carbon monoxide exposure from selected lower tar cigarette brands. *Pharmacol Biochem Behav* 1999;63:167-73.
 135. Wayne GF, Thomas BF, Connolly GN. Physical design analysis and mainstream smoke constituent yields of the new potential reduced exposure product, Marlboro UltraSmooth. *Nicotine Tob Res* 2007;9:1197-206.
 136. Breland AB, Acosta MC, Eissenberg T. Tobacco specific nitrosamines and potential reduced exposure products for smokers: a preliminary evaluation of Advance. *Tob Control* 2003;12:317-21.
 137. Breland AB, Evans SE, Buchhalter AR, Eissenberg T. Acute effects of Advance: a potential reduced exposure product for smokers. *Tob Control* 2002;11:376-8.
 138. Breland AB, Kleykamp BA, Eissenberg T. Clinical laboratory evaluation of potential reduced exposure products for smokers. *Nicotine Tob Res* 2006;8:727-38.
 139. Slade J, Connolly GN, Lymperis D. Eclipse: does it live up to its health claims? *Tob Control* 2002;11 Suppl 2:ii64-70.
 140. Lee EM, Malson JL, Moolchan ET, Pickworth WB. Quantitative comparisons between a nicotine delivery device (Eclipse) and conventional cigarette smoking. *Nicotine Tob Res* 2004;6:95-102.
 141. Hatsukami DK, Benowitz NL, Rennard SI, Oncken C, Hecht SS. Biomarkers to assess the utility of potential reduced exposure tobacco products. *Nicotine Tob Res* 2006;8:600-22.
 142. Tobin MJ, Jenouri G, Sackner MA. Subjective and objective measurement of cigarette smoke inhalation. *Chest* 1982;82:696-700.
 143. Sackner J, Nixon A, Davis B, Atkins N, Sackner M. Non-invasive measurement of ventilation during exercise using a respiratory inductive plethysmograph. *I. Am Rev Respir Dis* 1980;122:867-71.

144. Chadha TS, Watson H, Birch S, et al. Validation of respiratory inductive plethysmography using different calibration procedures. *Am Rev Respir Dis* 1982;125:644-9.
145. Charles FK, Krautter GR, Mariner DC. Post-puff respiration measures on smokers of different tar yield cigarettes. *Inhal Toxicol* 2009;1-7.
146. Armitage AK, Dixon M, Frost DE, Mariner DC, Sinclair NM. The effect of inhalation volume and breath-hold duration on the retention of nicotine and solanesol in the human respiratory tract and on subsequent plasma nicotine concentrations during cigarette smoking. *Contributions to Tobacco Research* 2004;21:240-9.
147. Feng S, Plunkett SE, Lam K, et al. A new method for estimating the retention of selected smoke constituents in the respiratory tract of smokers during cigarette smoking. *Inhal Toxicol* 2007;19:169-79.
148. Rickert WS, Robinson JC, Collishaw NE, Bray DF. Estimating the hazards of "less hazardous" cigarettes. III. A study of the effect of various smoking conditions on yields of hydrogen cyanide and cigarette tar. *J Toxicol Environ Health* 1983;12:39-54.
149. International Organization for Standardization—Ad Hoc Smoking Behavior Review Team Working Group 9 of the ISO TC 126. A review of human smoking behavior data and recommendations for a new ISO standard for the machine smoking of cigarettes. 2005.
150. Kozlowski LT. Application of some physical indicators of cigarette smoking. *Addict Behav* 1981;6:213-9.
151. Kozlowski LTOR, Sweeney C. Cigarette design. *Smoking and Tobacco Control Monograph No. 13: Risks Associated with Smoking Cigarettes with Low Machine-Measured Yields of Tar and Nicotine*. pp. 13-37. Bethesda, MD: National Cancer Institute; 2001, Ref Type: Serial (Book Monograph).
152. Harris JE. Smoke yields of tobacco-specific nitrosamines in relation to FTC tar level and cigarette manufacturer: analysis of the Massachusetts Benchmark Study. *Public Health Rep* 2001;116:336-43.
153. Rickert WS. Determination of Cigarette Yields under Realistic Conditions. 1996, Ref Type: Generic.
154. Schorp MK. Summary of Literature Data on Smoking Topography. In: A review of human smoking behavior data and recommendations for a new ISO standard for the machine smoking of cigarettes. 2005.
155. Hammond D, O'Connor RJ. Constituents in tobacco and smoke emissions from Canadian cigarettes. *Tob Control* 2008;17 Suppl 1:i24-31.
156. Rickert WS, Trivedi AH, Momin RA, Wright WG, Lauterbach JH. Effect of smoking conditions and methods of collection on the mutagenicity and cytotoxicity of cigarette mainstream smoke. *Toxicol Sci* 2007;96:285-93.
157. Counts ME, Morton MJ, Laffoon SW, Cox RH, Lipowicz PJ. Smoke composition and predicting relationships for international commercial cigarettes smoked with three machine-smoking conditions. *Regul Toxicol Pharmacol* 2005;41:185-227.
158. Roemer E, Stabbert R, Rustemeier K, et al. Chemical composition, cytotoxicity and mutagenicity of smoke from US commercial and reference cigarettes smoked under two sets of machine smoking conditions. *Toxicology* 2004;195:31-52.
159. Harris JE. Incomplete compensation does not imply reduced harm: yields of 40 smoke toxicants per milligram nicotine in regular filter versus low-tar cigarettes in the 1999 Massachusetts Benchmark Study. *Nicotine Tob Res* 2004;6:797-807.
160. Urban HJ, Gomm W, Schorp M. A modelling approach to develop machine smoking protocols reflecting human puffing behaviour for conventional cigarettes. *Contributions to Tobacco Research* 2008;23: 8-18.
161. Strasser AA, Pickworth WB, Patterson F, Lerman C. Smoking topography predicts abstinence following treatment with nicotine replacement therapy. *Cancer Epidemiol Biomarkers Prev* 2004;13: 1800-4.
162. Lerman C, Wileyto EP, Patterson F, et al. The functional mu opioid receptor (OPRM1) Asn40Asp variant predicts short-term response to nicotine replacement therapy in a clinical trial. *Pharmacogenomics J* 2004;4:184-92.
163. Lerman C, Shields PG, Wileyto EP, et al. Pharmacogenetic investigation of smoking cessation treatment. *Pharmacogen* 2002;12:627-34.
164. Lerman C, Berrettini W. Elucidating the role of genetic factors in smoking behavior and nicotine dependence. *Am J Med Genet* 2003; 118B:48-54.
165. Lerman C, Caporaso NE, Main D, et al. Evidence suggesting the role of specific genetic factors in cigarette smoking. *Health Psychol* 1999; 18:14-20.
166. Ring HZ, Valdes AM, Nishita DM, et al. Gene-gene interactions between CYP2B6 and CYP2A6 in nicotine metabolism. *Pharmacogenet Genomics* 2007;17:1007-15.
167. Mwenifumbo JC, Tyndale RF. Genetic variability in CYP2A6 and the pharmacokinetics of nicotine. *Pharmacogenomics* 2007;8:1385-402.
168. Benowitz NL, Swan GE, Jacob P, III, Lessov-Schlaggar CN, Tyndale RF. CYP2A6 genotype and the metabolism and disposition kinetics of nicotine. *Clin Pharmacol Ther* 2006;80:457-67.
169. Malaiyandi V, Goodz SD, Sellers EM, Tyndale RF. CYP2A6 genotype, phenotype, and the use of nicotine metabolites as biomarkers during ad libitum smoking. *Cancer Epidemiol Biomarkers Prev* 2006;15: 1812-9.
170. Guillerme R, Radziszewski E. A new method of analyzing the act of smoking. *Ann Tabac Etudes L'Equiment* 1975;13:101-10.
171. Rawbone RC, Murphy K, Tate ME. The analysis of smoking patterns. In: Thornton RE, editor. *Smoking Behaviour*. London: Churchill Livingstone; 1978, p. 171-94.
172. Sackner MA. Monitoring of ventilation without a physical connection to the airway. In: Sackner MA, editor. *Diagnostic techniques in pulmonary disease*. Pat 1. New-York: Marcel Dekker; 1980, p. 503-37.

Reconciling Human Smoking Behavior and Machine Smoking Patterns: Implications for Understanding Smoking Behavior and the Impact on Laboratory Studies

Catalin Marian, Richard J. O'Connor, Mirjana V. Djordjevic, et al.

Cancer Epidemiol Biomarkers Prev 2009;18:3305-3320.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/18/12/3305>

Cited articles This article cites 143 articles, 24 of which you can access for free at:
<http://cebp.aacrjournals.org/content/18/12/3305.full#ref-list-1>

Citing articles This article has been cited by 10 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/18/12/3305.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/18/12/3305>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.