

TP53 R249S Mutations, Exposure to Aflatoxin, and Occurrence of Hepatocellular Carcinoma in a Cohort of Chronic Hepatitis B Virus Carriers from Qidong, China

Katarzyna Szymańska,¹ Jian-Guo Chen,² Yan Cui,³ Yun Yun Gong,⁴ Paul Craig Turner,⁴ Stéphanie Villar,¹ Christopher Paul Wild,⁴ Donald Maxwell Parkin,⁵ and Pierre Hainaut¹

¹IARC, Lyon, France; ²Qidong Liver Cancer Institute and Qidong Cancer Registry, Department of Epidemiology, QDLICI, Jiangsu, P.R. China; ³Office of Health Assessment and Epidemiology, Los Angeles County Department of Public Health, Los Angeles, California; ⁴Molecular Epidemiology Unit, Centre for Epidemiology and Biostatistics, LIGHT Laboratories, University of Leeds, Leeds, United Kingdom; and ⁵Clinical Trial Service Unit and Epidemiological Studies Unit, University of Oxford, Oxford, United Kingdom

Abstract

Hepatocellular carcinoma (HCC) has a high mortality in East Asia and Sub-Saharan Africa, two regions where the main etiologic factors are chronic infections with hepatitis B virus and dietary exposure to aflatoxin. A single base substitution at the third nucleotide of codon 249 of *TP53* (*R249S*) is common in HCC in these regions and has been associated with aflatoxin-DNA adducts. To determine whether *R249S* may be detected in plasma DNA before HCC diagnosis, we conducted a case-control study nested in a cohort of adult chronic hepatitis B virus carriers from Qidong County, People's Republic of China. Of the 234 plasma specimens that yielded adequate DNA, only 2 (0.9%) were positive for *R249S* by restriction fragment length polymorphisms, and both of them were controls. Of the 249 subjects tested for aflatoxin-albumin adducts, 168 (67%) were positive, with

equal distribution between cases and controls. Aflatoxin-albumin adduct levels were low in the study, suggesting an overall low ongoing exposure to aflatoxin in this cohort. The *R249S* mutation was detected in 11 of 18 (61%) available tumor tissues. To assess whether low levels of mutant DNA were detectable in pre-diagnosis plasma, 14 plasma specimens from these patients were analyzed by short oligonucleotide mass analysis. Nine of them (64%) were found to be positive. Overall, these results suggest that HCC containing *R249S* can occur in the absence of significant recent exposure to aflatoxins. The use of short oligonucleotide mass analysis in the context of low ongoing aflatoxin exposure may allow the detection of *R249S* in plasma several months ahead of clinical diagnosis. (Cancer Epidemiol Biomarkers Prev 2009;18(5):1638–43)

Introduction

Hepatocellular carcinoma (HCC) is a major public health problem in many parts of the world, with high incidence areas in East Asia and Sub-Saharan Africa, where the main etiologic factors are chronic infection with hepatitis viruses [mostly hepatitis B virus (HBV)] and dietary exposure to aflatoxins, including aflatoxin B₁. Almost 54% of all liver cancer cases occur in the People's Republic of China (PRC; ref. 1). In Qidong County, PRC, about 16% of the adult population are seropositive for HBsAg (2). A study on 181 consecutive HCC cases from Qidong showed markers of HBV infection in all of them, whereas only 6 of 119 HCC cases were coinfecting with hepatitis C virus (HCV). Dietary exposure to aflatoxins was ubiquitous (3). A missense mutation at codon 249 in *TP53*, *AGG* to *AGT*, leading to a substitution of an arginine for a serine (*R249S*), is extremely common in HCC in areas with high prevalence of HBV chronic carriage and

aflatoxin exposure, but not in areas with high HBV prevalence alone (4, 5). This mutation is considered to be a consequence of aflatoxin N⁷-guanine adducts formed at the third base of codon 249 of the *TP53* gene (6). However, the reasons why this particular mutation is selected in HCC are not known. There is evidence of a synergistic effect between HBV carriage and aflatoxin exposure in inducing this mutation in HCC (4). However, despite ecological correlations (7), there is limited evidence for an association between recent aflatoxin exposure and the mutation at the individual level (8).

Recent studies have shown that *R249S* is detectable in free DNA extracted from serum or plasma of HCC patients. Using a sensitive method based on mass spectrometry, Jackson et al. (9) detected *R249S* at least 1 year before diagnosis in the sera of four of eight Qidong HCC patients who were positive for this mutation at the time of diagnosis. In a case-control study in The Gambia, another area of high incidence of HCC and common exposure to HBV and aflatoxin B₁, Kirk et al. (10) found that *R249S* was sometimes detectable in subjects with liver cirrhosis (15.3%) or in control subjects (3.5%), although with a lower prevalence than in HCC patients (39.8%). In our study, we have assessed the prevalence of *R249S* mutations in plasma and tumor DNA, HCV infection, and exposure to aflatoxin [based on quantitation of aflatoxin-albumin (Af-alb) adducts in the serum]

Received 11/18/08; revised 1/29/09; accepted 3/4/09; published OnlineFirst 4/14/09.

Grant support: French National Cancer Institute (INCa; CircBioCancer program). C.P. Wild was supported by National Institute of Environmental Health Sciences grant ES06052.

Note: Current address of C.P. Wild: IARC, Lyon, France.

Requests for reprints: Pierre Hainaut, Group of Molecular Carcinogenesis and Biomarkers, IARC, 150 Cours Albert Thomas, 69372 Lyon Cedex 08, France. Phone: 33-4-7273-8532; Fax: 33-4-7273-8322. E-mail: hainaut@iarc.fr

Copyright © 2009 American Association for Cancer Research.

doi:10.1158/1055-9965.EPI-08-1102

in a case-control study nested within a cohort of HBV chronic carriers in Qidong, PRC, undergoing follow-up for the occurrence of HCC and other cancers.

Materials and Methods

Study Design and Subjects. In 1989, a program was launched to screen for liver cancer among men aged 30 to 59 resident in Qidong County, Jiangsu Province, PRC. Within this program, 36,000 subjects were tested for carriage of HBsAg, and 3,712 subjects were followed up by testing for increased serum α -fetoprotein at 6-month intervals for up to 6 years, with a record of the occurrence of liver cancers (11). Plasma specimens (~0.5-1.0 ml) taken at the time of the initial recruitment and at subsequent screenings were stored at -20°C . A total of 130 primary liver cancer patients were identified between 1993 and 1998. These cases were identified among the 3,712 individuals either via the screening process or by follow-up of the cohort through the Qidong Cancer Registry (11). Most patients were diagnosed by a combination of α -fetoprotein measurement, alanine aminotransferase tests, and ultrasound. For 20 subjects, tumor biopsies (12 cases) or surgical specimens (8 cases) were available, the latter with both tumoral and peritumoral tissue specimens. For each case, one age-matched control was randomly selected from within the same study cohort. To be eligible as a control, a subject had to be in the same age category as the case, be still alive and free of cancer at the date of cancer diagnosis for the corresponding case, and have the same number of collected specimens as the case in the period before the case was diagnosed. Considering the rarity of HCV infection in the Chinese population, two more controls for each case were selected to explore the possible synergism of HCV and HBV in the development of HCC. The presence of HCV antibody was detected using a second-generation ELISA assay.

TP53 Analysis. The tissue specimens were histologically evaluated, classified according to the Edmonson-Steiner criteria (ES grades), and immunostained for p53 using a polyclonal antibody CM1 (NovoCastra). Immunostaining, DNA extraction from tissue and plasma specimens, and TP53 analyses by restriction fragment length polymorphism (RFLP) and sequencing were done as described elsewhere (12). Briefly, exon 7 of the TP53 gene was amplified and the PCR products were digested with a restriction endonuclease, *Hae*III, recognizing sequence CCGG, encompassing the 249 codon. Digestion of the wild-type exon 7 generates two bands (92 and 66 bp), whereas mutant material, in which the restriction site has been destroyed by the mutation, yields one band of 158 bp. These mutant fragments were cut out of the gel, reamplified, and sequenced by automated dideoxy sequencing to confirm the presence of the mutation. The samples were classified as mutant if the same result was found for two different PCR products. Mutations in plasma specimens were confirmed by short oligonucleotide mass analysis (SOMA) as described elsewhere (9, 13). Tissue specimens appearing wild type at digestion were directly sequenced (from a third PCR product) to screen for possible other mutations in exon 7. To search for mutations in other exons of the TP53 gene,

additional analyses were done using temporal temperature gradient electrophoresis as described elsewhere (14, 15). Additionally, 15 plasma specimens corresponding to the analyzed tumors were analyzed for the presence of the R249S DNA by SOMA (9, 13).

Aflatoxin-albumin Adducts. The concentration of Aflatoxin-albumin adducts in plasma was measured as previously described (16). Albumin (2 mg) was digested overnight using Pronase, and aflatoxin-containing residues were isolated by solid-phase extraction using a Sep-Pak cartridge. Samples were analyzed in quadruplicate by ELISA on two occasions on separate days using aflatoxin B₁-lysine in the standard curve. The limit of detection was 3 pg aflatoxin B₁-lysine equivalent/mg of albumin.

Statistical Analysis. Logistic regression models (including exact age) were used to estimate the odds ratios and 95% confidence intervals associated with exposure to HCV and Af-alb adducts and the development of liver cancer.

Results

Detection of R249S DNA in Tumor Tissues. To determine the prevalence of R249S TP53 mutation in HCC cases appearing among chronic HBV carriers, 20 available tumor specimens were analyzed. Ten tumors (50%: 6 of 12 biopsies and 4 of 8 surgical specimens) stained positive for p53 (at least 10% of stained tumor cells) with antibody CM1. Only two tumors (10%) showed positivity in >50% of tumor cells and both were diagnosed as HCC ES 2 (Table 1). DNA of sufficient quality for PCR amplification was obtained from 18 specimens. Eleven of these 18 tumors (61%) contained G-to-T transversions at the third nucleotide of codon 249 (the R249S mutation; Table 1). The distribution of mutations was uneven between the two series of specimens: Whereas only 5 of 11 biopsies (ID nos. 1-12 in Table 1) contained the R249S DNA, the mutation was found in 6 of 7 surgical specimens (the only one without the mutation contained a majority of nontumoral cells, so that the wild-type allele may have masked the mutant allele). Peritumoral liver tissues were available for eight cases, including five cirrhotic and three fibrotic tissues. R249S mutations were detected in three of the five cirrhotic tissues but not in fibrotic tissues. In addition to R249S, two other mutations were found by sequencing of exon 7, both in R249S-negative tumors (CCC→TCC, Pro→Ser, at codon 250 in patient no. 1, and GTC→TTC at donor splice site in patient no. 11; Table 1). The presence of TP53 mutations did not correlate with histologic features of the tumors or with p53 immunohistochemistry.

Detection of R249S in Plasma DNA. Cell-free DNA was extracted from plasma collected during the follow-up of the cohort of chronic carriers. A total of 130 subjects who developed HCC during the follow-up ("cases") were matched with subjects who did not develop HCC ("controls"). For both cases and controls, the most recently collected plasma samples were analyzed for the presence of R249S DNA by RFLP. In cases, the time elapsed between plasma collection and cancer diagnosis ranged from 0 to 74 months (two plasma samples were obtained at the date of cancer diagnosis). Only three specimens were found to be positive by RFLP, and they

Table 1. Histologically confirmed HCCs arising within the Qidong cohort of hepatitis B chronic carriers: patients' characteristics, tumor grade, HCV, and TP53 status

ID	Sex	Age (y)	Diagnosis	Anti-HCV	p53 immunostaining (%)	TP53 mutations, exon 7
6	M	33	HCC, clear cell	—	—	AGG→AGT at 249
3	M	47	HCC, ES 2	ND	>50+	AGG→AGT at 249
8	M	36	HCC, ES 2	ND	—	Wt
9	M	38	HCC, ES 2	—	>50+	Wt
10	M	47	HCC, ES 2	ND	0-10+	AGG→AGT at 249
4	M	36	HCC, ES 2	ND	20-50+	NA
12	M	42	HCC, ES 2	—	—	AGG→AGT at 249
15CA	M	49	HCC, ES 2	—	0-10+	AGG→AGT at 249
15AF			Cirrhosis	—	—	Wt
16CA	M	61	HCC, ES 2	ND	—	AGG→AGT at 249
16AF			Fibrosis	—	—	Wt
13CA	M	39	HCC, ES 2	ND	0-10+	AGG→AGT at 249
13AF			Cirrhosis, inactive	—	—	AGG→AGT at 249
14CA	M	35	HCC, ES 2	—	20-50+	AGG→AGT at 249
14AF			Cirrhosis, moderately active	—	—	AGG→AGT at 249
19CA	M	59	HCC, ES 2	—	20-50+	AGG→AGT at 249
19AF			Cirrhosis, vascular neoplastic emboli	—	20-50+	AGG→AGT at 249
18CA	M	59	HCC ES 2	—	—	NA
18AF			Cirrhosis, active	—	—	Wt
1	M	41	HCC, ES 3	—	20-50+	CCC→TCC at 250
5	M	50	HCC, ES 3	—	—	Wt
7	M	38	HCC, ES 3	—	—	Wt
11	M	50	HCC, ES 3	—	—	GTC→TTC at donor splice site
17CA	M	46	HCC, ES 3	—	20-50+	AGG→AGT at 249
17AF			Fibrosis	—	—	Wt
2	M	37	HCC	—	20-50+	AGG→AGT at 249
20CA	M	62	HCC (vascular embolus, small tumor area)	—	20-50+	Wt
20AF			Fibrosis	—	—	Wt

NOTE: Samples 1 to 12 are biopsies and samples 13 to 20 are surgical specimens. In the series of surgical specimens, peritumoral liver specimens were also available (marked AF in the ID number as opposed to CA for tumor specimens).

were reanalyzed by SOMA to confirm the presence of the mutation, with two (0.9%) being confirmed as R249S-positive by RFLP/sequencing and SOMA. Both were controls, neither of whom developed cancer during the follow-up period, which was 6 years for one subject but only 1 year for the other. Plasma specimens were available for 14 of the 18 patients from whom tumors were analyzed for the presence of R249S. Only two of these matched plasma specimens were collected at the time of cancer diagnosis. Twelve of them were analyzed by RFLP and none of them was found positive for R249S DNA. Next, all 14 plasma specimens were analyzed by SOMA. Nine of 14 samples (64%) were found to contain R249S DNA at levels between 157 and 3,546 gene copies/ml of plasma. This result shows that low levels of R249S were detectable in the plasma ahead of cancer diagnosis. Due to a limited amount of plasma and logistical constraints, analysis by SOMA could not be extended to all specimens.

The concordance between positivity in the plasma as detected by SOMA and positivity in the tumor as detected by RFLP and sequencing was poor. Three of the SOMA-positive plasma specimens were from subjects who later developed R249S-negative tumors. In contrast, two of the plasma specimens that were negative by SOMA corresponded to tumors that were found to be positive for R249S by RFLP. Thus, the presence of R249S in the plasma was not systematically predictive of the detection of the same mutation in the tumor (Table 2).

Detection of HCV. To assess the possible contribution of HCV as a risk of HCC in hepatitis B chronic carriers, we analyzed the prevalence of antibodies against HCV in 127 cases and 380 controls. There were only 10 positive subjects among 507 tested (2%; 2 cases and 8 controls; Table 3A). The odds ratio associated with HCV infection was 0.94 (95% confidence interval: 0.45-1.97). Thus, HCV does not seem to substantially contribute to the risk of HCC in this cohort.

Detection of Aflatoxin Adducts. To determine whether exposure to aflatoxin in the months and years that precede diagnosis was an important risk factor for HCC, the plasma from the last available aliquot was also analyzed for Af-alb adducts. Testing was possible in 123 cases and 126 controls. Of the 249 individuals tested, 168 were positive for this biomarker of exposure to aflatoxin in one or more specimens taken before the diagnosis of the case or the same reference date for the controls (Table 3B). The adduct levels were uniformly low, with most of the positive specimens showing levels between 5 and 10 pg/mg and only six specimens with levels higher than 10 pg/mg. Using a value of 3 pg/mg to dichotomize individuals into exposed and unexposed, there was no difference between cases and controls (odds ratio: 0.90; 95% confidence interval: 0.52-1.56). It is considered that Af-alb adducts reflect recent past exposure (past 2 to 3 months). Our results therefore suggest that exposure to aflatoxin in the months before diagnosis was not a major contributor to the risk of HCC in this cohort. Because the

Table 2. R249S in plasma specimens matched with the analyzed tumors

plasma ID	Time lapse between plasma collection and cancer diagnosis (mo)	R249S in plasma by SOMA (copies/mL plasma)	Corresponding tumor ID	R249S in tumor by RFLP/seq	Other TP53 mutations (exon 7) in tumor
28-1	0	0	1	Wt	CCC>TCC at 250
162-1	2	0	6	+	
151-1	2	0	9	Wt	
12-4	7	0	15	+	
168-6	21	0	20	Wt	
87-2	4	157	7	Wt	
258-5	54	247	19	+	
170-6	16	490	12	+	
10-6	32	593	17	+	
137-1	1	725	2	+	
90-3	23	817	11	Wt	GT>TT at donor splice site
138-5	0	946	14	+	
40-1	4	2310	5	Wt	
48-1	1	3546	3	+	

region of Qidong is historically known as an area of high exposure to dietary aflatoxin, these results also suggest that exposure to aflatoxin in this area of China has recently decreased as a result of public health intervention and/or changes in dietary and lifestyle patterns.

Discussion

In this study, we have used a nested case-control design to examine the contribution of aflatoxin and HCV to the risk of HCC in a cohort of chronic HBV carriers from Qidong County, PRC, an area of traditionally high exposure to aflatoxins. As a marker of mutagenesis by aflatoxin, we have analyzed the R249S TP53 mutation, which is common in liver cancer in those parts of the world where exposure to aflatoxin is high. First, we showed that the R249S mutation was present in 61% (11 of 18) of HCC cases, confirming the results of previous studies in HCC from the Qidong area. Second, we described the occasional presence of low levels of free plasma DNA containing R249S in plasma specimens collected ahead of diagnosis. Detection of such mutant plasma DNA was, however, not a predictor of HCC development in this cohort. Third, the prevalence of HCV infection was low and does not seem to play a

significant role in HCC etiology in this population. Fourth, individual exposure to aflatoxin, as measured by the detection of Af-alb adducts in plasma, indicates moderate to low exposure in the years and months before HCC diagnosis.

In a previous study using plasma specimens from West Africa (10), we found a good concordance between R249S mutations as detected by RFLP and by SOMA. This was not the case in the present study, perhaps because mutant DNA levels in the plasma were particularly low. In the West African study, about 35% of liver cancer patients were found to carry up to 10,000 copies of mutant plasma DNA/ml of plasma. At these high levels, RFLP gives robust results that are in good agreement with more sensitive methods such as SOMA. At low levels, however, the signals generated by RFLP are below the detection threshold. This explains why some specimens containing trace amounts of R249S were found positive by SOMA but not by RFLP. Thus, on the basis of RFLP alone, we cannot consider that plasma was negative for R249S, but only that plasma may contain levels of R249S that are too low for detection by this method. Unfortunately, due to small initial amounts of plasma and to logistical constraints, it has not been possible to perform SOMA in the plasma of all cases and controls.

Table 3. Prevalence of HCV, R249S TP53, and Af-alb adducts in plasma albumin in cases and controls from the Qidong cohort

(A) HCV			
	Cases, n (%)	Controls, n (%)	Total n (%)
HCV ⁺	2 (1.6)	8 (2.1)	10 (2.0)
HCV ⁻	125 (98.4)	372 (97.9)	497 (98.0)
Total	127 (100.0)	380 (100.0)	507 (100.0)
	Odds ratio, 0.94 (95% confidence interval: 0.45-1.97)		
(B) Af-alb adducts ≥ 3 pg/mg albumin			
	Cases, n (%)	Controls, n (%)	Total n (%)
Af-alb ⁺	84 (66.7)	84 (68.3)	168 (67.5)
Af-alb ⁻	42 (33.3)	39 (31.7)	81 (32.5)
Total	126 (100.0)	123 (100.0)	249 (100.0)
	Odds ratio, 0.90 (95% confidence interval: 0.52-1.56)		

Two explanations, not mutually exclusive, have been proposed for the presence of R249S in the plasma: it may represent a marker of ongoing mutagenesis by aflatoxins or a marker of early carcinogenesis in the liver. In the first instance, it is expected that levels of R249S may increase in subjects with high levels of Af-alb adducts in the plasma. In the second instance, levels of R249S in the plasma should be proportional to the tumor mass and the amount of tumor material released into the bloodstream. In the present study, we found that levels of Af-alb adducts were moderate to low in the whole cohort, indicating that contamination by aflatoxins was not widespread anymore in this population.

The low levels of aflatoxin biomarkers we observed may be a result of the national campaign aimed at reducing the aflatoxin contamination of food in China, combined with the increasing affluence in this East Coast province that has resulted in a widespread change from a maize- to rice-based diet. Wang et al. (17) have shown that the dietary patterns in the Jiangsu province changed remarkably in the 1990s, with a gradual decrease in grain consumption and an important increase in consumption of meat. The uniformly low levels of aflatoxin exposure biomarkers also explain why there is no association in this study between aflatoxin exposure and HCC risk among HBV chronic carriers, in contrast to earlier prospective studies (18, 19). The presence of R249S in 61% of the tumors despite low levels of Af-alb adducts suggests that this mutation may have been acquired by liver cells well ahead of the recruitment of the subjects in the present cohort, at a time when exposure to aflatoxin was higher. A previous study in the same region has found mutations in 55% (15 of 20) of the cases using the same sensitive SOMA method as described here (20). Other studies, using less sensitive methods, have reported roughly similar prevalences (21-23). In all studies, G→T transversion was the most common if not the only mutation type observed. Interestingly, in our study, the mutation prevalence was higher in surgical specimens than in biopsies. This may be due to heterogeneity within tumors, which may not be captured in a small biopsy, or may reflect a selection bias because the surgical specimens were carefully selected by the surgeon as being representative for the whole tumor.

The absence of significant recent exposure to aflatoxin raises the question of the contribution of this toxin in the time sequence of events leading to HCC. In fact, TP53 mutations may occur early in life and persist in a subset of liver cells until cancer onset many years later. The fact that we detected the R249S mutation is found in peritumoral, cirrhotic liver supports the hypothesis that acquisition of the R249S mutation is an event that may take place well ahead of cancer development. This notion is in agreement with results of other studies showing that the mutation was detectable in the plasma of some patients with cirrhosis but not cancer (9, 24, 25). It has been suggested that exposure to aflatoxin in young children may be particularly significant for the permanent acquisition of R249S in liver cells. However, a recent study of 149 young children (aged 2-5 years), from a region in West Africa with a higher frequency of serum Af-alb positivity, did not identify R249S mutations (26). Further age-stratified analyses of plasma DNA, and

when applicable, liver tissues, are required to address the question of timing of occurrence of this mutation in the natural history of HCC.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We thank Nicole Lyandrat for expert immunostaining, Drs. Philippe Tanière and Jean-Yves Scoazec for histopathologic diagnosis, and Anne Sutcliffe for assistance in the analysis of aflatoxin-albumin adducts.

References

- Parkin DM. Global cancer statistics in the year 2000. *Lancet Oncol* 2001;2:533–43.
- Ming L, Thorgeirsson SS, Gail MH, et al. Dominant role of hepatitis B virus and cofactor role of aflatoxin in hepatocarcinogenesis in Qidong, China. *Hepatology* 2002;36:1214–20.
- Wang JS, Qian GS, Zarba A, et al. Temporal patterns of aflatoxin-albumin adducts in hepatitis B surface antigen-positive and antigen-negative residents of Daxin, Qidong County, People's Republic of China. *Cancer Epidemiol Biomarkers Prev* 1996;5:253–61.
- Montesano R, Hainaut P, Wild CP. Hepatocellular carcinoma: from gene to public health. *J Natl Cancer Inst* 1997;89:1844–51.
- Wild CP, Turner PC. The toxicology of aflatoxins as a basis for public health decisions. *Mutagenesis* 2002;17:471–81.
- Smela ME, Hamm ML, Henderson PT, Harris CM, Harris TM, Essigmann JM. The aflatoxin B₁ formamidopyrimidine adduct plays a major role in causing the types of mutations observed in human hepatocellular carcinoma. *Proc Natl Acad Sci U S A* 2002;99:6655–60.
- Stern MC, Umbach DM, Yu MC, London SJ, Zhang ZQ, Taylor JA. Hepatitis B, aflatoxin B₁, and p53 codon 249 mutation in hepatocellular carcinomas from Guangxi, People's Republic of China, and a meta-analysis of existing studies. *Cancer Epidemiol Biomarkers Prev* 2001;10:617–25.
- Hsieh DP, Atkinson DN. Recent aflatoxin exposure and mutation at codon 249 of the human p53 gene: lack of association. *Food Addit Contam* 1995;12:421–4.
- Jackson PE, Kuang SY, Wang JB, et al. Prospective detection of codon 249 mutations in plasma of hepatocellular carcinoma patients. *Carcinogenesis* 2003;24:1657–63.
- Kirk GD, Lesi OA, Mendy M, et al. 249(ser) TP53 mutation in plasma DNA, hepatitis B viral infection, and risk of hepatocellular carcinoma. *Oncogene* 2005;24:5858–67.
- Chen JG, Parkin DM, Chen QG, et al. Screening for liver cancer: results of a randomised controlled trial in Qidong, China. *J Med Screen* 2003;10:204–9.
- Szymanska K, Lesi OA, Kirk GD, et al. Ser-249TP53 mutation in tumour and plasma DNA of hepatocellular carcinoma patients from a high incidence area in the Gambia, West Africa. *Int J Cancer* 2004;110:374–9.
- Laken SJ, Jackson PE, Kinzler KW, et al. Genotyping by mass spectrometric analysis of short DNA fragments. *Nat Biotechnol* 1998;16:1352–6.
- Hamelin R, Jego N, Laurent-Puig P, Vidaud M, Thomas G. Efficient screening of p53 mutations by denaturing gradient gel electrophoresis in colorectal tumors. *Oncogene* 1993;8:2213–20.
- Tanière P, Martel-Planche G, Saurin JC, et al. TP53 mutations, amplification of P63 and expression of cell cycle proteins in squamous cell carcinoma of the oesophagus from a low incidence area in Western Europe. *Br J Cancer* 2001;85:721–6.
- Wild CP, Jiang YZ, Sabbioni G, Chapot B, Montesano R. Evaluation of methods for quantitation of aflatoxin-albumin adducts and their application to human exposure assessment. *Cancer Res* 1990;50:245–51.
- Wang CN, Liang Z, Wei P, et al. Changes in dietary patterns and certain nutrition-related diseases in urban and rural residents of Jiangsu Province, China, during the 1990s. *Biomed Environ Sci* 2002;15:271–6.

18. Qian GS, Ross RK, Yu MC, et al. A follow-up study of urinary markers of aflatoxin exposure and liver cancer risk in Shanghai, People's Republic of China. *Cancer Epidemiol Biomarkers Prev* 1994; 3:3-10.
19. Wang LY, Hatch M, Chen CJ, et al. Aflatoxin exposure and risk of hepatocellular carcinoma in Taiwan. *Int J Cancer* 1996;67: 620-5.
20. Jackson PE, Qian GS, Friesen MD, et al. Specific p53 mutations detected in plasma and tumors of hepatocellular carcinoma patients by electrospray ionization mass spectrometry. *Cancer Res* 2001;61: 33-5.
21. Hsia CC, Nakashima Y, Thorgeirsson SS, et al. Correlation of immunohistochemical staining and mutations of p53 in human hepatocellular carcinoma. *Oncol Rep* 2000;7:353-6.
22. Hsu IC, Metcalf RA, Sun T, Welsh JA, Wang NJ, Harris CC. Mutational hotspot in the p53 gene in human hepatocellular carcinomas. *Nature* 1991;350:427-8.
23. Scorsone KA, Zhou YZ, Butel JS, Slagle BL. p53 mutations cluster at codon 249 in hepatitis B virus-positive hepatocellular carcinomas from China. *Cancer Res* 1992;52:1635-8.
24. Kirk GD, Camus-Randon AM, Mendy M, et al. Ser-249 p53 mutations in plasma DNA of patients with hepatocellular carcinoma from The Gambia. *J Natl Cancer Inst* 2000;92:148-53.
25. Huang XH, Sun LH, Lu DD, et al. Codon 249 mutation in exon 7 of p53 gene in plasma DNA: maybe a new early diagnostic marker of hepatocellular carcinoma in Qidong risk area, China. *World J Gastroenterol* 2003;9:692-5.
26. Turner PC, Sylla A, Kuang SY, et al. Absence of TP53 codon 249 mutations in young Guinean children with high aflatoxin exposure. *Cancer Epidemiol Biomarkers Prev* 2005;14:2053-5.

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

TP53 R249S Mutations, Exposure to Aflatoxin, and Occurrence of Hepatocellular Carcinoma in a Cohort of Chronic Hepatitis B Virus Carriers from Qidong, China

Katarzyna Szymańska, Jian-Guo Chen, Yan Cui, et al.

Cancer Epidemiol Biomarkers Prev 2009;18:1638-1643.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/18/5/1638>

Cited articles This article cites 26 articles, 8 of which you can access for free at:
<http://cebp.aacrjournals.org/content/18/5/1638.full#ref-list-1>

Citing articles This article has been cited by 4 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/18/5/1638.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/18/5/1638>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.