

Research Article

Effects of Supplemental Vitamin D and Calcium on Oxidative DNA Damage Marker in Normal Colorectal Mucosa: A Randomized Clinical Trial

Veronika Fedirko^{1,3}, Roberd M. Bostick^{1,3}, Qi Long^{2,3}, W. Dana Flanders^{1,2,3}, Marjorie L. McCullough⁴, Eduard Sidelnikov^{1,3}, Carrie R. Daniel⁶, Robin E. Rutherford⁵, and Aasma Shaukat⁷

Abstract

The exact antineoplastic effects of calcium and vitamin D₃ in the human colon are unclear. Animal and *in vitro* studies show that these two agents reduce oxidative stress; however, these findings have never been investigated in humans. To address this, we conducted a pilot, randomized, double-blind, placebo-controlled, 2 × 2 factorial clinical trial to test the effects of calcium and vitamin D₃ on a marker of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine (8-OH-dG), in the normal colorectal mucosa. Patients (N = 92) with at least one pathology-confirmed colorectal adenoma were treated with 2 g/d calcium and/or 800 IU/d vitamin D₃ versus placebo over 6 months. Overall labeling and colorectal crypt distribution of 8-OH-dG in biopsies of normal-appearing rectal mucosa were detected by standardized automated immunohistochemistry and quantified by image analysis. After 6 months of treatment, 8-OH-dG labeling along the full lengths of colorectal crypts decreased by 22% (P = 0.15) and 25% (P = 0.10) in the calcium and vitamin D₃ groups, respectively, but not in the calcium plus vitamin D₃ group. The estimated treatment effects were strongest among participants with higher baseline colon crypt vitamin D receptor expression (P = 0.05). Overall, these preliminary results indicate that calcium and vitamin D₃ may decrease oxidative DNA damage in the normal human colorectal mucosa, support the hypothesis that 8-OH-dG labeling in colorectal crypts is a treatable oxidative DNA damage biomarker of risk for colorectal neoplasms, and provide support for further investigation of calcium and vitamin D₃ as chemopreventive agents against colorectal neoplasms. *Cancer Epidemiol Biomarkers Prev*; 19(1); 280–91. ©2010 AACR.

Introduction

Colorectal cancer, the second leading cause of cancer death in the United States (1), is a disease highly correlated with low vitamin D exposure and with the Western-style diet, which is characterized by relatively low calcium consumption (2). Twenty-fold variations in international colon cancer rates, and migration studies showing acquired high risk within a generation, emphasize the importance of environmental exposures, especially diet and physical activity, in the etiology of colorectal cancer (2) and thus to its preventability.

Authors' Affiliations: Departments of ¹Epidemiology and ²Biostatistics and Bioinformatics, Rollins School of Public Health, and ³Winship Cancer Institute, Emory University; ⁴Epidemiology and Surveillance Research Department, American Cancer Society; ⁵Division of Digestive Diseases, Emory University School of Medicine, Atlanta, Georgia; ⁶Nutritional Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, Department of Health and Human Services, Bethesda, Maryland; and ⁷Department of Medicine, GI Division, University of Minnesota, Minneapolis, Minnesota

Corresponding Author: Roberd M. Bostick, Department of Epidemiology, Rollins School of Public Health, Emory University, 1518 Clifton Road Northeast, Atlanta, GA 30322. Phone: 404-727-2671; Fax: 404-727-8737. E-mail: rmbosti@sph.emory.edu

doi: 10.1158/1055-9965.EPI-09-0448

©2010 American Association for Cancer Research.

Currently, there is no complete agreement as to what dietary factors protect against or promote the development of colorectal cancer nor any accepted preneoplastic biomarkers of risk. Further investigation of potential mechanisms whereby dietary agents lead to clinically relevant changes in normal colon tissue and the development of biomarkers of risk derived from such mechanistic understanding are urgently needed.

There is strong biological plausibility and animal experimental evidence for protection against colorectal cancer by calcium and vitamin D (3). Moreover, in epidemiologic studies, higher total calcium intakes have been consistently associated with reduced risk for colorectal neoplasms (4–8), and calcium supplementation reduced adenoma recurrence (9). Also, higher circulating 25-hydroxyvitamin D (25-OH-vitamin D) levels have been associated with reduced risk for colorectal neoplasms (8, 10). However, the antineoplastic effects of calcium and vitamin D on the normal colorectal epithelium remain unclear.

The proposed mechanisms of calcium against colorectal cancer include protection of colonocytes against free bile and fatty acids (11), direct effects on the cell cycle, and modulation of the APC colon carcinogenesis pathway (12). Beyond calcium homeostasis, vitamin D regulates cell cycle events; promotes bile acid degradation; influences growth factor signaling, cell adhesion, and

DNA repair; and modulates >200 genes (12, 13). Recent evidence also indicates that vitamin D and the vitamin D receptor (VDR) are involved in protection against oxidative damage (14-16).

Despite the basic science evidence, there are no published human trials of the effects of vitamin D and/or calcium supplementation on markers of oxidative DNA damage, such as 8-hydroxy-2'-deoxyguanosine (8-OH-dG), in the normal-appearing colorectal mucosa. To address this, we conducted a pilot, randomized, double-blind, placebo-controlled, 2 × 2 factorial chemoprevention clinical trial of supplemental calcium and vitamin D₃, alone and in combination versus placebo over 6 months, to estimate the efficacy of these agents on a panel of biomarkers (including 8-OH-dG) in the normal colorectal mucosa. We hypothesized that calcium and vitamin D₃, alone and in combination, decrease colorectal epithelial oxidative DNA damage.

Patients and Methods

Participant Population

The detailed protocol of study recruitment and procedures was published previously (17). Briefly, eligible patients, ages 30 to 75 years, in general good health, capable of informed consent, with a history of at least one pathology-confirmed adenomatous colorectal polyp within the past 36 months, and no contraindications to calcium or vitamin D supplementation or rectal biopsy procedures and no medical conditions, habits, or medication usage that would otherwise interfere with the study were recruited from the patient population attending the Digestive Diseases Clinic at the Emory Clinic, Emory University. Detailed specific study exclusion criteria were presented elsewhere (17). This study was approved by the Emory University Institutional Review Board. Written informed consent was obtained from each study participant.

Clinical Trial Protocol

Between April 2005 and January 2006, 522 patients passed initial chart screening for eligibility, and 224 (43%) patients were sent an introductory letter followed by a telephone interview. A total of 105 (47%) potential participants attended an eligibility visit during which they were interviewed, signed a consent form, completed questionnaires, provided a blood sample, and started a 1-month placebo run-in period. Diet was assessed with a semiquantitative food frequency questionnaire (18). Medical and pathology records were reviewed. After a 30-day placebo run-in trial, 92 (88%) participants without significant perceived side effects and who had taken at least 80% of their tablets were eligible for randomized assignment. Eligible participants then underwent a baseline rectal biopsy and were randomly assigned to the following four treatment groups: a placebo control group, a 2.0 g elemental calcium (as calcium carbonate in equal doses twice daily) supplementation group, an

800 IU vitamin D₃ supplementation group (400 IU twice daily), and a calcium plus vitamin D supplementation group taking 2.0 g elemental calcium plus 800 IU vitamin D₃ daily.

All study tablets were custom manufactured by Tishcon. The corresponding supplement and placebo pills were identical in size, appearance, and taste. The placebo was free of vitamin D, calcium, magnesium, and chelating agents. Additional details on the rationale for the doses and forms of calcium and vitamin D supplementation forms were described previously (17).

The treatment period was 6 months, and participants attended follow-up visits at 2 and 6 months after randomization and were contacted by telephone between the second and the final follow-up visits. Pill-taking adherence was assessed by questionnaire, interview, and pill count. Participants were instructed to remain on their usual diet and not take any nutritional supplements not in use on entry into the study. At each of the follow-up visits, participants were interviewed and filled out questionnaires. At the last visit, all participants underwent venipuncture and a rectal biopsy procedure. All participants were asked to abstain from aspirin use for 7 days before each biopsy visit. All visits for a given participant were scheduled at the same time of day to control for possible circadian variability in the outcome measures. Factors hypothesized to be related to 8-OH-dG levels in the normal colon mucosa (e.g., antioxidant micronutrient intakes) were assessed at baseline and at the final follow-up visit. Participants did not have to be fasting for their visits and did not take a bowel cleansing preparation or enema.

Tissue Collection and Processing

Six sextant 1.0-mm-thick biopsy specimens were taken from the rectal mucosa 10 cm proximal to the external anal aperture through a rigid sigmoidoscope with a jumbo cup flexible endoscopic forceps mounted on a semiflexible rod. The biopsies were then immediately placed in PBS, oriented under a dissecting microscope and placed in 10% normal buffered formalin, and then transferred to 70% ethanol 24 h after initial placement in formalin. Within 1 week, the biopsies were processed and embedded in paraffin blocks with three biopsies per block.

Laboratory Methods

The paraffin blocks were cut into 3.0- μ m-thick sections, with each level 40 μ m apart. Five slides with four section levels per patient per biomarker were prepared for immunostaining. To uncover the epitope, heat-mediated antigen retrieval was used: slides were placed in a preheated Pretreatment Module (Lab Vision) with 100× citrate buffer (pH 6.0; DAKO S1699) and steamed for 40 min. Then, slides were placed in a DAKO Automated Immunostainer and immunohistochemically processed using a labeled streptavidin-biotin method for 8-OH-dG (mouse monoclonal antibody to 8-OH-dG manufactured

by Abcam, clone number N45.1, at a concentration of 1:100; ref. 19). For each participant, baseline and follow-up biopsy slides were stained in the same batch, and each staining batch included a balance of participants from each treatment group. The slides were not counterstained. After staining, the slides were coverslipped with a Leica CV5000 Coverslipper (Leica Microsystems). In each staining batch of slides, positive and negative control slides were included. Colon adenocarcinoma was used as a control tissue. The negative and positive control slides were treated identically to the patient slides, except that antibody diluent was used rather than primary antibody on the negative control slide. For VDR, slides were processed as described previously but using mouse monoclonal D-6 antibody raised against amino acids 344 to 424 of human VDR (SC-13133; Santa Cruz Biotechnology) at a concentration of 1:7,500 (20, 21).

Image Analysis of Immunohistochemically Detected Biomarkers in Normal Colon Crypts

A quantitative image analysis method (“scoring”) was used to evaluate detected levels of the biomarkers in colon crypts, as depicted in Fig. 1. The major equipment and software for the image analysis procedures were Scanscope CS digital scanner (Aperio Technologies), computer, digital drawing board, Matlab software (MathWorks), CellularEyes Image Analysis Suite (DivEyes), and MySQL (Sun Microsystems). First, slides were scanned with the Aperio Scanscope CS digital scanner, and electronic images were reviewed in the CellularEyes program to identify colon crypts acceptable for analysis. A “scorable” crypt was defined as an intact crypt extending from the muscularis mucosa to the colon lumen (17, 22). Before analysis, images of negative and positive control slides were checked for staining adequacy. Standardized settings were used on all equipment throughout the scoring procedures. The technician reviewed slides in the CellularEyes program and selected two of three biopsies

with 16 to 20 “scorable” hemicypts (one half of the crypt) per biopsy. Using the digital drawing board, the borders of each selected hemicypt were traced. The program then divided the outline into the equally spaced segments with the average widths of normal colonocytes. Finally, the program measured the background-corrected optical density of the biomarker labeling across the entire hemicypt as well as within each segment. The range of optical density for 8-OH-dG labeling was set between 0.04 and 0.20. All resulting data were automatically transferred into the MySQL database. Then, the technician moved to the next identified hemicypt and repeated all the previously described analysis steps. A reliability control sample previously analyzed by the reader was re-analyzed during the course of the trial to determine intrareader scoring reliability by intraclass correlation coefficient, which was 0.94 for 8-OH-dG.

Protocol for Measuring Serum Vitamin D Levels

All laboratory assays for serum 25-OH-vitamin D and 1,25-dihydroxyvitamin D [1,25-(OH)₂-vitamin D] were done by Dr. Bruce Hollis (Medical University of South Carolina) using a RIA method as described previously (23). Serum samples for baseline and follow-up visits for all subjects were assayed together, ordered randomly, and labeled to mask treatment group, follow-up visit, and quality-control replicates. The average intra-assay coefficient of variation was 2.3% for serum 25-OH-vitamin D and 6.2% for 1,25-(OH)₂-vitamin D.

Statistical Analysis

We assessed treatment groups for comparability of characteristics at baseline and at final follow-up by Fisher's exact test for categorical variables and ANOVA for continuous variables.

Several outcome variables were defined to estimate the overall labeling and within-crypt distributions of 8-OH-dG in the crypts. The mean optical density of 8-OH-dG labeling in the crypts was calculated for each patient at




Figure 1. Quantitative image analysis using Aperio Scanscope and CellularEyes software to measure 8-OH-dG labeling in normal-appearing colorectal crypts. **A.** Choosing scorable crypts. **B.** Tracing borders of hemicypt and removing unwanted objects. **C.** Dividing hemicypt into sections. **D.** Storing resulting biomarker staining, tissue and goblet cell optical density data.

baseline and 6 months of follow-up by summing all the densities from all analyzed crypts from the biopsy specimens and dividing by the number of crypts analyzed. Measures of the within-crypt distributions of the marker were calculated for each patient by taking the means of the biomarker densities in various zones of the crypt (e.g., upper 40% and lower 60%).

Primary analyses were based on assigned treatment at the time of randomization regardless of adherence status (intent-to-treat analysis). Mean biomarker densities were calculated for each treatment group for the baseline and 6 months of follow-up visits. Treatment effects were evaluated by assessing the differences in the densities from baseline to the 6 months of follow-up visit between patients in each active treatment group and the placebo group by a repeated-measures linear MIXED effects model. The model included the intercept, follow-up visit effects (baseline and follow-up), and interactions between treatment groups and the follow-up visit effect (the absolute treatment effect). Because optical density is measured in arbitrary units, to provide perspective on the magnitude of the treatment effects, we also calculated relative effects, defined as [(treatment group follow-up mean) / (treatment group baseline mean)] / [(placebo follow-up mean) / (placebo baseline mean)]. The relative effect provides a conservative estimate of the proportional change in the treatment group relative to that in the placebo group. The interpretation of the relative effect is somewhat analogous to that of an odds ratio (e.g., a relative effect of 2.0 would mean that the proportional change in the treatment group was twice as great as that in the placebo group; refs. 17, 24). Because the treatment groups were balanced on risk factors at baseline, no adjustment was made for other covariates in the primary intent-to-treat analyses.

The distributions of 8-OH-dG staining density were graphically evaluated using the LOESS procedure with a smoothing parameter of 0.5 and local quadratic fitting. First, the number of sections within a hemicypt was standardized to 50. Then, the average for each section across all crypts was predicted by the LOESS model separately for each patient and then for each treatment group by follow-up visit. The results were plotted in the graphs along with smoothing lines.

A questionnaire-derived oxidative balance score (OBS) was calculated as described previously (25, 26). Briefly, continuous variables that reflect pro-oxidant (saturated fat and total iron intake) and antioxidant (total tocopherol, carotenoid, vitamin C, lycopene, lutein/zeaxanthin, and β -cryptoxanthin intake) exposures were divided into high and low categories based on the median value among all participants at baseline. Participants with low (below median) exposure to a particular pro-oxidant were awarded 1 point, whereas those with high (above median) exposure to the same pro-oxidant were awarded 0 point. For antioxidant exposure, 1 point was awarded for each high-level (above median) exposure and 0 point for each low-level (below median) exposure. For dichotomous variables ("yes" versus "no"), participants re-

ceived 1 point for each antioxidant exposure (regular use of nonsteroidal anti-inflammatory drugs (NSAID) and/or aspirin, supplementation with selenium, and never smokers). Then, the points assigned for each individual component of OBS were summed up to calculate the overall score. Lower OBS values indicate a higher prevalence of pro-oxidant exposures, whereas higher OBS values indicate a predominance of antioxidant exposures. The range of the baseline OBS in this study was between 3 and 10, and the median was 6. We dichotomized baseline OBS based on the median value and assigned each participant to a high OBS (above median, "antioxidant") or low OBS (below median, "pro-oxidant") category. Similarly, continuous variables (e.g., age and VDR expression) were dichotomized (into high/low categories) based on the median value in all study participants at baseline. Then, stratified analyses were conducted to explore differential treatment effects by baseline age (<60 and \geq 60 years), VDR expression (high/low), 8-OH-dG labeling (high/low), OBS (\leq 6 and >6), first-degree family history of colorectal cancer (yes/no), sex (male/female), regular NSAID use (yes/no), serum 25-OH-vitamin D levels (<22 and \geq 22 ng/mL), and adherence to treatment (<80% or \geq 80% treatment pills taken). Differences between categories were tested by including the category-intervention interaction term in the model.

Statistical analyses were done using SAS System software version 9.1.3 (SAS Institute). A cutoff level of $P \leq 0.05$ (two-sided) was used for assessing statistical significance.

Results

Characteristics of Study Participants

Treatment groups did not differ significantly on participant characteristics measured at baseline (Table 1) or at the end of the study (data not shown). The mean age of participants was 61 years, 64% were men, 71% were White, and 20% had a family history of colorectal cancer in a first-degree relative. Most participants were overweight, nonsmokers, college graduates, and had a single small mildly dysplastic tubular adenoma (Table 1).

Adherence to visit attendance averaged 92% and did not differ significantly among the four treatment groups. On average, at least 80% of pills were taken by 93% of participants at the first follow-up visit and by 84% at the final follow-up visit. There were no treatment or biopsy complications. Seven (8%) were lost to follow-up due to perceived drug intolerance ($n = 2$), unwillingness to continue participation ($n = 3$), physician's advice ($n = 1$), and cardiovascular disease death ($n = 1$). Dropouts included one person from the vitamin D supplementation group and two persons from each of the other three groups.

At baseline, there were no significant differences between the four study groups in serum 25-OH-vitamin D

Table 1. Selected baseline characteristics of the study participants (N = 92)

Characteristics	Treatment group				P*
	Placebo (n = 23)	Calcium (n = 23)	Vitamin D (n = 23)	Calcium + vitamin D (n = 23)	
Demographics, medical history, habits, and anthropometrics					
Age, y	58.5 (8.2)	61.9 (8.2)	60.2 (8.1)	62.1 (7.5)	0.39
Men (%)	70	70	70	70	1.00
White (%)	74	83	65	61	0.39
College graduate (%)	65	61	57	44	0.53
History of colorectal cancer in first-degree relative (%)	17	30	17	13	0.60
Take NSAID regularly [†] (%)	22	13	9	22	0.60
Take aspirin regularly [†] (%)	22	52	30	56	0.05
If woman (n = 28), taking estrogens (%)	4	9	4	4	1.00
Current smoker (%)	9	4	0	0	0.61
Take multivitamin (%)	30	30	26	39	0.86
Body mass index, kg/m ²	30.6 (7.2)	29.4 (5.5)	28.9 (5.6)	31.6 (6.0)	0.44
Mean dietary intakes					
Total energy intake, kcal/d	1,596 (528)	1,788 (691)	1,848 (821)	1,845 (752)	0.59
Total [‡] calcium, mg/d	618 (308)	746 (335)	843 (526)	824 (714)	0.41
Total [‡] vitamin D, IU/d	277 (230)	336 (202)	360 (317)	415 (316)	0.40
Total fat, g/d	67 (32)	72 (35)	70 (32)	74 (28)	0.59
Dietary fiber, g/d	15 (7)	17 (9)	18 (9)	17 (11)	0.97
Alcohol, g/d	9 (14)	11 (15)	14 (18)	10 (20)	0.84
OBS [§]	6 (2)	7 (2)	7 (2)	7 (2)	0.46
Adenoma characteristic					
Multiple adenomas (%)	17	22	39	26	0.45
Large adenoma ≥1 cm [¶] (%)	19	32	17	9	0.32
Villous/tubulovillous adenoma ^{**} (%)	4	9	9	4	1.00
Mild dysplasia (%)	100	96	100	100	1.00

NOTE: Data are given as mean (SD) unless otherwise specified.

*By Fisher's exact χ^2 test for categorical variables and ANOVA for continuous variables.

[†]At least once a week.

[‡]Diet plus supplements.

[§]See Statistical Analysis for details.

^{||}At least two adenomas.

[¶]At least one large adenoma.

^{**}At least one villous or tubulovillous adenoma.

or 1,25-(OH)₂-vitamin D levels. At the study end, the vitamin D and calcium plus vitamin D groups had significantly higher levels of serum 25-OH-vitamin D ($P < 0.001$), whereas the placebo and calcium groups had slight nonsignificant decreases in 25-OH-vitamin D levels (Table 2A). As expected, serum levels of 1,25-(OH)₂-vitamin D at the end of follow-up period did not differ significantly between study groups (17).

Graphical Assessment of Changes over 6 Months in the Distribution of 8-OH-dG Labeling along Normal Colorectal Crypts

The distribution of 8-OH-dG staining optical density ("labeling") along the colorectal crypts at the baseline and 6 months of follow-up visits is shown in Fig. 2. In

each treatment group, 8-OH-dG labeling appeared to be the highest in the lower 20% to 30% of the crypts, to decrease in the middle part of the crypts, and then to increase somewhat again toward the colon lumen. The baseline distribution of 8-OH-dG along the crypts in all four treatment groups appeared to be almost identical in shape and optical density range. In the placebo group, from baseline to follow-up, 8-OH-dG labeling appeared to increase slightly in the middle part of the crypts (Fig. 2A). A large postsupplemental decrease in 8-OH-dG labeling along the full lengths of the crypt was noted in the calcium and vitamin D groups (Fig. 2B and C). In the vitamin D plus calcium group, similar to the placebo group, 8-OH-dG labeling slightly increased from baseline to follow-up (Fig. 2D).

Effects of Calcium and/or Vitamin D Supplementation on 8-OH-dG Labeling in Normal Colorectal Crypts

At baseline, there were no differences in 8-OH-dG labeling along the full lengths of crypts among the four treatment groups. Relative to placebo, 8-OH-dG labeling along the full lengths of crypts decreased by 22% ($P = 0.14$) in the calcium group and by 25% ($P = 0.10$) in the vitamin D group and increased by 6% ($P = 0.70$) in the calcium plus vitamin D group (Table 2B). The findings for the upper 40% and the lower 60% of the crypts (the differentiation and proliferation zones, respectively; Table 2B) and for the upper and lower 20% (areas closest to and furthest from colon lumen exposures, respectively; data not shown) did not differ substantially from those for the entire crypts.

Stratified Analyses

We investigated whether change in 25-OH-vitamin D levels, adherence to treatment, family history of colorectal cancer, sex, age, smoking, NSAID use, baseline OBS, and baseline batch-standardized VDR expression or 8-OH-dG labeling modified response to treatment; however, the sample size was too small for most of these results to be reliable. The effect of treatment on 8-OH-dG variables did not vary by age, smoking status, family history of colorectal cancer, NSAID use, or change in serum 25-OH-vitamin D levels (data not shown). In women, 8-OH-dG labeling decreased only in the calcium group (-25% ; $P = 0.43$); however, in men, 8-OH-dG labeling decreased in all three active treatment groups (Table 3). In those with a high ("antioxidant") baseline OBS, 8-OH-dG labeling decreased in all three active treatment groups

Table 2. Serum 25-OH-vitamin D and optical density of immunohistochemically detected 8-OH-dG in colorectal crypts at baseline and 6 months of follow-up

	Baseline			6 mo follow-up			Absolute treatment effect*			Relative effect†
	n	Mean (SE)	P‡	n	Mean (SE)	P‡	n	Mean (SE)	P‡	
(A) Serum vitamin D measurements										
25-OH-vitamin D, ng/mL										
Placebo	23	20.7 (1.7)		21	18.2 (1.8)		21	0		1.00
Calcium	23	25.7 (1.7)	0.05	21	23.4 (1.7)	0.03	21	0.2 (1.8)	0.92	1.03
Vitamin D	23	21.0 (1.7)	0.81	22	29.5 (1.7)	<0.0001	22	10.9 (1.8)	<0.0001	1.59
Calcium + vitamin D	23	20.9 (1.7)	0.84	21	28.9 (1.7)	<0.0001	21	10.5 (1.8)	<0.0001	1.57
(B) 8-OH-dG [§] labeling optical density in colorectal crypts										
Entire crypts										
Placebo	23	2,360.8 (193.2)		21	2,509.0 (202.1)		21	0		1.00
Calcium	23	2,349.4 (193.2)	0.97	21	1,946.2 (202.1)	0.05	21	-551.5 (374.4)	0.14	0.78
Vitamin D	23	2,318.4 (193.2)	0.88	22	1,847.3 (197.5)	0.02	22	-619.3 (372.0)	0.10	0.75
Calcium + vitamin D	23	2,347.8 (193.2)	0.96	21	2,642.6 (202.1)	0.64	21	146.5 (264.8)	0.70	1.06
Upper 40% of crypts										
Placebo	22	677.7 (64.1)		21	655.9 (67.0)		21	0		1.00
Calcium	23	704.7 (64.1)	0.77	21	525.1 (67.0)	0.17	21	-157.8 (125.8)	0.21	0.77
Vitamin D	23	684.4 (64.1)	0.94	22	505.9 (65.5)	0.11	22	-156.8 (125.0)	0.21	0.76
Calcium + vitamin D	23	655.0 (64.1)	0.80	21	741.5 (67.0)	0.37	21	108.3 (125.8)	0.39	1.17
Lower 60% of crypts										
Placebo	22	1,418.7 (112.0)		21	1,459.7 (117.2)		21	0		1.00
Calcium	23	1,431.8 (112.0)	0.93	21	1,201.5 (117.2)	0.12	21	-271.3 (213.5)	0.21	0.82
Vitamin D	23	1,450.7 (112.0)	0.84	22	1,145.1 (114.5)	0.06	22	-346.6 (212.1)	0.11	0.77
Calcium + vitamin D	23	1,390.0 (112.0)	0.86	21	1,556.8 (117.2)	0.56	21	125.8 (213.5)	0.56	1.09

*Absolute treatment effect = [(treatment group follow-up) - (treatment group baseline)] - [(placebo group follow-up) - (placebo group baseline)].

†Relative effect = [(treatment group follow-up) / (treatment group baseline)] / [(placebo follow-up) / (placebo baseline)]; interpretation similar to that for an odds ratio (e.g., a relative effect of 1.7 indicates a proportional increase of 70% in the treatment group relative to that in the placebo group).

‡P value for difference between each active treatment group and placebo group from repeated-measures MIXED model.

§Biomarker was detected immunohistochemically and then its labeling optical density was quantified by image analysis (see text for details).




Figure 2. Distribution of 8-OH-dG staining optical density along normal colorectal crypts by treatment group at baseline and follow-up. **A.** Placebo group. **B.** Calcium group. **C.** Vitamin D group. **D.** Vitamin D plus calcium group.

after 6 months of treatment, whereas in those with low (“pro-oxidant”) baseline OBS, 8-OH-dG decreased only in the vitamin D group (-19% ; $P = 0.40$; Table 3).

There were no substantial differences in the estimated treatment effects according to baseline levels of 8-OH-dG labeling (data not shown). Among those with high baseline colorectal crypt VDR expression, 8-OH-dG labeling decreased by 35% ($P = 0.09$) in the calcium group, 54% ($P = 0.003$) in the vitamin D group, and 17% ($P = 0.34$) in the calcium plus vitamin D group relative to the placebo, whereas there were no decreases seen in those with low baseline VDR expression and there was a 75% increase in 8-OH-dG labeling in the calcium plus vitamin D group relative to the placebo. The test for interaction for treatment effect by VDR status was statistically significant ($P = 0.05$; Table 3).

Discussion

The results from this pilot, randomized, controlled clinical trial suggest that supplementation with calcium or

vitamin D_3 , but not with both agents combined, may decrease oxidative DNA damage, as indicated by decreased 8-OH-dG immunohistochemical labeling, in the normal-appearing colorectal epithelium of sporadic adenoma patients. These findings are consistent with the hypothesis that high intakes of calcium or vitamin D_3 may decrease oxidative stress and oxidative DNA damage in the colon and thus reduce risk for colorectal neoplasms. Our findings also suggest that vitamin D_3 combined with calcium may have either a lesser or no treatment effect on 8-OH-dG labeling than does either calcium or vitamin D alone. Consistent with existing animal data (15, 27), we found evidence that baseline VDR expression levels may modify treatment effects of calcium and vitamin D_3 , such that those with higher colorectal crypt VDR expression may be more strongly responsive to treatment. Finally, the treatment effect of calcium and vitamin D_3 tended to be stronger in men and those with higher baseline antioxidant relative to pro-oxidant exposures.

Oxidative stress, a condition characterized by an imbalance of pro-oxidants to antioxidants, which results in macromolecular damage and disruption of redox signaling

and control (28), may play a role in colon carcinogenesis, inducing protein and DNA damage and lipid peroxidation and impairing intracellular signaling. Under normal conditions, reactive oxygen species have an important role as intracellular signaling molecules that regulate many genes (29). However, under inflammatory conditions, increased generation of reactive oxygen species products leads to cell molecule damage such as oxidation of DNA

(29). The most abundant product of oxidative DNA modifications by reactive oxygen species is 8-OH-dG (30). This oxidized base is a useful biomarker of oxidative stress that can be measured in urine, blood, and tissues (19, 31). Several studies showed increased levels of oxidatively modified DNA in colorectal adenocarcinomas when compared with adenomas and adjacent normal epithelium (32, 33). This suggests that inhibition of oxidative stress in the

Table 3. 8-OH-dG labeling in colorectal crypts stratified by sex, baseline OBS, and baseline colorectal crypt VDR expression

	<i>n</i>	Absolute treatment effect* Mean (SE)	<i>P</i> [†]	Relative effect [‡]	<i>P</i> [§]
Women					
Placebo	7	0		1.00	
Calcium	7	-577.4 (714.7)	0.43	0.75	
Vitamin D	6	253.3 (729.4)	0.73	1.11	
Calcium + vitamin D	7	508.3 (714.7)	0.48	1.23	
Men					
Placebo	14	0		1.00	
Calcium	14	-526.1 (422.9)	0.22	0.80	
Vitamin D	16	-959.0 (414.3)	0.02	0.62	
Calcium + vitamin D	14	-26.2 (422.9)	0.95	0.99	0.35
High OBS					
Placebo	8	0		1.00	
Calcium	13	-898.0 (594.8)	0.14	0.67	
Vitamin D	12	-879.0 (599.7)	0.15	0.67	
Calcium + vitamin D	11	-120.5 (615.1)	0.85	0.94	
Low OBS					
Placebo	13	0		1.00	
Calcium	8	25.1 (509.7)	0.96	1.02	
Vitamin D	9	-421.6 (498.6)	0.40	0.81	
Calcium + vitamin D	10	383.2 (478.1)	0.43	1.19	0.71
High baseline colorectal crypt VDR expression					
Placebo	10	0		1.00	
Calcium	10	-827.0 (477.0)	0.09	0.68	
Vitamin D	8	-1,626.1 (506.7)	0.003	0.46	
Calcium + vitamin D	11	-443.6 (462.5)	0.34	0.83	
Low baseline colorectal crypt VDR expression					
Placebo	8	0		1.00	
Calcium	9	264.6 (532.4)	0.62	1.11	
Vitamin D	13	63.5 (498.8)	0.90	1.00	
Calcium + vitamin D	8	1,420.4 (551.5)	0.02	1.75	0.05

*Absolute treatment effect = [(treatment group follow-up) – (treatment group baseline)] – [(placebo group follow-up) – (placebo group baseline)].

[†]*P* value for difference between each active treatment group and placebo group from repeated-measures MIXED model.

[‡]Relative effect = [(treatment group follow-up) / (treatment group baseline)] / [(placebo follow-up) / (placebo baseline)]; interpretation similar to that for an odds ratio (e.g., a relative effect of 1.7 indicates a proportional increase of 70% in the treatment group relative to that in the placebo group).

[§]*P* value for the category-intervention interaction term.

^{||}OBS and baseline colorectal crypt VDR expression were dichotomized into high/low categories based on the median value in all study participants at baseline. OBS was calculated as described in Statistical Analysis. VDR was detected immunohistochemically and then its labeling optical density was quantified by image analysis (see text for details).

normal colorectal epithelium may slow down or prevent carcinogenesis and prompts the development of chemopreventive agents such as calcium and vitamin D that target oxidative stress in the colon.

There are several lines of evidence to support our hypotheses that calcium and vitamin D may act as antioxidants and DNA damage-reducing agents in the colon. Bile acids damage cell membranes, at least in part through an oxidative mechanism (34, 35), provoking an inflammatory response and causing DNA damage (36), and both calcium and vitamin D can reduce the free bile acid load in the colon lumen. Calcium directly binds bile acids, rendering them inert (37). Vitamin D activation of the ubiquitous VDR in the colon upregulates CYP3A4, which in turn catabolizes the secondary bile acid, lithocholic acid (38, 39). Furthermore, high blood 25-OH-vitamin D levels provide a pool of vitamin D that is available for various tissues, such as the colorectal epithelium. In colonocytes, vitamin D increases expression of enzymes involved in antioxidant response, inhibits iron-dependent lipid peroxidation in liposomes, lowers glutathione reductase levels, induces glutathione peroxidase and manganese-dependent superoxide dismutase activity, and elevates glutathione levels (refs. 16, 40; also reviewed in ref. 14), thereby decreasing oxidative stress in the colorectal epithelium. The results of this study, combined with the biological evidence, support calcium and vitamin D₃ as oxidative DNA damage-reducing agents.

Contrary to our original hypothesis and to what has been described in some epidemiologic and clinical studies (41-45), we did not observe a treatment effect in the calcium plus vitamin D group. We also previously reported that vitamin D combined with calcium may have lesser treatment effects on colorectal epithelial apoptosis and differentiation than does calcium or vitamin D separately (17, 46). There are several possible explanations for this finding. Considering the small sample size of the study, the lack of treatment effect in the calcium and vitamin D group may have been due to chance. It is also possible that the two agents may have attenuated the effects of one another. 1,25-(OH)₂-vitamin D₃ regulates calcium homeostasis (47). As calcium concentration decreases, the production of 1,25-(OH)₂-vitamin D₃ increases, which in turn increases intestinal calcium absorption (47). Elevated calcium in the diet may suppress 1,25-(OH)₂-vitamin D₃ synthesis at the cellular level, which in turn may also attenuate activation of vitamin D-responsive detoxifying enzymes. One animal study (48) found that calcium and vitamin D were more potent inhibitors of colon tumorigenesis when given separately, but several other animal studies reported synergistic effects with calcium and vitamin D combined (49, 50). A large adenoma recurrence trial also supported an enhanced chemopreventive effect of vitamin D with calcium (42). Taken altogether, the combined effect of calcium and vitamin D on oxidative DNA damage in colorectal epithelium is unclear and will require clarification through larger studies.

In contrast to that in men, there was no evidence for a treatment effect of vitamin D alone and in combination with calcium on colorectal crypt 8-OH-dG labeling levels in women. There are several possible explanations for this finding, including a very low statistical power to detect treatment effects due to the small sample size. Another possible explanation may be that women in our study may have had decreased estrogen levels, as the majority of them were postmenopausal and not taking estrogens. The Women's Health Initiative Hormone Replacement Therapy Trial (51) found that endogenous estrogen plus progestin therapy, but not estrogen alone therapy, reduced risk for colorectal cancer (52). However, one human study found that an estrogen intervention activated the VDR pathway and downregulated inflammatory and immune signaling pathways in the rectal mucosa of postmenopausal women (53). Therefore, the findings of our study are consistent with the hypothesis that low estrogen levels may interfere with VDR signaling in the colorectal mucosa, resulting in no changes in 8-OH-dG levels after supplementation with vitamin D; however, further studies are needed to clarify these issues.

Epidemiologic studies have not consistently found associations of colorectal neoplasms with individual pro-oxidant and antioxidant factors despite the strong biological rationale and basic science evidence. Several analyses (25, 26) suggested that oxidative stress, inflammation, diet, and other risk factors may synergistically or individually affect risk of colorectal neoplasia through multiple pathways. Therefore, we used the OBS to categorize patients into different oxidative stress profiles based on multiple determinants of oxidative stress. We hypothesized that patients with different oxidative stress profiles may respond differently to calcium and/or vitamin D. Those with a high baseline OBS (higher balance of antioxidant to pro-oxidant exposures) had greater estimated calcium and calcium plus vitamin D treatment effects on 8-OH-dG labeling than those with a low OBS. A low OBS reflects low total intakes of antioxidants such as vitamin C and carotene combined with high pro-oxidant exposures such as high fat or iron intakes. In the colon lumen, free calcium directly binds bile acids (37), thereby reducing procarcinogenic effects of bile acids on the colorectal epithelium. Persons with high fat intake have higher colonic lumen levels of deoxycholic and lithocholic bile acids (36) and may require more calcium to neutralize the DNA-damaging bile acids than do persons on a low-fat diet. Antioxidant enzymes in humans function in combination with low-weight antioxidant compounds such as vitamin C, α -tocopherol, and β -carotene (54). In the colorectal epithelium, vitamin D activates the expression of antioxidant enzymes (14), which may not function properly in the antioxidant-depleted environment. Therefore, it is possible that calcium and vitamin D effects on the oxidative DNA damage marker 8-OH-dG are modified by the presence or absence of various pro-oxidant or antioxidant exposures.

On the other hand, our findings may have been due to chance and further investigations are required to confirm them.

Because complete loss of the VDR significantly increased 8-OH-dG labeling in the mouse colon (15, 27), we hypothesized that different VDR expression levels in the normal-appearing colorectal mucosa modify vitamin D treatment effects. Consistent with this hypothesis, we observed substantial decreases in 8-OH-dG labeling in study participants with high, but not low, baseline VDR expression.

This study has several limitations. First, treatment effects of vitamin D and calcium on the oxidative DNA damage marker 8-OH-dG in parts of the colon other than the rectum are unclear, as we did not collect tissue biopsies from different parts of the colon and there are no published studies of 8-OH-dG labeling throughout the colon. Another potential limitation of this study is that it is not known whether oxidative stress markers are associated with risk for colon cancer in humans. However, substantial published literature supports the plausibility of an important role for increased oxidative DNA damage in colon carcinogenesis, especially for the transition from colorectal adenoma to carcinoma (32, 33). Persistent oxidative stress leads to protein and DNA damage and lipid peroxidation, which can cause genetic and epigenetic alterations, and may facilitate the development of neoplasia from the normal colorectal mucosa (29). Therefore, 8-OH-dG in the normal colorectal mucosa may serve as a biomarker of risk for colorectal neoplasms. Finally, the most obvious limitation of the study is the small sample size, which may have increased the probability of chance findings in detecting or not detecting a treatment effect.

The strengths of this study include the randomized, double-blind, placebo-controlled trial design; high protocol adherence by study participants; examination of both independent and combined effects of calcium and vitamin D₃ on an oxidative stress marker; automated standardized biopsy handling and immunostaining procedures; and use of cutting-edge technologies to conduct the quantitative image analyses. Another strength of this study is that we used immunohistochemical detection of 8-OH-dG in the colorectal epithelium as it was important to detect 8-OH-dG in colonocytes but not in infiltrating lymphocytes or other intermingled cells. Such detection was made possible by the development of a specific monoclonal antibody against 8-OH-dG (19) and our novel image analysis methods. High-performance liquid chromatography, an alternative method of measuring 8-OH-dG in colon tissue, may overestimate oxidative DNA damage in the colonocytes, especially in the presence of inflammation. Finally, this study is the first human study to test the effect of

calcium and/or vitamin D₃ on an oxidative DNA damage marker in the normal-appearing colorectal mucosa.

Overall, these preliminary results from this pilot clinical trial suggest that calcium and vitamin D, given separately, may decrease oxidative DNA damage in the normal-appearing colorectal epithelium; the treatment effects of calcium and vitamin D on oxidative DNA damage marker 8-OH-dG may be strongest in those with higher VDR expression in the colon; and 8-OH-dG may be a modifiable biomarker of oxidative stress that can be used in colon cancer-related chemoprevention trials to assess treatment efficacy. Our results support further investigations of calcium and vitamin D as chemopreventive agents against colorectal neoplasms.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Jill Joelle Woodard and Bonita Feinstein for managing the study, Dr. Bruce W. Hollis for conducting blood vitamin D assays, Vaunita Cohen and Eileen Veronica Smith for excellent technical assistance, Christopher Farino and Stuart Myerberg for development of the study database, John Melonakos and Tauseef Rehman from DivEyes for development of the scoring software, the physicians of the Emory Clinic for work on biopsy procurement, and all study participants for their time and dedication to the study.

Grant Support

National Cancer Institute, NIH grants RO1 CA104637 and R03 CA136113 and Georgia Cancer Coalition Distinguished Scholar Award (R.M. Bostick); Franklin Foundation; and Emory Graduate School supplemental research funds (V. Fedirko).

The National Cancer Institute, the Georgia Cancer Coalition, the Franklin Foundation, and the Emory Graduate School had no influence on the design of the study; the collection, analysis, and interpretation of the data; the decision to submit the manuscript for publication; or the writing of the manuscript.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 5/11/09; revised 9/23/09; accepted 10/22/09; published online 1/7/10.

References

1. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ. Cancer statistics, 2009. *CA Cancer J Clin* 2009;59:225–49.
2. Potter JD, Slattery ML, Bostick RM, Gapstur SM. Colon cancer: a review of the epidemiology. *Epidemiol Rev* 1993;15:499–545.
3. Lipkin M, Lamprecht SA. Mechanisms of action of vitamin D: recent findings and new questions. *J Med Food* 2006;9:135–7.
4. Oh K, Willett WC, Wu K, Fuchs CS, Giovannucci EL. Calcium and

- vitamin D intakes in relation to risk of distal colorectal adenoma in women. *Am J Epidemiol* 2007;165:1178–86.
5. Kesse E, Boutron-Ruault MC, Norat T, Riboli E, Clavel-Chapelon F. Dietary calcium, phosphorus, vitamin D, dairy products and the risk of colorectal adenoma and cancer among French women of the E3N-EPIC prospective study. *Int J Cancer* 2005;117:137–44.
 6. Peters U, Chatterjee N, McGlynn KA, et al. Calcium intake and colorectal adenoma in a US colorectal cancer early detection program. *Am J Clin Nutr* 2004;80:1358–65.
 7. Cho E, Smith-Warner SA, Spiegelman D, et al. Dairy foods, calcium, and colorectal cancer: a pooled analysis of 10 cohort studies. *J Natl Cancer Inst* 2004;96:1015–22.
 8. Bostick RM, Goodman M, Sidelnikov E. Calcium and vitamin D. In: Potter JD, Lindor NM, editors. *Genetics of colorectal cancer*. New York (NY): Springer Science + Business Media; 2009, p. 277–96.
 9. Weingarten MA, Zalmanovici A, Yaphe J. Dietary calcium supplementation for preventing colorectal cancer and adenomatous polyps. *Cochrane Database Syst Rev* 2008;1:CD003548.
 10. Wei MY, Garland CF, Gorham ED, Mohr SB, Giovannucci E. Vitamin D and prevention of colorectal adenoma: a meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2008;17:2958–69.
 11. Newmark HL, Lipkin M. Calcium, vitamin D, colon cancer. *Cancer Res* 1992;52:2067–70s.
 12. Lamprecht SA, Lipkin M. Chemoprevention of colon cancer by calcium, vitamin D and folate: molecular mechanisms. *Nat Rev Cancer* 2003;3:601–14.
 13. Ebert R, Schutze N, Adamski J, Jakob F. Vitamin D signaling is modulated on multiple levels in health and disease. *Mol Cell Endocrinol* 2006;248:149–59.
 14. Chatterjee M. Vitamin D and genomic stability. *Mutat Res* 2001;475:69–87.
 15. Kallay E, Bareis P, Bajna E, et al. Vitamin D receptor activity and prevention of colonic hyperproliferation and oxidative stress. *Food Chem Toxicol* 2002;40:1191–6.
 16. Kutuzova GD, DeLuca HF. 1,25-Dihydroxyvitamin D₃ regulates genes responsible for detoxification in intestine. *Toxicol Appl Pharmacol* 2007;218:37–44.
 17. Fedirko V, Bostick RM, Flanders WD, et al. Effects of vitamin D and calcium supplementation on markers of apoptosis in normal colon mucosa: a randomized, double-blind, placebo-controlled clinical trial. *Cancer Prev Res* 2009;2:213–23.
 18. Willett WC, Sampson L, Browne ML, et al. The use of a self-administered questionnaire to assess diet four years in the past. *Am J Epidemiol* 1988;127:188–99.
 19. Toyokuni S, Tanaka T, Hattori Y, et al. Quantitative immunohistochemical determination of 8-hydroxy-2'-deoxyguanosine by a monoclonal antibody N45.1: its application to ferric nitrilotriacetate-induced renal carcinogenesis model. *Lab Invest* 1997;76:365–74.
 20. Hidalgo AA, Paredes R, Garcia VM, et al. Altered VDR-mediated transcriptional activity in prostate cancer stroma. *J Steroid Biochem Mol Biol* 2007;103:731–6.
 21. Tishkoff DX, Nibbelink KA, Holmberg KH, Dandu L, Simpson RU. Functional vitamin D receptor (VDR) in the t-tubules of cardiac myocytes: VDR knockout cardiomyocyte contractility. *Endocrinology* 2008;149:558–64.
 22. Bostick RM, Fosdick L, Lillemo T, et al. Methodological findings and considerations in measuring colorectal epithelial cell proliferation in humans. *Cancer Epidemiol Biomarkers Prev* 1997;6:931–42.
 23. Hollis BW. Quantitation of 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D by radioimmunoassay using radioiodinated tracers. *Methods Enzymol* 1997;282:174–86.
 24. Bostick RM, Fosdick L, Wood JR, et al. Calcium and colorectal epithelial cell proliferation in sporadic adenoma patients: a randomized, double-blinded, placebo-controlled clinical trial. *J Natl Cancer Inst* 1995;87:1307–15.
 25. Goodman M, Bostick RM, Dash C, Flanders WD, Mandel JS. Hypothesis: oxidative stress score as a combined measure of pro-oxidant and antioxidant exposures. *Ann Epidemiol* 2007;17:394–9.
 26. Goodman M, Bostick RM, Dash C, Terry P, Flanders WD, Mandel J. A summary measure of pro- and anti-oxidant exposures and risk of incident, sporadic, colorectal adenomas. *Cancer Causes Control* 2008;19:1051–64.
 27. Kallay E, Pietschmann P, Toyokuni S, et al. Characterization of a vitamin D receptor knockout mouse as a model of colorectal hyperproliferation and DNA damage. *Carcinogenesis* 2001;22:1429–35.
 28. Sies H, Jones DP. Oxidative stress. In: Fink G, editor. *Encyclopedia of stress*. 2nd ed. Amsterdam, Boston: Elsevier, Academic Press; 2007, p. 45–8.
 29. Roessner A, Kuester D, Malfertheiner P, Schneider-Stock R. Oxidative stress in ulcerative colitis-associated carcinogenesis. *Pathol Res Pract* 2008;204:511–24.
 30. Evans M, Cooke M. *Oxidative damage to nucleic acids*. Illustrated ed. Austin, Tex.; New York: Springer 2007.
 31. Kasai H. Analysis of a form of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis. *Mutat Res* 1997;387:147–63.
 32. Kondo S, Toyokuni S, Iwasa Y, et al. Persistent oxidative stress in human colorectal carcinoma, but not in adenoma. *Free Radic Biol Med* 1999;27:401–10.
 33. Ribeiro ML, Priolli DG, Miranda DD, Arcari DP, Pedrazzoli J, Jr., Martinez CA. Analysis of oxidative DNA damage in patients with colorectal cancer. *Clin Colorectal Cancer* 2008;7:267–72.
 34. Venturi M, Hambly RJ, Glinghammar B, Rafter JJ, Rowland IR. Genotoxic activity in human faecal water and the role of bile acids: a study using the alkaline comet assay. *Carcinogenesis* 1997;18:2353–9.
 35. Babbs CF. Free radicals and the etiology of colon cancer. *Free Radic Biol Med* 1990;8:191–200.
 36. Bernstein H, Bernstein C, Payne CM, Dvorakova K, Garewal H. Bile acids as carcinogens in human gastrointestinal cancers. *Mutat Res* 2005;589:47–65.
 37. Newmark HL, Wargovich MJ, Bruce WR. Colon cancer and dietary fat, phosphate, and calcium: a hypothesis. *J Natl Cancer Inst* 1984;72:1323–5.
 38. Harris DM, Go VL. Vitamin D and colon carcinogenesis. *J Nutr* 2004;134:3463–71S.
 39. Lamprecht SA, Lipkin M. Cellular mechanisms of calcium and vitamin D in the inhibition of colorectal carcinogenesis. *Ann N Y Acad Sci* 2001;952:73–87.
 40. Sardar S, Chakraborty A, Chatterjee M. Comparative effectiveness of vitamin D₃ and dietary vitamin E on peroxidation of lipids and enzymes of the hepatic antioxidant system in Sprague-Dawley rats. *Int J Vitam Nutr Res* 1996;66:39–45.
 41. Lappe JM, Travers-Gustafson D, Davies KM, Recker RR, Heaney RP. Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial. *Am J Clin Nutr* 2007;85:1586–91.
 42. Grau MV, Baron JA, Sandler RS, et al. Vitamin D, calcium supplementation, and colorectal adenomas: results of a randomized trial. *J Natl Cancer Inst* 2003;95:1765–71.
 43. Peters U, McGlynn KA, Chatterjee N, et al. Vitamin D, calcium, and vitamin D receptor polymorphism in colorectal adenomas. *Cancer Epidemiol Biomarkers Prev* 2001;10:1267–74.
 44. Wu K, Willett WC, Fuchs CS, Colditz GA, Giovannucci EL. Calcium intake and risk of colon cancer in women and men. *J Natl Cancer Inst* 2002;94:437–46.
 45. Zheng W, Anderson KE, Kushi LH, et al. A prospective cohort study of intake of calcium, vitamin D, other micronutrients in relation to incidence of rectal cancer among postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 1998;7:221–5.
 46. Fedirko V, Bostick RM, Flanders WD, et al. Effects of vitamin D and calcium on proliferation and differentiation in normal colon mucosa: a randomized clinical trial. *Cancer Epidemiol Biomarkers Prev* 2009;18:2933–41.
 47. Quarles LD. Endocrine functions of bone in mineral metabolism regulation. *J Clin Invest* 2008;118:3820–8.

48. Pence BC, Buddingh F. Inhibition of dietary fat-promoted colon carcinogenesis in rats by supplemental calcium or vitamin D₃. *Carcinogenesis* 1988;9:187–90.
49. Sitrin MD, Halline AG, Abrahams C, Brasitus TA. Dietary calcium and vitamin D modulate 1,2-dimethylhydrazine-induced colonic carcinogenesis in the rat. *Cancer Res* 1991;51:5608–13.
50. Beaty MM, Lee EY, Glauert HP. Influence of dietary calcium and vitamin D on colon epithelial cell proliferation and 1,2-dimethylhydrazine-induced colon carcinogenesis in rats fed high fat diets. *J Nutr* 1993;123:144–52.
51. Chlebowski RT, Wactawski-Wende J, Ritenbaugh C, et al. Estrogen plus progestin and colorectal cancer in postmenopausal women. *N Engl J Med* 2004;350:991–1004.
52. Anderson GL, Limacher M, Assaf AR, et al. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA* 2004; 291:1701–12.
53. Protiva P, Cross HS, Hopkins ME, et al. Chemoprevention of colorectal neoplasia by estrogen: potential role of vitamin D activity. *Cancer Prev Res (Phila Pa)* 2009;2:43–51.
54. Harris ED. Regulation of antioxidant enzymes. *FASEB J* 1992;6: 2675–83.

Effects of Supplemental Vitamin D and Calcium on Oxidative DNA Damage Marker in Normal Colorectal Mucosa: A Randomized Clinical Trial

Veronika Fedirko, Roberd M. Bostick, Qi Long, et al.

Cancer Epidemiol Biomarkers Prev 2010;19:280-291.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/19/1/280>

Cited articles This article cites 51 articles, 11 of which you can access for free at:
<http://cebp.aacrjournals.org/content/19/1/280.full#ref-list-1>

Citing articles This article has been cited by 18 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/19/1/280.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/19/1/280>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.