
Correction

Correction: Effect of Differing Levels of Tobacco-Specific Nitrosamines in Cigarette Smoke on the Levels of Biomarkers in Smokers

In this article (1), which was published in the June 2010 issue of *Cancer Epidemiology, Biomarkers & Prevention*, in Fig. 1, the x-axis is labeled with units of μg . The correct units, as given in the figure legend, are ng.

Reference

1. Ashley DL, O'Connor RJ, Bernert JT, Watson CH, Polzin GM, Jain RB, Hammond D, Hatsukami DK, Giovino GA, Cummings KM, McNeill A, Shahab L, King B, Fong GT, Zhang L, Xia Y, Yan X, McCraw JM. Effect of differing levels of tobacco-specific nitrosamines in cigarette smoke on the levels of biomarkers in smokers. *Cancer Epidemiol Biomarkers Prev* 2010;19:1389–98.

©2010 American Association for Cancer Research.
doi: 10.1158/1055-9965.EPI-10-0731

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Correction: Effect of Differing Levels of Tobacco-Specific Nitrosamines in Cigarette Smoke on the Levels of Biomarkers in Smokers

Cancer Epidemiol Biomarkers Prev 2010;19:2416.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/19/9/2416.1>

Cited articles This article cites 1 articles, 1 of which you can access for free at:
<http://cebp.aacrjournals.org/content/19/9/2416.1.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/19/9/2416.1>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.