

Trends in Colorectal Cancer Test Use among Vulnerable Populations in the United States

Carrie N. Klabunde¹, Kathleen A. Cronin², Nancy Breen¹, William R. Waldron³, Anita H. Ambis¹, and Marion R. Nadel⁴

Abstract

Background: Evaluating trends in colorectal cancer (CRC) screening use is critical for understanding screening implementation, and whether population groups targeted for screening are receiving it, consistent with guidelines. This study examines recent national trends in CRC test use, including among vulnerable populations.

Methods: We used the 2000, 2003, 2005, and 2008 National Health Interview Survey to examine national trends in CRC screening use overall and for fecal occult blood test (FOBT), sigmoidoscopy, and colonoscopy. We also assessed trends by race/ethnicity, educational attainment, income, time in the United States, and access to health care.

Results: During 2000 to 2008, significant declines in FOBT and sigmoidoscopy use and significant increases in colonoscopy use and in the percentages of adults up-to-date with CRC screening occurred overall and for most population subgroups. Subgroups with consistently lower rates of colonoscopy use and being up-to-date included Hispanics; people with minimal education, low income, or no health insurance; recent immigrants; and those with no usual source of care or physician visits in the past year. Among up-to-date adults, there were few subgroup differences in the type of test by which they were up-to-date (i.e., FOBT, sigmoidoscopy, or colonoscopy).

Conclusions: Although use of CRC screening and colonoscopy increased among U.S. adults, including those from vulnerable populations, 45% of adults aged 50 to 75—or nearly 35 million people—were not up-to-date with screening in 2008.

Impact: Continued monitoring of CRC screening rates among population subgroups with consistently low utilization is imperative. Improvement in CRC screening rates among all population groups in the United States is still needed. *Cancer Epidemiol Biomarkers Prev*; 20(8); 1611–21. ©2011 AACR.

Introduction

Since the mid-1990s, multiple expert groups in the United States have recommended screening asymptomatic average-risk adults for colorectal cancer (CRC; refs. 1–3). In its most recent guidance published in 2008, the U.S. Preventive Services Task Force (USPSTF) strongly recommends screening average-risk adults aged 50 to 75 for CRC with annual high-sensitivity fecal occult blood

test (FOBT), sigmoidoscopy every 5 years, or colonoscopy every 10 years (4). Although CRC screening rates in the United States have increased from 20% to 30% in 1997 to nearly 55% in 2008, millions of eligible adults are not screened (5). Moreover, certain population subgroups—including Hispanics, people with low income or low educational attainment, recent immigrants, and those without health insurance, a usual source of care, or recent physician contact—have shown especially low CRC screening rates (6–10). For purposes of this report, we have defined these 7 subgroups as vulnerable populations.

Evaluating trends in the use of CRC screening is important for understanding how screening is being implemented in practice in the United States; to assess whether the population groups targeted for screening are receiving it, consistent with guidelines; and to identify potential problems with underuse, overuse, and misuse of screening (11). Although some prior studies have examined trends in use of CRC screening, their focus has been on the Medicare population, or they have not included more contemporaneous data or comprehensively assessed trends among vulnerable populations

Authors' Affiliations: ¹Applied Research Program and ²Surveillance Research Program, Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda; ³Information Management Services, Inc., Silver Spring, Maryland; ⁴Division of Cancer Prevention and Control, Centers for Disease Control and Prevention, Atlanta, Georgia

Note: This study was presented in part at the NIH State of the Science Conference: Enhancing Use and Quality of Colorectal Cancer Screening in Bethesda, MD, on February 2, 2010.

Corresponding Author: Carrie Klabunde, Health Services and Economics Branch, Applied Research Program, National Cancer Institute, 6130 Executive Boulevard, Room 4005, Bethesda, MD 20892-7344. Phone: 301-402-3362; Fax: 301-435-3710; E-mail: klabundC@mail.nih.gov

doi: 10.1158/1055-9965.EPI-11-0220

©2011 American Association for Cancer Research.

(12–18). Following implementation by the Medicare program in 2001 of coverage for CRC screening with colonoscopy for average-risk enrollees, colonoscopy use has increased rapidly (5); of particular interest but relatively unexplored is whether vulnerable populations use colonoscopy at rates comparable with non-vulnerable groups. In the present study, we aim to (i) describe recent national trends in CRC test use overall and among 7 population subgroups that historically have shown lower rates of cancer screening; (ii) assess whether vulnerable subgroups are less likely to receive colonoscopy than their less vulnerable counterparts; and (iii) provide baseline data for assessing trends in CRC test use following implementation of health care reform measures.

Methods

Data source

We used data from the 2000, 2003, 2005, and 2008 National Health Interview Survey (NHIS) to examine national trends in the overall use of CRC screening and in use of FOBT, sigmoidoscopy, and colonoscopy. The NHIS is a multipurpose health survey sponsored by the National Center for Health Statistics, Centers for Disease Control and Prevention. Conducted annually since 1957, the NHIS is the principal source of information on the health of the civilian, noninstitutionalized household population of the United States. The majority of NHIS interviews are conducted in person by trained personnel from the U.S. Census Bureau. Response rates for the NHIS survey years included in this analysis were 72% (2000), 74% (2003), 69% (2005), and 63% (2008); more information about response rates is available in the appendices for each year's survey description (19–22). Survey responses were weighted to reflect the probability of selection into the sample and survey nonresponse.

Men and women aged 40 or older were asked whether they ever had a sigmoidoscopy, colonoscopy, or proctoscopy, and when they had their most recent examination. They also were asked whether they ever had a blood stool test using a home test kit, and when they had their most recent test. Brief descriptions of each test were provided. Respondents were also asked to report their age, race/ethnicity, educational attainment, household income, time in the United States, type of health care coverage, whether there was a place that they usually went when they were sick or needed advice about their health, and the number of times they saw a physician in the past year. Complete survey instrumentation for the NHIS is available from the National Center for Health Statistics (23).

Measures

CRC test use. Consistent with current USPSTF recommendations (4), we defined CRC test use as having had a home FOBT in the past year, sigmoidoscopy in the

past 5 years, colonoscopy in the past 10 years, or any of these (up-to-date with CRC screening). We included in this definition tests done for any reason, not just as part of a routine exam, because the reported reason for having the test may not be accurate, and having the test within the recommended time interval, regardless of the reason, essentially means that the individual has been screened.

Race/ethnicity. We used the racial/ethnic designations approved by the federal Office of Management and Budget (24) to categorize respondents as non-Hispanic white, non-Hispanic black, Hispanic, non-Hispanic Asian, and other race/ethnicity.

Education. Educational attainment was based on the highest level of education achieved by the respondent and categorized as more than high school (HS), HS graduate, and not attaining an HS diploma.

Income. Annual income was recorded for each family and reported as a percentage of the federal poverty level (FPL). Families with a computed FPL at or below 100% are considered impoverished. We categorized respondents according to the FPL of their interview year, as follows: <200% FPL, 200% to <300% FPL, 300% to <400% FPL, 400% to <500% FPL, and \geq 500% FPL and higher. Multiple imputation was used to impute missing data for NHIS income questions (25).

Time in the United States. Time in the United States was measured as born in the United States, immigrated to the United States 10 or more years ago, and immigrated to the United States less than 10 years ago.

Health insurance. We developed separate categories for type of health insurance for respondents aged 50 to 64 and 65 to 75. For the 50 to 64 subgroup, respondents were categorized as having private Health Maintenance Organization (HMO), private non-HMO, public (i.e., Medicaid, Medicare, and other government-sponsored insurance), or no insurance. For the 65 to 75 subgroup, respondents were categorized as having Medicare HMO; Medicare plus private supplemental insurance; Medicare with no supplemental insurance; Medicaid, military, or other government-sponsored insurance; or no insurance or Medicare Part A only.

Usual source of care. Respondents who indicated that they had a place where they usually went when they were sick or needed advice about their health were defined as having a usual source of care. Those who either reported that they did not have a place or the place they went to most often was an emergency room (ER) were defined as having no usual source of care.

Physician visits. We used 3 categories to measure the respondent's number of physician visits in the past year: none, 1, and 2 or more.

Data analysis

We used descriptive statistics to characterize overall use of CRC tests and use by specific test modality. Consistent with current USPSTF recommendations (4), we restricted the analysis to adults who were 50 to 75 years of age at the time of the interview. To

approximate an average-risk screening population, we excluded individuals who reported a personal history of CRC ($n = 320$). Respondents who reported being up-to-date for multiple tests were counted as receiving the test when calculating percentages for each test option reported. All percentages reported in the tables and figure were standardized to the 2000 U.S. population by 5-year age groups. The statistical significance of overall trends and each subgroup category trend was tested using logistic regression modeling; separate models were estimated for each test modality and subgroup category. The models included respondent age in 5-year categories to control for changes in the age structure of the U.S. population over time, and survey year for assessment of the statistical significance of the trend. Wald chi-square tests for association of independent variables with each dependent variable (i.e., had home FOBT in the past year versus did not; had sigmoidoscopy in the past 5 years versus did not; had colonoscopy in the past 10 years versus did not; and up-to-date with CRC screening versus not) were computed. Statistical tests were 2-sided, and a P value of 0.05 was considered to be statistically significant.

We conducted a subsidiary analysis among respondents who were up-to-date with CRC screening to assess whether there were subgroup differences in the type of test by which they were up-to-date (i.e., FOBT only, sigmoidoscopy only, colonoscopy only, or up-to-date by multiple-test modalities). We used the 2000 and 2008 NHIS data to calculate percentages and 95% CIs for each type of test and compared them across the 2 time periods to assess trends in test use patterns.

Survey weights were applied in the analyses to reflect the probability of selection into the sample and adjust for nonresponse and poststratification; the weighted data yield national estimates of the prevalence of CRC test use overall, by test modality, and by population subgroup. SUDAAN version 9.0 was used to analyze the data; this statistical package accounts for the complex design of the NHIS and allows for calculation of accurate standard errors, from which 95% CIs around point estimates were derived.

Results

Sample and U.S. population estimates for adults aged 50 to 75

Sample size and U.S. population estimates for each year and by population subgroup are shown in Table 1. The U.S. civilian and noninstitutionalized population aged 50 to 75 grew from an estimated 60 million people in 2000 to nearly 77 million people in 2008. The population subgroups shown in the table also increased in size over this time period, except for people with less than an HS education, with private HMO or Medicare HMO health insurance, or who were aged 65 to 75 and uninsured or had only Medicare Part A coverage. The number of people aged 65 to 75 with Medicare plus private

supplemental coverage remained essentially the same over the period 2000 to 2008.

Trends in home FOBT use

Use of home FOBT among U.S. adults aged 50 to 75 declined during 2000 to 2008 by 6.5 percentage points; in 2008, 11% of adults reported that they had completed a home FOBT in the past year (Table 2). Statistically significant declines ranging from 3.6 to 13.6 percentage points occurred for non-Hispanic whites and non-Hispanic blacks; all education and income groups; people born in the United States; people with private health insurance or Medicare coverage; those reporting a usual source of health care; and those who had seen a physician at least once in the past year.

Trends in sigmoidoscopy use

Similar trends were observed for sigmoidoscopy (Table 3). The proportion of U.S. adults aged 50 to 75 who reported having a sigmoidoscopy in the past 5 years declined by 7 percentage points during 2000 to 2008, to 2.4%. Statistically significant declines ranging from 1.9 to 10.9 percentage points occurred for all population subgroups, with the exception of non-Hispanic Asians, immigrants who had been in the United States less than 10 years, the uninsured, and people without a usual source of health care.

Trends in colonoscopy use

In contrast to the trends for home FOBT and sigmoidoscopy, use of colonoscopy increased markedly among U.S. adults aged 50 to 75: in 2000, 19.0% reported having a colonoscopy in the past 10 years, whereas in 2008, 47.5% indicated that they had undergone this procedure (Table 4). Statistically significant increases in colonoscopy use over the period 2000 to 2008 were seen for all population subgroups, with 1 exception: people aged 65 to 75 who were uninsured or reported having only Medicare Part A coverage. The sample size for this group, however, was small ($n < 90$).

Although colonoscopy use increased significantly among nearly all population subgroups, there were differences in use rates (Table 4). The percentages of Hispanics reporting colonoscopy use were consistently lower than those of non-Hispanic whites, and the percentage point increase in use among Hispanics was less than that of non-Hispanic whites. Similar patterns were observed for people with less than an HS education (versus those with more than an HS education); recent immigrants (versus individuals born in the United States); people with family income less than 200% of FPL (versus those with family income $\geq 500\%$ of FPL); the uninsured (versus people with health insurance coverage); those with no usual source of care (versus individuals who reported having a usual source of care); and people who had no physician visits in the past year (versus those with 2 or more visits).

Table 1. Sample size and U.S. population estimates^a for adults aged 50 to 75, NHIS 2000–2008

	2000		2003		2005		2008	
	<i>n</i>	<i>N</i>	<i>n</i>	<i>N</i>	<i>n</i>	<i>N</i>	<i>n</i>	<i>N</i>
Overall	10,041	60,120,556	9,928	66,367,948	10,733	70,151,788	7,776	76,763,249
Race/ethnicity								
NH white	7,233	47,913,678	7,139	52,543,643	7,639	54,401,706	5,236	57,922,863
NH black	1,305	5,569,419	1,258	6,391,692	1,429	6,957,214	1,165	7,957,136
Hispanic	1,186	4,399,116	1,218	5,151,211	1,237	5,654,680	911	6,728,265
NH Asian	205	1,547,048	202	1,589,687	283	2,210,807	329	2,949,205
Education								
>HS	4,392	27,615,842	4,734	33,165,278	5,442	36,643,584	4,127	42,405,243
HS graduate	3,088	19,204,541	2,995	20,652,409	3,232	21,813,778	2,259	22,310,729
<HS	2,433	12,514,502	2,026	11,374,206	1,930	10,849,488	1,332	11,400,937
Family income (% FPL)								
500%+	2,798	19,670,541	2,754	21,676,932	3,188	24,234,310	2,389	27,674,983
400%–<500%	992	6,411,053	1,014	7,198,802	1,047	7,330,034	761	8,130,622
300%–<400%	1,259	7,901,910	1,301	8,962,111	1,463	9,929,025	1,057	10,825,327
200%–<300%	1,747	10,360,448	1,698	11,189,666	1,727	10,961,088	1,219	11,510,251
<200%	3,245	15,776,604	3,162	17,340,437	3,309	17,697,331	2,349	18,622,066
Time in the United States								
Born in United States	8,748	53,500,959	8,579	58,696,863	9,247	61,359,559	6,588	66,551,312
Immigrant, in United States 10+ years	1,031	5,170,983	1,164	6,505,103	1,300	7,605,197	1,073	9,145,404
Immigrant, in United States <10 years	134	776,187	137	892,854	142	928,520	90	835,973
Health insurance—ages 50–64								
Private non-HMO	2,862	19,663,349	3,192	24,057,110	3,627	26,505,422	2,643	29,692,875
Private HMO	1,910	12,199,153	1,648	11,612,087	1,657	11,652,507	992	10,182,499
Public	813	3,947,810	975	5,429,748	1,046	5,734,111	915	8,149,749
Uninsured	886	4,627,743	940	5,486,960	1,041	6,126,622	718	6,930,625
Health insurance—ages 65–75								
Medicare + private	1,944	11,631,560	1,693	11,517,482	1,700	10,864,474	1,195	11,513,029
Medicare HMO	597	3,128,678	351	1,978,735	396	2,263,677	329	2,572,604
Medicare, no supplemental	516	2,614,259	540	3,190,164	693	4,032,079	559	4,565,195
Medicaid, military, and other government	386	1,607,414	482	2,499,011	478	2,456,428	367	2,649,194
Uninsured or Medicare Part A only	87	458,553	69	359,875	62	365,189	47	345,054
Has usual source of care								
Yes (excluding ER)	9,137	55,174,724	9,038	61,025,909	9,718	64,039,052	7,008	69,503,281
No	838	4,622,595	814	4,850,947	907	5,463,296	661	6,266,798
# Physician visits in past year								
2 or more	7,249	43,606,924	7,272	48,865,333	7,788	51,105,829	5,669	55,837,224
1	1,334	8,357,917	1,236	8,586,391	1,478	9,896,285	1,042	10,766,323
None	1,296	7,276,414	1,236	7,770,913	1,246	7,869,436	888	8,583,344

NOTE: *n* = sample size; *N* = U.S. population estimate.^aFor the U.S. civilian and noninstitutionalized population.**Trends in being up-to-date with CRC screening**

The proportion of the U.S. population aged 50 to 75 that is up-to-date with CRC screening increased by nearly 16 percentage points over the period 2000 to 2008 (Table 5). In 2000, 38.6% were up-to-date; by 2008, nearly 55% reported being up-to-date. Statistically significant increases in the proportion up-to-date ranging from 1.8 to 28.3 percentage points occurred for all population

subgroups, with the exception of people aged 65 to 75 who were uninsured or reported having only Medicare Part A coverage.

Although the percentages of people up-to-date with CRC screening increased significantly among nearly all population subgroups, there were subgroup differences (Table 5). The percentages of up-to-date Hispanics were consistently lower than those of non-Hispanic whites,

Table 2. Reported home FOBT in the past year, U.S. adults aged 50 to 75, NHIS 2000–2008^a

	2000 % (95% CI)	2003 % (95% CI)	2005 % (95% CI)	2008 % (95% CI)	Percentage point change, 2000–2008
Overall	17.4 (16.4–18.3)	14.8 (13.8–15.9)	12.7 (11.8–13.6)	10.9 (10.0–12.0)	–6.5
Race/ethnicity					
NH white	18.3 (17.2–19.4)	15.0 (13.9–16.2)	13.4 (12.4–14.5)	11.2 (10.0–12.4)	–7.1
NH black	14.7 (12.0–17.8)	15.2 (12.7–18.1)	10.0 (8.0–12.4)	10.3 (8.0–13.2)	–4.4
Hispanic	9.8 (7.5–12.5)	11.2 (8.2–15.0)	8.6 (6.4–11.5)	8.9 (6.5–12.0)	–0.9 ^b
NH Asian	15.8 (8.2–28.4)	11.3 (7.3–17.1)	14.8 (8.3–25.2)	12.2 (9.1–16.2)	–3.6 ^b
Education					
>HS	21.0 (19.4–22.6)	16.9 (15.4–18.5)	14.9 (13.6–16.4)	12.7 (11.1–14.5)	–8.3
HS graduate	16.2 (14.7–17.9)	14.2 (12.6–15.9)	11.7 (10.5–13.1)	9.7 (8.2–11.5)	–6.5
<HS	11.6 (10.1–13.4)	10.8 (9.1–12.8)	9.7 (8.0–11.9)	7.5 (6.0–9.3)	–4.1
Time in the United States					
Born in United States	18.0 (17.0–19.1)	15.4 (14.3–16.5)	13.1 (12.1–14.2)	11.1 (10.0–12.3)	–6.9
Immigrant, in United States 10+ years	13.9 (11.0–17.3)	11.1 (8.7–14.0)	10.4 (8.1–13.4)	10.3 (8.0–13.1)	–3.6 ^b
Immigrant, in United States <10 years	1.1 (0.2–5.2)	5.8 (2.7–12.1)	2.9 (1.1–7.4)	4.8 (1.7–12.7)	+3.7 ^b
Family income (% FPL)					
500%+	22.4 (20.0–25.1)	18.5 (16.0–21.2)	14.9 (13.0–17.0)	14.0 (11.0–17.6)	–8.4
400%–<500%	18.5 (14.7–23.0)	15.2 (12.4–18.4)	10.7 (8.1–14.1)	9.9 (6.9–14.0)	–8.6
300%–<400%	16.9 (14.0–20.2)	15.8 (13.1–19.1)	13.8 (11.2–17.0)	10.8 (8.5–13.6)	–6.1
200%–<300%	17.1 (14.7–19.7)	12.9 (10.7–15.6)	14.0 (11.5–16.9)	9.9 (7.7–12.7)	–7.2
<200%	12.8 (11.2–14.6)	11.3 (9.7–13.0)	10.2 (8.7–11.9)	9.1 (7.7–10.9)	–3.6
Health insurance—ages 50–64					
Private non-HMO	17.0 (15.5–18.6)	14.4 (12.9–16.0)	10.9 (9.7–12.3)	9.1 (7.8–10.5)	–7.9
Private HMO	16.0 (14.1–18.0)	14.7 (12.7–17.0)	13.4 (11.6–15.3)	11.6 (9.5–14.1)	–4.4
Public	13.8 (10.9–17.2)	15.9 (13.2–19.0)	14.0 (11.5–16.9)	10.8 (8.7–13.4)	–3.0 ^b
Uninsured	7.0 (5.2–9.5)	4.7 (3.5–6.4)	4.0 (2.9–5.5)	5.3 (3.4–8.0)	–1.7 ^b
Health insurance—ages 65–75					
Medicare + private	21.1 (18.7–23.7)	16.8 (14.4–19.6)	15.6 (13.3–18.1)	15.7 (12.4–19.7)	–5.4
Medicare HMO	29.6 (24.6–35.2)	22.7 (16.8–29.8)	17.5 (12.3–24.4)	15.9 (10.7–22.8)	–13.6
Medicare, no supplemental	13.3 (10.0–17.7)	15.4 (11.6–20.1)	14.5 (11.1–18.8)	8.2 (5.4–12.1)	–5.1
Medicaid, military, and other government	11.3 (6.7–18.3)	15.4 (12.2–19.2)	15.8 (11.3–21.6)	11.4 (7.7–16.6)	+0.1 ^b
Uninsured or Medicare Part A only	28.1 (15.9–44.7)	6.6 (2.9–14.5)	4.2 (1.5–11.3)	12.9 (5.8–26.4)	–15.2 ^b
Has usual source of care					
Yes (excluding ER)	18.3 (17.3–19.3)	15.6 (14.6–16.7)	13.4 (12.5–14.3)	11.5 (10.5–12.6)	–6.8
No	5.8 (3.7–9.1)	3.5 (2.2–5.6)	4.5 (2.2–9.0)	2.6 (1.4–4.6)	–3.2 ^b
# Physician visits in past year					
2 or more	20.1 (19.0–21.3)	17.3 (16.1–18.5)	14.8 (13.8–15.9)	12.2 (11.1–13.5)	–7.9
1	15.9 (13.4–18.7)	11.6 (9.5–14.1)	8.9 (7.2–10.9)	9.9 (7.7–12.7)	–6.0
None	2.2 (1.4–3.4)	2.0 (1.3–3.3)	2.7 (1.6–4.5)	2.1 (1.3–3.4)	–0.1 ^b

^aOverall and subgroup trends are statistically significant at $\alpha = 0.05$, unless otherwise noted.

^bTrend for this subgroup is not statistically significant at $\alpha = 0.05$.

and the percentage point increase in being up-to-date was lower for Hispanics than non-Hispanic whites. Similar patterns were evident for people with less than an HS education (versus those with more than an HS education); people with family income less than 200% of FPL

(versus those with family income $\geq 500\%$ of FPL); individuals aged 50 to 64 who are uninsured (versus people with health insurance coverage); those with no usual source of care (versus individuals who reported having a usual source of care); and people who had no physician

Table 3. Reported sigmoidoscopy in the past 5 years, United States adults aged 50 to 75, NHIS 2000–2008^a

	2000 % (95% CI)	2003 % (95% CI)	2005 % (95% CI)	2008 % (95% CI)	Percentage point change, 2000–2008
Overall	9.4 (8.7–10.2)	6.6 (5.9–7.4)	4.1 (3.6–4.7)	2.4 (1.9–3.0)	–7.0
Race/ethnicity					
NH white	10.1 (9.3–11.0)	7.1 (6.3–8.0)	4.3 (3.7–5.0)	2.4 (1.8–3.1)	–7.7
NH black	5.7 (4.3–7.5)	5.7 (4.0–8.0)	3.3 (2.1–5.1)	1.4 (0.8–2.5)	–4.3
Hispanic	5.5 (4.1–7.3)	3.6 (2.4–5.5)	2.73 (1.8–4.2)	2.9 (1.5–5.3)	–2.6
NH Asian	8.7 (3.9–18.4)	6.8 (3.2–13.7)	3.3 (1.5–7.0)	3.1 (1.6–6.1)	–5.6 ^b
Education					
>HS	12.4 (11.0–13.9)	9.1 (8.0–10.3)	5.5 (4.6–6.6)	2.8 (2.2–3.7)	–9.6
HS graduate	8.5 (7.2–10.0)	5.7 (4.4–7.3)	2.5 (1.9–3.3)	1.7 (1.2–2.6)	–6.8
<HS	4.8 (3.8–6.1)	2.2 (1.6–3.1)	3.6 (2.6–4.9)	1.7 (0.7–4.1)	–3.1
Time in the United States					
Born in United States	9.9 (9.1–10.7)	7.1 (6.3–7.9)	4.3 (3.7–4.9)	2.5 (1.9–3.2)	–7.4
Immigrant, in United States 10+ years	6.2 (4.2–8.9)	3.4 (2.4–4.9)	3.0 (2.1–4.3)	1.8 (0.9–3.4)	–4.4
Immigrant, in United States <10 years	1.2 (0.3–5.6)	1.4 (0.4–5.4)	1.9 (0.4–8.6)	0.0 (0.0–0.0)	–1.2 ^b
Family income (% FPL)					
500%+	14.5 (12.2–17.2)	10.4 (8.5–12.7)	6.2 (5.0–7.8)	3.6 (2.5–5.2)	–10.9
400%–<500%	8.5 (6.5–11.0)	6.1 (4.3–8.7)	3.7 (1.9–7.0)	3.2 (1.7–5.8)	–5.3
300%–<400%	9.5 (7.14–12.6)	6.6 (4.8–9.1)	3.6 (2.4–5.4)	1.6 (0.8–3.2)	–7.9
200%–<300%	7.9 (6.3–9.8)	4.9 (3.6–6.7)	2.9 (1.9–4.4)	1.5 (0.4–5.3)	–6.4
<200%	5.8 (4.7–7.0)	3.7 (2.7–5.0)	2.9 (2.1–3.8)	1.6 (0.7–3.5)	–4.2
Health insurance—ages 50–64					
Private non-HMO	9.4 (8.2–10.7)	6.9 (5.9–7.9)	3.3 (2.7–4.0)	2.0 (1.4–2.8)	–7.4
Private HMO	9.8 (8.3–11.4)	7.4 (6.0–9.1)	6.9 (5.5–8.7)	3.5 (2.4–5.2)	–6.3
Public	4.8 (3.1–7.4)	3.8 (2.6–5.6)	3.0 (1.9–4.7)	1.4 (0.7–3.0)	–3.4
Uninsured	2.2 (1.3–3.8)	1.7 (1.0–3.0)	1.2 (0.5–3.0)	1.5 (0.6–3.6)	–0.7 ^b
Health insurance—ages 65–75					
Medicare + private	12.6 (10.4–15.2)	7.6 (5.9–9.9)	5.0 (3.5–7.1)	2.0 (1.1–3.9)	–10.6
Medicare HMO	10.5 (7.9–13.7)	15.7 (11.0–21.8)	4.6 (3.1–6.8)	5.1 (2.6–9.7)	–5.4
Medicare, no supplemental	8.4 (5.4–13.0)	3.6 (1.8–7.0)	2.8 (1.3–5.7)	3.7 (1.4–9.5)	–4.7
Medicaid, military, and other government	9.1 (4.9–16.5)	5.2 (3.4–7.8)	5.9 (3.6–9.5)	2.6 (1.0–6.7)	–6.5
Uninsured or Medicare Part A only	8.7 (4.5–16.2)	20.1 (7.6–43.5)	2.4 (0.6–8.9)	0.9 (0.1–6.3)	–7.8 ^b
Has usual source of care					
Yes (excluding ER)	10.0 (9.2–10.8)	7.0 (6.3–7.8)	4.4 (3.8–5.0)	2.4 (2.0–3.1)	–7.6
No	2.4 (1.4–4.3)	1.3 (0.6–2.6)	0.8 (0.4–1.7)	1.1 (0.4–3.0)	–1.3 ^b
# Physician visits in past year					
2 or more	10.5 (9.6–11.4)	7.4 (6.6–8.3)	4.6 (4.0–5.4)	2.4 (1.8–3.1)	–8.1
1	8.6 (6.8–10.9)	5.0 (3.7–6.6)	3.5 (2.5–5.0)	3.1 (1.9–4.9)	–5.6
None	3.3 (2.3–4.7)	2.7 (1.8–4.0)	1.1 (0.6–1.9)	1.4 (0.8–2.5)	–1.9

^aOverall and subgroup trends are statistically significant at $\alpha = 0.05$, unless otherwise noted.^bTrend for this subgroup is not statistically significant at $\alpha = 0.05$.

visits in the past year (versus those with 1 or more visits). The percentages of recent immigrants who were up-to-date were consistently lower than those of individuals born in the United States, as were the

percentages of individuals aged 65 to 75 who have Medicare but no supplemental coverage compared with those covered by Medicare plus private supplemental insurance.

Table 4. Reported colonoscopy in the past 10 years, U.S. adults aged 50 to 75, NHIS 2000–2008^a

	2000 % (95% CI)	2003 % (95% CI)	2005 % (95% CI)	2008 % (95% CI)	Percentage point change, 2000–2008
Overall	19.0 (18.0–20.1)	29.9 (28.6–31.2)	39.4 (38.0–40.9)	47.5 (45.9–49.0)	+28.5
Race/ethnicity					
NH white	19.8 (18.6–21.0)	31.3 (29.9–32.8)	42.3 (40.7–43.9)	50.0 (48.2–51.9)	+30.2
NH black	18.3 (15.3–21.7)	24.7 (21.7–27.9)	30.3 (27.2–33.5)	45.4 (41.0–49.9)	+27.1
Hispanic	13.9 (11.0–17.5)	21.3 (17.5–25.7)	25.3 (20.4–30.8)	31.7 (27.4–36.3)	+17.8
NH Asian	12.5 (6.9–21.5)	23.3 (16.1–32.3)	25.3 (17.4–35.3)	41.7 (34.8–48.9)	+29.2
Education					
>HS	22.4 (20.7–24.2)	34.6 (32.7–36.6)	44.5 (42.7–46.4)	54.4 (52.1–56.8)	+32.0
HS graduate	18.7 (16.9–20.6)	29.5 (27.4–31.7)	37.7 (35.3–40.2)	44.1 (41.4–46.8)	+25.4
<HS	12.9 (11.2–14.7)	19.6 (17.4–22.1)	27.7 (25.1–30.6)	31.3 (28.2–34.6)	+18.4
Time in the United States					
Born in United States	19.6 (18.5–20.8)	30.5 (29.1–31.9)	40.7 (39.2–42.3)	49.2 (47.5–50.9)	+29.6
Immigrant, in United States 10+ years	15.4 (12.8–18.5)	26.9 (23.2–31.0)	31.1 (27.1–35.4)	36.8 (32.6–41.1)	+21.4
Immigrant, in United States <10 years	8.0 (4.7–13.2)	9.2 (4.6–17.7)	12.0 (6.6–20.9)	19.1 (11.9–29.2)	+11.1
Family income (% FPL)					
500%+	23.0 (20.4–25.8)	37.4 (34.4–40.5)	47.0 (44.1–50.0)	57.8 (54.2–61.3)	+34.8
400%–<500%	20.2 (15.7–25.6)	31.1 (26.2–36.5)	45.7 (39.6–51.8)	52.2 (46.8–57.5)	+32.0
300%–<400%	18.1 (15.0–21.6)	31.7 (28.3–35.3)	40.0 (36.1–44.1)	46.0 (42.3–49.9)	+27.9
200%–<300%	17.5 (14.8–20.6)	27.5 (24.6–30.5)	36.5 (33.2–40.0)	44.2 (39.3–49.2)	+26.7
<200%	15.8 (14.0–17.8)	22.0 (20.0–24.2)	28.8 (26.6–31.2)	33.9 (31.1–36.8)	+18.1
Health insurance—ages 50–64					
Private non-HMO	17.4 (16.0–19.0)	28.2 (26.4–30.3)	36.7 (34.8–38.5)	49.2 (46.9–51.5)	+31.8
Private HMO	16.9 (14.9–19.2)	29.0 (26.4–31.7)	36.2 (33.4–39.1)	47.5 (43.8–51.3)	+30.6
Public	17.2 (14.2–20.7)	24.3 (21.1–27.9)	32.1 (29.0–35.3)	39.0 (35.4–42.7)	+21.8
Uninsured	7.2 (5.4–9.5)	11.2 (8.8–14.1)	13.2 (10.9–15.9)	14.9 (11.7–18.9)	+7.7
Health insurance—ages 65–75					
Medicare + private	27.8 (25.0–30.9)	40.0 (36.6–43.6)	53.8 (49.9–57.6)	59.4 (55.0–63.6)	+31.6
Medicare HMO	23.6 (18.9–29.1)	35.0 (28.2–42.4)	43.7 (35.9–51.9)	50.7 (43.2–57.7)	+27.1
Medicare, no supplemental	13.8 (10.1–18.4)	26.8 (21.3–33.1)	45.0 (39.8–50.3)	45.8 (39.4–52.4)	+32.0
Medicaid, military, and other government	13.3 (9.7–18.0)	34.3 (27.0–42.4)	45.5 (37.6–53.6)	49.6 (42.1–57.1)	+36.3
Uninsured or Medicare Part A only	13.7 (6.5–26.7)	11.9 (6.4–21.1)	34.9 (29.9–40.3)	42.6 (29.7–56.7)	+28.9 ^b
Has usual source of care					
Yes (excluding ER)	19.9 (18.8–21.0)	31.2 (29.9–32.5)	41.1 (39.7–42.6)	50.0 (48.3–51.6)	+30.1
No	7.6 (5.4–10.5)	12.1 (7.7–18.4)	18.7 (14.4–24.0)	14.3 (11.4–17.8)	+6.7
# Physician visits in past year					
2 or more	22.3 (21.0–23.6)	34.3 (32.8–35.9)	44.4 (42.8–46.0)	54.2 (52.5–55.9)	+31.9
1	11.6 (9.5–14.1)	20.6 (17.1–24.5)	31.9 (27.9–36.1)	34.7 (31.2–38.3)	+23.1
None	5.8 (4.4–7.6)	12.2 (8.5–17.3)	15.4 (12.3–19.0)	15.8 (12.3–20.2)	+10.0

^aOverall and subgroup trends are statistically significant at $\alpha = 0.05$, unless otherwise noted.^bTrend for this subgroup is not statistically significant at $\alpha = 0.05$.

Among those up-to-date with CRC screening, there were few subgroup differences in the type of test they reported using (Fig. 1). In 2000 and in 2008, no significant differences were evident in the percentages of non-Hispanic whites, non-Hispanic blacks, Hispanics, and Asians

who were up-to-date by FOBT, sigmoidoscopy, or colonoscopy (Fig. 1A). By education subgroup, there were no significant differences in the percentages up-to-date by FOBT, sigmoidoscopy, or colonoscopy in 2000, although in 2008, people with less than an HS education were more

Table 5. Up-to-date¹ with colorectal cancer screening, U.S. adults aged 50 to 75, NHIS 2000–2008^a

	2000 % (95% CI)	2003 % (95% CI)	2005 % (95% CI)	2008 % (95% CI)	Percentage point change, 2000–2008
Overall	38.6 (37.4–39.9)	43.3 (41.9–44.7)	48.6 (47.2–50.1)	54.5 (52.9–56.2)	+15.9
Race/ethnicity					
NH white	40.5 (39.1–41.8)	45.3 (43.7–46.8)	51.7 (50.0–53.3)	57.0 (55.1–58.9)	+16.5
NH black	33.0 (29.6–36.7)	37.7 (34.0–41.6)	38.7 (35.3–42.3)	51.4 (47.1–55.6)	+18.4
Hispanic	26.4 (22.4–30.9)	30.3 (25.9–35.0)	32.4 (27.5–37.9)	39.1 (34.8–43.6)	+12.7
NH Asian	32.2 (25.8–39.3)	33.6 (25.6–42.5)	40.7 (34.4–47.4)	50.8 (43.2–58.4)	+18.6
Education					
>HS	45.3 (43.4–47.3)	50.2 (48.3–52.1)	55.3 (53.5–57.1)	62.0 (60.0–64.0)	+16.7
HS graduate	37.4 (35.1–39.8)	42.2 (39.9–44.4)	45.8 (43.4–48.2)	50.5 (47.7–53.2)	+13.1
<HS	26.0 (24.0–28.2)	28.4 (25.8–31.2)	35.3 (32.3–38.5)	37.5 (34.1–41.0)	+11.5
Time in the United States					
Born in United States	39.9 (38.5–41.2)	44.5 (43.1–46.0)	50.2 (48.6–51.8)	56.5 (54.7–58.3)	+16.6
Immigrant, in United States 10+ years	32.0 (28.2–35.9)	35.5 (31.6–39.6)	39.2 (35.3–43.3)	42.4 (38.5–46.7)	+10.4
Immigrant, in United States <10 years	10.0 (6.1–15.9)	16.0 (10.0–24.7)	16.6 (10.2–26.0)	25.7 (17.2–36.6)	+15.7
Family income (% FPL)					
500%+	49.1 (45.8–52.4)	55.1 (52.5–57.6)	58.3 (55.4–61.1)	66.0 (63.2–68.8)	+16.9
400%–<500%	39.8 (34.7–45.2)	43.1 (38.4–48.0)	53.7 (47.9–59.5)	60.3 (55.1–65.3)	+20.5
300%–<400%	37.5 (33.9–41.3)	45.7 (42.1–49.4)	48.5 (44.9–52.1)	53.0 (49.0–56.8)	+15.5
200%–<300%	36.2 (32.8–39.8)	38.3 (35.3–41.4)	45.5 (41.7–49.2)	50.9 (47.0–54.9)	+14.7
<200%	29.8 (27.6–32.1)	32.3 (30.0–34.6)	37.1 (34.7–39.5)	40.1 (37.2–43.1)	+10.3
Health insurance—ages 50–64					
Private non-HMO	36.9 (34.9–38.9)	41.8 (39.5–44.0)	45.2 (43.3–47.2)	55.2 (52.8–57.5)	+18.3
Private HMO	36.5 (34.1–39.1)	43.1 (40.2–46.0)	48.5 (45.5–51.5)	55.9 (52.1–59.7)	+19.4
Public	30.7 (26.8–34.8)	37.3 (33.5–41.2)	42.1 (38.6–45.7)	45.3 (41.4–49.2)	+14.6
Uninsured	14.7 (12.0–17.8)	16.2 (13.5–19.3)	17.2 (14.6–20.2)	19.9 (16.1–24.3)	+5.2
Health insurance—ages 65–75					
Medicare + private	51.4 (48.1–54.7)	54.6 (50.9–58.1)	64.0 (60.2–67.6)	67.5 (63.3–71.4)	+16.1
Medicare HMO	51.8 (45.6–57.9)	57.3 (49.5–64.7)	52.6 (44.8–60.4)	62.9 (54.8–70.3)	+11.1
Medicare, no supplemental	31.9 (26.1–38.3)	36.9 (30.8–43.4)	52.3 (47.0–57.6)	52.9 (46.7–59.0)	+21.0
Medicaid, military, and other government	28.5 (22.2–35.9)	49.0 (41.6–56.4)	56.8 (48.3–64.9)	56.8 (48.9–64.3)	+28.3
Uninsured or Medicare Part A only	41.3 (27.8–56.2)	36.3 (19.6–57.2)	39.3 (33.1–45.9)	51.3 (36.4–66.0)	+10.0 ^b
Has usual source of care					
Yes (excluding ER)	40.5 (39.2–41.8)	45.4 (44.0–46.8)	50.8 (49.3–52.3)	57.4 (55.7–59.0)	+16.9
No	14.5 (11.4–18.2)	14.9 (10.3–21.2)	21.7 (17.2–27.0)	16.3 (13.1–20.1)	+1.8
# Physician visits in past year					
2 or more	44.6 (43.1–46.1)	49.5 (47.9–51.2)	54.8 (53.1–56.4)	61.7 (59.9–63.4)	+17.1
1	29.2 (26.2–32.3)	31.4 (27.6–35.6)	40.1 (36.0–44.4)	43.3 (39.4–47.3)	+14.1
None	10.6 (8.7–12.8)	15.9 (11.9–21.0)	17.9 (14.7–21.6)	18.1 (14.5–22.4)	+7.5

¹Up-to-date is defined as having had a home FOBT in the past year, sigmoidoscopy in the past 5 years, and/or colonoscopy in the past 10 years.

^aOverall and subgroup trends are statistically significant at $\alpha = 0.05$, unless otherwise noted.

^bTrend for this subgroup is not statistically significant at $\alpha = 0.05$.

likely than those having more education to be up-to-date by FOBT only (Fig. 1B). For those aged 50 to 64, there were no significant differences in either year in the percentages up-to-date by FOBT, sigmoidoscopy, or

colonoscopy by type of health care coverage (Fig. 1C). Similarly, in 2000 and in 2008, there were no significant differences in the percentages of people with or without a usual source of health care who were up-to-date by




Figure 1. Type of colorectal cancer test used by U.S. adults aged 50 to 75 who met Colorectal Cancer Screening Guidelines.¹

FOBT, sigmoidoscopy, or colonoscopy (Fig. 1D). Among those up-to-date, for all population subgroups, the most notable trend was the substantial increase in colonoscopy use.

Discussion

This analysis of nationally representative data for adults aged 50 to 75, the age group targeted by the USPSTF as benefiting the most from CRC screening (4), showed that the proportion up-to-date with CRC screening recommendations increased from 39% in 2000 to 55% in 2008. Moreover, most CRC screening in the United States during 2000 to 2008 was undertaken with colonoscopy, whereas use of FOBT and sigmoidoscopy declined. Despite the 16 percentage point increase in the proportion of U.S. adults up-to-date with CRC screening, though, it is important to note that in 2008, 45% of the

population aged 50 to 75—or nearly 35 million people—were not current with screening as recommended by the USPSTF.

A key aim of our study was to examine recent national trends in CRC test use among 7 population groups that historically have had lower rates of cancer screening: Hispanics; people with low income or low educational attainment; recent immigrants; and those without health insurance, a usual source of health care, or recent physician contact. Over the period 2000 to 2008, we observed statistically significant increases in the proportions of people in these vulnerable populations who were up-to-date with CRC screening. The 1 exception was individuals aged 65 to 75 who were uninsured or reported having only Medicare Part A coverage. However, because of small sample sizes for this group, we may have lacked sufficient statistical power to detect a significant increase.

Nevertheless, it is important to highlight that there are still disparities in CRC screening rates. Among the 7 vulnerable populations examined in our analysis, all showed lower rates of colonoscopy use and in being up-to-date with CRC screening, compared with more advantaged populations. For example, in 2008, 39% of Hispanics were up-to-date with CRC screening, compared with 57% of non-Hispanic whites; 32% of Hispanics reported having colonoscopy in the past 10 years, compared with 50% of non-Hispanic whites. Other groups with CRC screening rates lower than 40% in 2008 included people with less than an HS education (38% up-to-date), immigrants who had been in the United States less than 10 years (26% up-to-date), the uninsured (20% up-to-date), people with no physician visits in the past year (18% up-to-date), and those with no usual source of care (16% up-to-date).

Our analysis further revealed that, even among vulnerable subgroups, most CRC screening during 2000 to 2008 was undertaken with colonoscopy, whereas FOBT and sigmoidoscopy use declined. Although concerns have been raised that individuals from vulnerable populations might not have access to colonoscopy equivalent to that of more advantaged individuals (26), we found no significant subgroup differences in test use patterns among individuals who were up-to-date with CRC screening. In other words, in 2008, colonoscopy use predominated among all groups, including racial/ethnic minorities, people with low educational attainment, the uninsured, and those lacking a usual source of care.

Our study has limitations. Data are self-reported; validation studies have shown that adults may over-report screening behaviors in surveys (27). Institutionalized and

noncivilian populations are not captured in the NHIS. American Indian, Alaska Native, and Pacific Islander samples were too small to analyze separately and therefore were excluded from the study.

Despite its limitations, our study provides important national benchmark data for monitoring progress toward improved delivery of CRC screening, particularly following the 2010 National Institutes of Health State of the Science Conference on Enhancing Use and Quality of CRC Screening (5, 28)—which outlined research and public health agendas for addressing underuse, overuse, and misuse of CRC screening—and implementation of national health care reform legislation, which includes provisions for reducing or removing barriers to preventive services, including CRC screening (29). Demonstration of improved rates of CRC screening among historically underserved groups will be a critical marker of success for public health efforts that target CRC screening. As our study documents, although CRC screening rates in the United States increased during 2000 to 2008, considerable improvement among all population groups is still needed.

Disclosure of Potential Conflicts of Interest

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the National Cancer Institute or the Centers for Disease Control and Prevention.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received March 3, 2011; revised May 2, 2011; accepted May 31, 2011; published OnlineFirst June 8, 2011.

References

- Byers T, Levin B, Rothenberger D, Dodd GD, Smith RA. American Cancer Society guidelines for screening and surveillance for early detection of colorectal polyps and cancer: update 1997. American Cancer Society Detection and Treatment Advisory Group on Colorectal Cancer. *CA Cancer J Clin* 1997;47:154-60.
- U.S. Preventive Services Task Force. 2nd ed. DiGiuseppe C, Atkins D, Woolf SH, editors. Guide to clinical preventive services. Baltimore, MD: Williams & Wilkins; 1996.
- Winawer SJ, Fletcher RH, Miller L, Godlee F, Stolar MH, Mulrow CD, et al. Colorectal cancer screening: clinical guidelines and rationale. *Gastroenterology* 1997;112:594-642.
- U.S. Preventive Services Task Force. Screening for colorectal cancer: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med* 2008;149:627-37.
- Steinwachs D, Allen JD, Barlow WE, Duncan RP, Egede LE, Friedman LS, et al. National Institutes of Health State-of-the-Science Conference Statement: enhancing use and quality of colorectal cancer screening. *Ann Intern Med* 2010;152:663-7.
- Swan J, Breen N, Graubard BI, McNeel TS, Blackman D, Tangka FK, et al. Data and trends in cancer screening in the United States. *Cancer* 2010;116:4872-81.
- Centers for Disease Control and Prevention. Vital signs: colorectal cancer screening among adults aged 50-75 years—United States, 2008. *MMWR* 2010;59:808-12.
- Shapiro JA, Seeff LC, Thompson TD, Nadel MR, Klabunde CN, Vernon SW. Colorectal cancer test use from the 2005 National Health Interview Survey. *Cancer Epidemiol Biomarkers Prev* 2008;17:1623-30.
- Seeff LC, Nadel MR, Klabunde CN, Thompson T, Shapiro JA, Vernon SW, et al. Patterns and predictors of colorectal cancer test use in the adult U.S. population. *Cancer* 2004;100:2093-103.
- Swan J, Breen N, Coates RJ, Rimer BK, Lee NC. Progress in cancer screening practices in the United States: results from the 2000 National Health Interview Survey. *Cancer* 2003;97:1528-40.
- Institute of Medicine. Crossing the quality chasm: a new health system for the 21st century. Washington, DC: National Academy Press; 2001.
- Doubeni CA, Laiyemo AO, Young AC, Klabunde CN, Reed G, Field TS, et al. Primary care, economic barriers to health care, and use of colorectal cancer screening tests among Medicare enrollees over time. *Ann Fam Med* 2010;8:299-307.
- Doubeni CA, Laiyemo AO, Reed G, Field TS, Fletcher RH. Socio-economic and racial patterns of colorectal cancer screening among Medicare enrollees in 2000 to 2005. *Cancer Epidemiol Biomarkers Prev* 2009;18:2170-5.
- Schenck AP, Peacock SC, Klabunde CN, Lapin P, Coan JF, Brown ML. Trends in colorectal cancer test use in the Medicare population, 1998-2005. *Am J Prev Med* 2009;37:1-7.
- Chen X, White MC, Peipins LA, Seeff LC. Increase in screening for colorectal cancer in older Americans: results from a national survey. *J Am Geriatr Soc* 2008;56:1511-6.

16. Nadel MR, Blackman DK, Shapiro JA, Seeff LC. Are people being screened for colorectal cancer as recommended? Results from the National Health Interview Survey. *Prev Med* 2002;35:199–206.
17. Fenton JJ, Cai Y, Green P, Beckett LA, Franks P, Baldwin LM. Trends in colorectal cancer testing among Medicare subpopulations. *Am J Prev Med* 2008;35:194–202.
18. Phillips KA, Liang SY, Ladabaum U, Haas J, Kerlikowske K, Lieberman D, et al. Trends in colonoscopy for colorectal cancer screening. *Med Care* 2007;45:160–7.
19. Centers for Disease Control and Prevention, National Center for Health Statistics. NHIS Public Use Data Release, NHIS Survey Description. [2000]. Available from: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHIS/2000/srvydesc.pdf (Appendix 1, page 65).
20. Centers for Disease Control and Prevention, National Center for Health Statistics. NHIS Public Use Data Release, NHIS Survey Description. [2003]. Available from: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHIS/2003/srvydesc.pdf (Appendix 1, page 65).
21. Centers for Disease Control and Prevention, National Center for Health Statistics. NHIS Public Use Data Release, NHIS Survey Description. [2005]. Available from: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHIS/2005/srvydesc.pdf (Appendix 1, page 72).
22. Centers for Disease Control and Prevention, National Center for Health Statistics. NHIS Public Use Data Release, NHIS Survey Description. [2008]. Available from: ftp://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/NHIS/2008/srvydesc.pdf (Appendix 1, page 78).
23. Centers for Disease Control and Prevention, National Center for Health Statistics. National Health Interview Survey: Questionnaires, Datasets, and Related Documentation. Available from: http://www.cdc.gov/nchs/nhis/nhis_questionnaires.htm.
24. The White House, Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. Available from: http://www.whitehouse.gov/omb/fedreg_1997standards/.
25. Centers for Disease Control and Prevention, National Center for Health Statistics. National Health Interview Survey: 2008 Imputed Family Income/Personal Earnings Files. Available from: <http://www.cdc.gov/nchs/nhis/2008imputedincome.htm>.
26. Bastani R. Intervention strategies in diverse populations. NIH State-of-the-Science Conference: Enhancing Use and Quality of Colorectal Cancer Screening, Program and Abstracts. Available from: http://consensus.nih.gov/2010/images/colorectal/colorectal_abstracts.pdf.
27. Cronin KA, Miglioretti DL, Krapcho M, Yu B, Geller BM, Carney PA, et al. Bias associated with self-report of prior screening mammography. *Cancer Epidemiol Biomarkers Prev* 2009;18:1699–705.
28. Holden DJ, Jonas DE, Porterfield DS, Reuland D, Harris R. Systematic review: enhancing the use and quality of colorectal cancer screening. *Ann Intern Med* 2010;152:668–76.
29. Patient Protection and Affordable Care Act. Public Law 111-148. [March 23, 2010]. Available from: <http://www.gpo.gov/fdsys/pkg/PLAW-111publ148.pdf>.

Cancer Epidemiology, Biomarkers & Prevention

Trends in Colorectal Cancer Test Use among Vulnerable Populations in the United States

Carrie N. Klabunde, Kathleen A. Cronin, Nancy Breen, et al.

Cancer Epidemiol Biomarkers Prev 2011;20:1611-1621. Published OnlineFirst June 8, 2011.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-11-0220](https://doi.org/10.1158/1055-9965.EPI-11-0220)

Cited articles This article cites 17 articles, 4 of which you can access for free at:
<http://cebp.aacrjournals.org/content/20/8/1611.full#ref-list-1>

Citing articles This article has been cited by 26 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/20/8/1611.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/20/8/1611>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.