

Research Article

Prognostic Impact of Cigarette Smoking on the Survival of Patients with Established Nasopharyngeal Carcinoma

Pu-Yun OuYang^{1,2}, Zhen Su^{1,2}, Yan-Ping Mao^{1,2}, Xue-Xia Liang^{1,2}, Qing Liu^{1,3}, Wuguo Deng¹, and Fang-Yun Xie^{1,2}

Abstract

Background: Cigarette smoking is associated with the etiology of nasopharyngeal carcinoma; however, the influence of smoking on survival in patients with established nasopharyngeal carcinoma remains unknown.

Methods: We retrospectively analyzed 1,849 patients with nasopharyngeal carcinoma who were categorized as never, former, and current smokers. Cumulative effect of smoking was defined in terms of pack-years. Associations between cigarette exposure and survival were estimated by Cox proportional hazards model.

Results: The risks of death, progression, locoregional relapse, and distant metastasis were significantly higher for former and current smokers (all $P \leq 0.002$) than never smokers. Heavy smokers with high pack-years had HRs for death of 3.31 [95% confidence interval (CI), 2.58–4.26; $P < 0.001$], for progression of 2.53 (95% CI, 2.03–3.16; $P < 0.001$), and for distant metastasis of 2.65 (95% CI, 1.89–3.70; $P < 0.001$). Specifically, in the cohort of 495 patients treated with intensity-modulated radiotherapy/three-dimensional conformal radiotherapy, we obtained similarly significant results. All of the survival outcomes remained significant in multivariate analyses.

Conclusions: Pretreatment cigarette smoking is an independent, poor prognostic factor for patients with nasopharyngeal carcinoma, which is associated with increased risk of death, progression, locoregional relapse, and distant metastasis, with the risk increasing with pack-years.

Impact: It is clear that cigarette smoking not only promotes carcinogenesis in the normal nasopharyngeal epithelium, but also affects the survival of patients with nasopharyngeal carcinoma. *Cancer Epidemiol Biomarkers Prev*; 22(12); 2285–94. ©2013 AACR.

Introduction

Nasopharyngeal carcinoma is a nonlymphomatous, squamous cell carcinoma that occurs in the epithelial lining of the nasopharynx. Despite improvements in the locoregional control rate because of the development of more precise imaging and radiotherapy techniques, and eradication of potential metastases by chemotherapy (1, 2), the survival of patients with advanced nasopharyngeal carcinoma remains unsatisfactory. Therefore, it is necessary to identify prognostic factors to recognize patients at high risk of failure. Recently, associations between cigarette smoking and survival have been demonstrated in several

types of cancer, including colon cancer (3), renal cell carcinoma (4), squamous cell carcinoma of the head and neck (HNSCC; refs. 5 and 6), and oropharyngeal cancer (7).

However, nasopharyngeal carcinoma has a distinct epidemiology, etiology (8), and clinical manifestation (9). The highest rates of incidence are observed in Southeast Asia, especially in Southern China where the incidence of nasopharyngeal carcinoma can be as high as 20 to 30 per 100,000 (10). In contrast, nasopharyngeal carcinoma is a relatively rare disease in Europe and the United States, with an incidence of 0.5 to 2 per 100,000 (11). The gender, age, and ethnic distribution of patients with nasopharyngeal carcinoma from different regions of China are far from uniform (8). The nasopharyngeal carcinoma–endemic populations have a particularly high intake of salt-preserved food, which is a unique risk factor for nasopharyngeal carcinoma; the potential of a high intake of salt-preserved food to lead to the development of malignant nasal and nasopharyngeal tumors is supported by data from rat models (12). More importantly, Epstein-Barr virus (EBV) plays a strongly causal role in the occurrence and development of nasopharyngeal carcinoma (13), whereas human papillomavirus is related to the etiology and prognosis of HNSCC (14). In addition, widely documented patterns of familial aggregation have demonstrated that some individuals have a genetic susceptibility to nasopharyngeal carcinoma (8). To

Authors' Affiliations: ¹State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine; and Departments of ²Radiation Oncology and ³Epidemiology, Sun Yat-sen University Cancer Center, Guangzhou, China

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Authors: Fang-Yun Xie, Department of Radiation Oncology, Sun Yat-sen University Cancer Center, State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, No. 651 Dongfeng Road East, Guangzhou 510060, China. Phone: 86-20-87343484; Fax: 86-20-87343484; E-mail: xiefy@sysucc.org.cn; and Wuguo Deng, dengwg@sysucc.org.cn

doi: 10.1158/1055-9965.EPI-13-0546

©2013 American Association for Cancer Research.

date, numerous case-control studies (15–17) examining cigarette smoking and the etiology of nasopharyngeal carcinoma have established that tobacco smoking is a consensus risk factor for this type of cancer (18); however, it is worth noting that an estimated two thirds of cases of World Health Organization type I (keratinizing squamous cell carcinoma) nasopharyngeal carcinoma are attributable to smoking, whereas type II (nonkeratinizing squamous carcinoma) and type III (undifferentiated carcinomas) nasopharyngeal carcinoma are not associated with smoking (19). The key issue related to nasopharyngeal carcinoma is that around 25%, 12%, and 63% of patients in North America have type I, II, and III, respectively whereas the histological distribution in Southern Chinese patients is 2%, 3%, and 95%, respectively (9). In addition, radiotherapy with or without chemotherapy, as the standard treatment for nasopharyngeal carcinoma (9), is more likely to be affected by smoking compared with surgery, which is the major therapeutic strategy for other types of head and neck cancers.

The association between cigarette smoking history and survival in patients with nasopharyngeal carcinoma remains to be explored. Therefore, we performed this study to elucidate the effect of cigarette smoking history on the clinicopathologic features and survival of patients with nasopharyngeal carcinoma.

Materials and Methods

Patient characteristics

Between January 2005 and May 2007, all newly diagnosed with biopsy-proven nasopharyngeal carcinoma (International Classification of Diseases for Oncology, 3rd Edition [ICD-O-3] C11, C11.0 to C11.9) patients who were hospitalized at Sun Yat-sen University Cancer Center were entered into our study. We excluded patients who were diagnosed with distant metastases before initial treatment by clinical symptoms, physical examinations, and imaging methods including chest radiography, bones scan, magnetic resonance imaging (MRI), and abdominal sonography. Medical records were reviewed to extract data on the clinicopathological features and smoking history of the patients, including age, gender, histological type, titer of immunoglobulin A against viral capsid antigen (VCA-IgA) before treatment, alcohol drinking status, smoking status at diagnosis, number of cigarettes smoked per day, number of years of smoking, and number of years since cessation. All patients were restaged according to the seventh edition of the AJCC/UICC Staging System for nasopharyngeal carcinoma (20).

Treatment and follow-up

All patients were treated by definitive radiotherapy. Details of the radiation techniques have been described previously (21, 22). In addition, institutional guidelines recommended no chemotherapy for patients with stage I or II disease, and both induction or adjuvant chemotherapy and concomitant chemotherapy for patients with stage III to IV disease, as defined by the sixth edition of the AJCC/

UICC Staging System for nasopharyngeal carcinoma or the Chinese 1992 Staging System for nasopharyngeal carcinoma at that time. Induction or adjuvant chemotherapy consisted of cisplatin with 5-fluorouracil, cisplatin with taxoids, or a triplet of cisplatin and 5-fluorouracil plus taxoids every 3 weeks for 2 to 3 cycles. Concomitant chemotherapy consisted of cisplatin on weeks 1, 4, and 7 of radiotherapy, or cisplatin given weekly.

Patients were followed up every 3 months during the first 2 years, and every 6 months thereafter until death. Patients with relapse, distant metastasis, or persistent disease were administered salvage treatments including re-irradiation, chemotherapy, and surgery. The follow-up duration was calculated from the first day of therapy to either the day of death or the day of the last examination.

Study endpoints

We mainly explored the associations between survival and cigarette smoking in terms of: (i) smoking status at diagnosis—patients were classified as never smokers (defined as never smoking before treatment), former smokers (defined as former smokers who had stopped smoking for 1 year or more before treatment), and current smokers (defined as smoking until the day of hospitalization or smokers who had stopped smoking for less than 1 year); and (ii) the cumulative effects of smoking in terms of pack-years, which was defined as the equivalent of smoking one pack of cigarettes per day for 1 year.

Our primary endpoint was overall survival (OS), defined as the time from treatment to death from any cause. Secondary endpoints were progression-free survival (PFS), defined as the time from treatment to disease progression or death from any cause, whichever occurred first; locoregional relapse-free survival (LRFS), defined as the time from treatment to the first locoregional relapse; and distant metastasis-free survival (DMFS), defined as the time from treatment to the first distant metastasis.

Statistical methods

All endpoints were examined using Kaplan–Meier methods and the log-rank test. Univariate survival analyses were performed in terms of smoking status at diagnosis and pack-years. Multivariate analyses were performed using the Cox proportional hazards model adjusting for basic characteristics such as age, gender, etc. Comparisons of demographic, clinical, and pathologic variables were performed using the χ^2 test or Fisher exact test for nominal variables as appropriate, or the Kruskal–Wallis test for ordinal categorical variables like T-stage, N-stage, and clinical stage. For continuous variables such as pack-years and age, we explored the effect using restricted cubic splines nested within Cox models by RCS-macro of SAS (SAS Institute); if there was linear effect, cutoff scores of the continuous variable were subsequently selected based on receiver operating characteristic (ROC) curve analysis (23). The sensitivity and specificity of each endpoint was plotted, thus generating ROC curves. The score closest to the point with both

maximum sensitivity and specificity (i.e., the point [0.0, 1.0] on the curve) was selected as the optimal cutoff score for prediction of survival. Two-sided *P* values less than 0.05 were considered significant.

Results

Treatment profiles, patterns of treatment failure, and survival

A total of 1,849 nasopharyngeal carcinoma patients were included in this study. After restaging according to the seventh edition of the AJCC/UICC Staging System, the clinical stage distribution of the 1,849 patients was: stage I, 90 (4.9%); stage II, 481 (26.0%); stage III, 796 (43.1%), and stage IV 482 (26.0%). Overall, 433/1,849 (23.4%) patients were treated with radiotherapy alone and 1,416/1,849 (76.6%) received radiotherapy plus chemotherapy. Of these 1,416 patients, 455 patients (32.1%) received induction chemotherapy and 522 patients (36.9%) received concomitant chemotherapy; a combination of induction and concomitant chemotherapy, concomitant and adjuvant chemotherapy, or the triplet of induction, concomitant, and adjuvant chemotherapy were administered to 363/1,416 (25.6%), 50/1,416 (3.5%), and 26/1,416 (1.8%) patients, respectively. With respect to radiotherapy, 1,354/1,849 (73.2%) were treated with conventional techniques, 457/1,849 (24.7%) with intensity-modulated radiotherapy (IMRT), and 38/1,849 (2.1%) with three-dimensional conformal radiotherapy (3DCRT).

Within a median follow-up duration of 73.5 months (range, 1.7–96.8 months), 150/1,849 (8.1%) patients developed locoregional relapse, 233/1,849 (12.6%) developed distant metastases, and 378/1,849 (20.4%) died. Twenty patients (1.1%) developed both locoregional relapse and distant metastases. The 3- and 5-year survival rates were as follows: OS, 88.8% and 82.2%; PFS, 79.5% and 74.2%; LRFS, 93.6% and 91.8%; and DMFS, 89.3% and 87.0%.

Demographic and clinicopathologic characteristics

The proportions of former smokers, current smokers, and never smokers were 9.1% (168/1849), 39.7% (734/1849) versus 51.2% (947/1849) in the entire population, and 9.7% (48/495), 36.4% (180/495) versus 53.9% (267/495) in the cohort of patients treated with IMRT/3DCRT. As shown in Table 1, there were no differences in the distribution of histological type or radiotherapy techniques for the entire patient cohort when stratified by smoking status. However, significant differences were observed in terms of age, gender, drinking status, VCA-IgA titer, T-stage, N-stage, clinical stage, and chemotherapy approach. Patients older than 40 years of age, and male patients with a drinking history were more frequent in the former and current smokers. There was a trend for a higher VCA-IgA titer among former and current smokers. In addition, the proportions of patients with advanced T-stage, N-stage, or clinical stage were higher for former and current smokers than never smokers. Accordingly, the proportion of patients adopting chemotherapy was higher for former and current smokers.

In the cohort of patients treated with IMRT/3DCRT, there were no significant differences in the distributions of histological type and chemotherapy when stratified by smoking status. Similarly to the entire population, the proportions of patients treated with IMRT/3DCRT who were older than 40 years of age, male with a positive drinking history, with a higher VCA-IgA titer, or with advanced T-stage, N-stage, or clinical stage were higher for former and current smokers.

Impact of cigarette smoking on survival in univariate analysis

In the entire population, OS, PFS, LRFS, and DMFS were all significantly poorer for former and current smokers than never smokers (Fig. 1). OS at 5 years was 75.4% for former smokers and 75.6% for current smokers versus 88.4% for never smokers (log-rank test, $P < 0.001$ and $P < 0.001$, respectively); the 5-year PFS rates were 67.8% and 65.0% versus 82.4% ($P < 0.001$ and $P < 0.001$, respectively); the 5-year LRFS rates were 88.6% and 89.1% versus 94.4% ($P < 0.001$, respectively); and the 5-year DMFS rates were 81.1% and 83.5% versus 90.7% ($P < 0.001$ and $P < 0.001$, respectively). Considering small number of former smokers and the similar survival rates between former and current smokers, we combined them into a single group—the smoking history group, the 5-year survival rates of which were as follow: OS 75.6%, PFS 65.6%, LRFS 89.0%, and DMFS 83.0% (Supplementary Fig. S1).

The cumulative effect of smoking was also strongly associated with the survival of patients with nasopharyngeal carcinoma. Among patients with smoking history, 32, 22, and 22 pack-years were identified as the cutoff scores for heavy and light smokers associated with OS, PFS, and DMFS, respectively. Heavy smokers had an HR of death of 3.31 [95% confidence interval (CI), 2.58–4.26; log-rank test, $P < 0.001$], HR of progression of 2.53 (95% CI, 2.03–3.16; $P < 0.001$), and HR of distant metastasis of 2.65 (95% CI, 1.89–3.70; $P < 0.001$) compared with light smokers (Fig. 2).

In the cohort of patients treated with IMRT/3DCRT, current smokers or former smokers also had higher risks of death ($P < 0.001$, $P = 0.002$) and disease progression ($P < 0.001$, $P = 0.004$) than never smokers (Supplementary Fig. S2a and S2b). The significant differences remained unchanged when combined current and former smokers into the smoking history group (Supplementary Fig. S3). And among this group, heavy smokers with more than 25 pack-years of cigarettes had an HR of death of 3.61 (95% CI, 2.12–6.14; $P < 0.001$), and those with more than 16 pack-years had an HR of progression of 3.01 (95% CI, 1.83–4.97; $P < 0.001$) compared with light smokers (Supplementary Fig. S2c and S2d).

Impact of cigarette smoking on survival in multivariate analysis

Using restricted cubic splines nested within Cox models, the variables of age and pack-years were tested in multivariate analysis in continuous and nonlinear fashion. Resultantly, both of them showed linear effects in

Table 1. Demographic and clinicopathologic characteristics of the study population stratified by smoking status

| Factor | The entire population | | | P | The IMRT/3DCRT cohort | | | P |
|-------------------|------------------------------------|-------------------------------------|--------------------------------------|---------------------|------------------------------------|------------------------------------|--------------------------------------|---------------------|
| | Never smoker (N = 947) N (%) | Former smoker (N = 168) N (%) | Current smoker (N = 734) N (%) | | Never smoker (N = 267) N (%) | Former smoker (N = 48) N (%) | Current smoker (N = 180) N (%) | |
| Age group | | | | <0.001 ^a | | | | <0.001 ^a |
| ≤30 | 85 (9.0) | 2 (1.2) | 36 (4.9) | | 25 (9.4) | 1 (2.1) | 9 (5.0) | |
| 31–40 | 311 (32.8) | 25 (14.9) | 167 (22.8) | | 85 (31.8) | 6 (12.5) | 47 (26.1) | |
| 41–50 | 289 (30.5) | 57 (33.9) | 243 (33.1) | | 88 (33.0) | 22 (45.8) | 56 (31.1) | |
| 51–60 | 174 (18.4) | 55 (32.7) | 192 (26.2) | | 43 (16.1) | 9 (18.8) | 38 (21.1) | |
| 61–70 | 73 (7.7) | 24 (14.3) | 82 (11.2) | | 22 (8.2) | 8 (16.7) | 25 (13.9) | |
| ≥71 | 15 (1.6) | 5 (3.0) | 14 (1.9) | | 4 (1.5) | 2 (4.2) | 5 (2.8) | |
| Gender | | | | <0.001 | | | | <0.001 |
| Male | 509 (53.7) | 163 (97.0) | 726 (98.9) | | 157 (58.8) | 45 (93.8) | 175 (97.2) | |
| Female | 438 (46.3) | 5 (3.0) | 8 (1.1) | | 110 (41.2) | 3 (6.2) | 5 (2.8) | |
| Drinking status | | | | <0.001 ^b | | | | <0.001 ^b |
| Never | 908 (95.9) | 125 (74.4) | 537 (73.2) | | 259 (97.0) | 37 (77.1) | 120 (66.7) | |
| Former | 6 (0.6) | 17 (10.1) | 5 (0.7) | | 3 (1.1) | 5 (10.4) | 3 (1.7) | |
| Current | 33 (3.5) | 26 (15.5) | 192 (26.2) | | 5 (1.9) | 6 (12.5) | 57 (31.6) | |
| VCA-IgA | | | | 0.001 | | | | 0.005 |
| ≤1:160 | 465 (49.1) | 81 (48.2) | 292 (39.8) | | 149 (55.8) | 24 (50.0) | 72 (40.0) | |
| >1:160 | 482 (50.9) | 87 (51.8) | 442 (60.2) | | 118 (44.2) | 24 (50.0) | 108 (60.0) | |
| Histological type | | | | 0.348 ^b | | | | 0.296 ^b |
| I | 4 (0.4) | 1 (0.6) | 1 (0.1) | | 1 (0.4) | 0 (0) | 0 (0) | |
| II | 56 (5.9) | 7 (4.2) | 32 (4.4) | | 20 (7.5) | 1 (2.1) | 7 (3.9) | |
| III | 887 (93.7) | 160 (95.2) | 701 (95.5) | | 246 (92.1) | 47 (97.9) | 173 (96.1) | |
| T-stage | | | | 0.042 ^a | | | | 0.004 ^a |
| T1 | 144 (15.2) | 25 (14.9) | 80 (10.9) | | 51 (19.1) | 8 (16.7) | 21 (11.7) | |
| T2 | 250 (26.4) | 41 (24.4) | 198 (27.0) | | 74 (27.7) | 10 (20.8) | 38 (21.1) | |
| T3 | 356 (37.6) | 68 (40.5) | 271 (36.9) | | 99 (37.1) | 21 (43.8) | 75 (41.7) | |
| T4 | 197 (20.8) | 34 (20.2) | 185 (25.2) | | 43 (16.1) | 9 (18.8) | 46 (25.6) | |
| N-stage | | | | <0.001 ^a | | | | 0.066 ^a |
| N0 | 175 (18.5) | 54 (32.1) | 94 (12.8) | | 61 (22.8) | 18 (37.5) | 38 (21.1) | |
| N1 | 555 (58.6) | 82 (48.8) | 390 (53.1) | | 150 (56.2) | 22 (45.8) | 96 (53.3) | |
| N2 | 190 (20.1) | 25 (14.9) | 203 (27.7) | | 45 (16.9) | 7 (14.6) | 37 (20.6) | |
| N3 | 27 (2.9) | 7 (4.2) | 47 (6.4) | | 11 (4.1) | 1 (2.1) | 9 (5.0) | |
| Clinical stage | | | | <0.001 ^a | | | | 0.015 ^a |
| I | 51 (5.4) | 14 (8.3) | 25 (3.4) | | 23 (8.6) | 6 (12.5) | 11 (6.1) | |
| II | 269 (28.4) | 42 (25.0) | 170 (23.2) | | 81 (30.3) | 9 (18.8) | 38 (21.1) | |
| III | 408 (43.1) | 72 (42.9) | 316 (43.1) | | 111 (41.6) | 23 (47.9) | 78 (43.3) | |
| IVa | 192 (20.3) | 33 (19.6) | 177 (24.1) | | 41 (15.4) | 9 (18.8) | 44 (24.4) | |
| IVb | 27 (2.9) | 7 (4.2) | 46 (6.3) | | 11 (4.1) | 1 (2.1) | 9 (5.0) | |
| Chemotherapy | | | | <0.001 | | | | 0.098 |
| No | 225 (23.8) | 59 (35.1) | 149 (20.3) | | 69 (25.8) | 18 (37.5) | 40 (22.2) | |
| Yes | 722 (76.2) | 109 (64.9) | 585 (79.7) | | 198 (74.2) | 30 (62.5) | 140 (87.8) | |
| Radiotherapy | | | | 0.504 ^b | | | | |
| IMRT | 246 (26.0) | 45 (26.8) | 166 (22.6) | | | | | |
| 3DCRT | 21 (2.2) | 3 (1.8) | 14 (1.9) | | | | | |
| CRT | 680 (71.8) | 120 (71.4) | 554 (75.5) | | | | | |

NOTE: A former smoker was defined as an individual who had not smoked for 12 months or more at enrollment.
^aKruskal–Wallis test.
^bFisher's exact probabilities test.




Figure 1. (A) Overall survival, (B) progression-free survival, (C) locoregional relapse-free survival, and (D) distant metastasis-free survival curves of patients in the entire population according to smoking status at diagnosis. A former smoker was defined as an individual who had not smoked for 12 months or more at enrollment.

most cases. However, the age was excluded as a covariate when we analyzed the impact of smoking history in PFS in the entire population; the pack-years was showed to be a nonsignificant prognosis of LRFS for patients with smoking history in the entire population (see Supplementary Material).

In multivariate analysis, smoking history (former and current smokers) versus no smoking history (never smokers) and high pack-years versus low pack-years, together with age (continuous), T-stage, and N-stage were found to be significant, independent predictors of overall survival for both the entire population and

patients treated with IMRT/3DCRT (Table 2). In addition, we also assessed the association between smoking history and OS across strata of other potential predictors of patient outcome in the entire population (Table 3). The effect of smoking history on the risk of death was not significantly modified by age, titer of VCA-IgA, clinical stage of disease, chemotherapy approach, or radiation technique. However, the impact of smoking history in increasing risk of death was not observed among female patients (adjusted HR = 1.75; 95% CI, 0.66–4.64; $P = 0.257$) and patients with drinking history (adjusted HR = 1.78; 95% CI, 0.76–4.16; $P = 0.187$).




Figure 2. (A) Overall survival, (B) progression-free survival, and (C) distant metastasis-free survival curves of patients according to pack-years. A pack-year was defined as the equivalent of smoking one pack of cigarettes per day for 1 year.

In second analysis (Table 4), former smokers had an HR of death of 1.46 (95% CI, 1.20–1.78; $P < 0.001$) and HR of progression of 1.41 (95% CI, 1.19–1.68; $P < 0.001$)

compared with never smokers; whereas current smokers had an HR of death of 1.67 (95% CI, 1.26–2.21; $P < 0.001$) and HR of progression of 1.74 (95% CI, 1.37–2.21; $P < 0.001$) compared with never smokers. The risks of death and progression for heavy smokers were higher than light smokers, with HRs of 2.10 (95% CI, 1.58–2.79; $P < 0.001$) and 1.64 (95% CI, 1.23–2.19; $P < 0.001$), respectively. When the pack-years was evaluated as a continuous variable, the HRs for death and progression increased by 2% per pack-year.

In the cohort of patients treated with IMRT/3DCRT, former smokers had an HR of death of 1.51 (95% CI, 1.05–2.18; $P = 0.026$) and HR of progression of 1.44 (95% CI, 1.03–2.01; $P = 0.033$) compared with never smokers, whereas current smokers had an HR of death of 1.90 (95% CI, 1.12–3.22; $P = 0.017$) and HR of progression of 2.37 (95% CI, 1.50–3.75; $P < 0.001$) compared with never smokers. In addition, compared with light smokers, the HRs for death and progression for heavy smokers were extremely high, at 2.41 (95% CI, 1.17–4.96; $P = 0.017$) and 2.60 (95% CI, 1.44–4.69; $P = 0.002$), respectively. Similarly, when the pack-years was evaluated as a continuous variable, the HRs for death and progression increased by 3% per pack-year (Table 4).

Discussion

Numerous case-control studies have convincingly demonstrated that cigarette smoking greatly increases the risk of developing nasopharyngeal carcinoma (15–17); however, the possibility of cigarette smoking being a predictor of overall survival in nasopharyngeal carcinoma has only been shown in one study (24), the analyzed data extracted from an epidemiological investigation which was limited with respect to the treatment regimens, follow-up care, disease stage, treatment outcomes and other prognostic factors. This study is the first investigation of a large number of patients ($N = 1,849$) to demonstrate that cigarette smoking is an independent, poor prognostic factor for survival in patients with established nasopharyngeal carcinoma after adjusting for age (continuous), gender, drinking status, histological type, T-stage, N-stage, VCA-IgA titer (\leq and $>1:160$), radiotherapy techniques, and chemotherapy regimens. We subsequently evaluated the increased risk of death and progression in terms of pack-years of cigarettes. In addition, we performed a specific analysis of the cohort of 495 patients treated with IMRT/3DCRT to account for the heterogeneity of radiotherapy techniques. According to these analyses, it is clear that cigarette smoking not only promotes carcinogenesis in the normal nasopharyngeal epithelium (15–17), but also affects the survival of patients with nasopharyngeal carcinoma. This result is not unexpected. In a recent study of patients with oral, pharyngeal, and larynx cancer that did not include patients with nasopharyngeal carcinoma, smoking was associated with an increased cancer-specific mortality (25). Previous studies have shown that smoking status at diagnosis is associated with the survival of patients with HNSCC (5, 6), and

Table 2. Multivariable Cox proportional hazards models for overall survival in terms of smoking history (smoking history vs. no smoking history) and pack-years (heavy vs. light smokers)

| Variable | The entire population | | | | The IMRT/3DCRT cohort | | | |
|----------------------|--------------------------------|--------|--------------------------------------|--------|------------------------------|--------|--------------------------------------|-------|
| | Smoking history (N = 1,849) | | Pack-years (N = 902) ^a | | Smoking history (N = 495) | | Pack-years (N = 228) ^b | |
| | HR (95% CI) | P | HR (95% CI) | P | HR (95% CI) | P | HR (95% CI) | P |
| Age (continuous) | 1.04 (1.03–1.05) | <0.001 | 1.03 (1.02–1.04) | <0.001 | 1.05 (1.03–1.07) | <0.001 | 1.03 (0.99–1.06) | 0.115 |
| Gender | 0.96 (0.70–1.33) | 0.811 | 0.98 (0.45–2.16) | 0.961 | 1.15 (0.57–1.65) | 0.693 | 1.96 (0.69–5.57) | 0.210 |
| Drinking | 1.02 (0.82–1.28) | 0.847 | 0.95 (0.74–1.21) | 0.681 | 0.97 (0.57–1.65) | 0.923 | 0.85 (0.47–1.54) | 0.598 |
| VCA-IgA ^c | 1.02 (0.83–1.26) | 0.830 | 1.02 (0.78–1.32) | 0.899 | 1.13 (0.71–1.81) | 0.612 | 1.00 (0.57–1.77) | 0.990 |
| Histological type | 1.08 (0.70–1.65) | 0.738 | 1.01 (0.60–1.72) | 0.965 | 1.30 (0.41–4.17) | 0.658 | 2.12 (0.28–15.91) | 0.467 |
| T-stage | 1.62 (1.43–1.82) | <0.001 | 1.45 (1.25–1.68) | <0.001 | 1.62 (1.39–2.49) | <0.001 | 1.52 (1.08–2.14) | 0.016 |
| N-stage | 1.70 (1.49–1.95) | <0.001 | 1.64 (1.40–1.93) | <0.001 | 1.53 (1.12–2.08) | 0.007 | 1.64 (1.12–2.40) | 0.011 |
| Radiotherapy | 0.94 (0.82–1.07) | 0.331 | 0.99 (0.85–1.16) | 0.924 | 1.15 (0.41–3.21) | 0.794 | 1.15 (0.34–3.94) | 0.821 |
| Chemotherapy | 0.96 (0.90–1.02) | 0.194 | 0.98 (0.91–1.07) | 0.683 | 0.91 (0.78–1.07) | 0.267 | 0.88 (0.73–1.06) | 0.179 |
| Smoking history | 1.73 (1.32–2.27) | <0.001 | – | – | 2.08 (1.14–3.79) | 0.017 | – | – |
| Pack-years | – | – | 2.10 (1.58–2.79) | <0.001 | – | – | 2.41 (1.17–4.96) | 0.017 |

^aPack-years: ≤ vs. > 32.^bPack-years: ≤ vs. > 25.^cTiter of VCA-IgA: ≤ vs. > 1:160.

the risk of death increased with increasing exposure to tobacco as measured in pack-years or years of smoking (26). Smoking was also identified as an indicator of poor prognosis in patients with oropharyngeal cancer, regardless of the tumor p16 status or treatment (7). In addition, patients with renal cell carcinoma and a history of smoking were more likely to have advanced pathologic features and poorer survival outcomes (4). Similar significant associations between smoking and poorer disease-free survival and time to recurrence have also been observed in patients with colon cancer (3).

Recently, Duffy and colleagues found that smoking was independently associated with a higher serum interleukin-6 (IL-6) level among patients with head and neck cancer (27), and a higher IL-6 level was predictive of an increased risk of recurrence and poorer OS (28). In nasopharyngeal carcinoma, IL-6 facilitates tumor carcinogenesis and malignancy enhancement via activation of the *STAT3*, and a higher IL-6 level is associated with advanced stage nasopharyngeal carcinoma (29). Second, it has been demonstrated that smoking is involved in EBV activation and smokers have increased seropositivity for the EBV VCA-IgA antibody (16); and EBV status is strongly associated with the risk of developing nasopharyngeal carcinoma (30) and patients' LRFS and OS (31, 32). Therefore, via EBV activation, smoking may also contribute to the survival differences between patients who smoke and those who do not. Moreover, evidence indicates that smoking exacerbates tissue hypoxia and can lead to smoking-induced tissue hypoxia in healthy human smokers (33), and inhalation of carbon monoxide—one component of cigarette smoke—can reduce the control of

tumors by radiotherapy in animal models (34). In addition, patients with head and neck cancer (including nasopharyngeal carcinoma) with a higher smoking status have higher venous blood levels of carboxyhemoglobin, which results in reduced oxygen supply to the tumors (35). It is known that hypoxia induces the expression of a variety of genes associated with an aggressive malignant phenotype, including genes involved in stem cell maintenance, invasion, angiogenesis, and extravasation (36). The transcription of hypoxia-related genes, which is predominantly mediated by the hypoxia-inducible factor-1 α (*HIF-1 α*) in cooperation with *HIF-2 α* , promotes tumor angiogenesis and the proliferation, invasion, and metastasis of tumor cells (37). Furthermore, tumor hypoxia has been acknowledged to affect the responses to both radiotherapy (38) and chemotherapy (39), and has been associated with poorer OS in HNSCC (40). In addition, basic research has demonstrated that cigarette smoke condensate promotes chemoresistance via *Akt*-mediated regulation of the activity of the ATP-binding cassette transporter G2, and may also contribute to tumor recurrence, invasion, or metastasis by increasing the proportion of cancer stem-like cells (41). Thus, the presence of residual smoke condensate in former and current smokers may reduce the effect of chemotherapy and promote tumor progression.

In this study, the impact of smoking in increasing risk of death was not observed among female patients and patients with drinking history. This may have been because of small number of patients in the individual subgroups. Moreover, it was interesting that there were no significant differences in the survival outcomes of former and current smokers; both groups had similarly

Table 3. Subgroup analysis of OS in terms of smoking history vs. no smoking history by the characteristics of patients in the entire population

| Factor | 5-year OS rate (%) | P for OS by each factor | No. of events/no. at risk | | Adjusted HR (95% CI) ^a | P for smoking history vs. no smoking history |
|---------------------|--------------------|-------------------------|---------------------------|-----------------|-----------------------------------|--|
| | | | No smoking history | Smoking history | | |
| Overall | 82.2 | | 126/947 | 252/902 | 1.73 (1.32–2.27) | <0.001 |
| Age (year) | | <0.001 | | | | |
| ≤52 | 86.4 | | 83/723 | 126/589 | 1.79 (1.26–2.54) | 0.001 |
| >52 | 71.8 | | 43/224 | 126/313 | 2.02 (1.31–3.14) | 0.002 |
| Gender | | <0.001 | | | | |
| Male | 80.4 | | 65/509 | 245/889 | 1.69 (1.27–2.24) | <0.001 |
| Female | 87.6 | | 61/438 | 7/13 | 1.75 (0.66–4.64) | 0.257 |
| Drinking status | | 0.017 | | | | |
| Yes | 77.8 | | 7/39 | 65/240 | 1.78 (0.76–4.16) | 0.187 |
| No | 83.0 | | 119/908 | 187/662 | 1.79 (1.34–2.38) | <0.001 |
| VCA-IgA | | 0.004 | | | | |
| ≤1:160 | 84.7 | | 53/465 | 94/373 | 1.77 (1.25–2.52) | 0.001 |
| >1:160 | 80.1 | | 73/482 | 158/529 | 1.78 (1.34–2.36) | <0.001 |
| Clinical stage | | <0.001 | | | | |
| I + II | 92.6 | | 11/320 | 37/251 | 3.67 (1.54–8.77) | 0.003 |
| III + IV | 77.4 | | 115/627 | 215/651 | 1.76 (1.32–2.35) | <0.001 |
| Chemotherapy | | 0.001 | | | | |
| Yes | 80.5 | | 104/722 | 210/694 | 1.74 (1.28–2.35) | <0.001 |
| No | 87.5 | | 22/225 | 42/208 | 2.10 (1.05–4.18) | 0.035 |
| Radiation technique | | 0.007 | | | | |
| IMRT/3DCRT | 86.9 | | 23/267 | 55/228 | 2.08 (1.14–3.79) | 0.017 |
| Conventional | 80.4 | | 103/680 | 197/674 | 1.60 (1.18–2.17) | 0.003 |

^aAdjusted for age (continuous), gender, drinking status, histological type, T-stage, N-stage, VCA-IgA titer (≤ and > 1:160), radiation techniques, and chemotherapy regimens.

poor survival outcomes compared with never smokers, even after adjusting for factors such as drinking behavior. Heavy smokers—in terms of high pack-years—are likely to have been subjected to the prolonged, severe, cumulative effects of smoking, and their relatively poor survival rates also demonstrated that the influence of smoking develops after long-term exposure. Therefore, despite the fact that former smokers were patients who had stopped smoking for at least 1 year, they may not have totally eliminated the residual condensates of cigarette smoke during this time, and the ability of smoking to enhance the IL-6 level, activate EBV, exacerbate tissue hypoxia, and promote chemoresistance and tumor progression may have already occurred and are likely to be maintained after stopping smoking. Finally, a recent study (42) found that 13% of cancer patients who did not smoke in the 7 days before surgery had resumed smoking within 12 months after surgery; their resumption of smoking was related to a higher perceived difficulty of quitting and lower perceptions of their cancer-related risk. Research has shown that continued smoking after diagnosis has immediate adverse impacts, including reduced efficacy of cancer treatment (5, 41), increased proportions of cancer

stem-like cells (41), higher rates of treatment complications and side effects (43, 44), higher treatment-related weight loss (45), and a poorer quality of life (46). Unfortunately, we did not collect data on the patients who resumed or continued smoking during treatment or follow-up in this study; therefore, the possibility that some former smokers resumed smoking during treatment or follow-up cannot be ignored.

The following limitations of this study deserve comment. First, like other retrospective studies in nasopharyngeal carcinoma, the treatment regimens were not totally consistent with the latest NCCN guidelines, for example, the number of patients with stage I and II disease (Table 1) was not equal to the number of patients who received radiotherapy alone. This is mainly because the patients were initially staged according to the sixth edition of the AJCC/UICC Staging System or the Chinese 1992 Staging System for nasopharyngeal carcinoma before making treatment decisions, whereas we restaged the patients according to the seventh edition of the AJCC/UICC Staging System during the retrospective analysis. In addition, during the period when the patients were treated, many patients were encouraged to participate in

Table 4. Effect of various smoking exposure measures on overall survival and progression-free survival in patients with nasopharyngeal carcinoma after adjusting for significant prognostic factors^a

| Variables | The entire population | | The IMRT/3DCRT cohort | |
|-----------------------------|---------------------------------|--|---------------------------------|--|
| | Overall survival HR (95% CI) | Progression-free survival HR (95% CI) | Overall survival HR (95% CI) | Progression-free survival HR (95% CI) |
| Smoking status at diagnosis | | | | |
| Former vs. never | 1.46 (1.20–1.78) | 1.41 (1.19–1.68) | 1.51 (1.05–2.18) | 1.44 (1.03–2.01) |
| Current vs. never | 1.67 (1.26–2.21) | 1.74 (1.37–2.21) | 1.90 (1.12–3.22) | 2.37 (1.50–3.75) |
| Pack-years | | | | |
| Heavy vs. light | 2.10 ^b (1.58–2.79) | 1.64 ^c (1.23–2.19) | 2.41 ^d (1.17–4.96) | 2.60 ^e (1.44–4.69) |
| Pack-years (continuous) | 1.02 (1.01–1.02) | 1.02 (1.01–1.02) | 1.03 (1.01–1.04) | 1.03 (1.02–1.04) |

^aAdjusted for age (continuous), gender, drinking status, histological type, T-stage, N-stage, VCA-IgA titer (\leq and $>1:160$), radiation techniques, and chemotherapy regimens.

^bPack-years: \leq versus >32 .

^cPack-years: \leq versus >22 .

^dPack-years: \leq versus >25 .

^ePack-years: \leq versus >16 .

randomized trials, which also resulted in heterogeneous treatment strategies. However, we conducted multivariate analyses accounting for radiation techniques and chemotherapy approaches, and specifically analyzed the IMRT/3DCRT cohort independently. Second, we were unable to collect adequate information about the patients' pretreatment EBV DNA copy number, which has been shown to be superior to the serum EBV VCA-IgA antibody titer for making prognostic predictions in nasopharyngeal carcinoma (47). However, the VCA-IgA antibody titer, and not the EBV DNA copy number, has been demonstrated to be highly associated with smoking status in nasopharyngeal carcinoma (16). Future studies based on the EBV DNA copy number are being planned. Finally, smoking and drinking status at diagnosis were extracted from medical records, rather than determined by standardized questionnaires at enrollment. This is an inevitable limitation caused by retrospective study. However, this sort of data are of high reliability because of medical records' strong validity of law in China.

In conclusion, this study of 1,849 patients demonstrated that pretreatment smoking history was an independent, poor prognostic factor for nasopharyngeal carcinoma patients; smoking was associated with an increased risk of death, progression, locoregional relapse and distant

metastasis, with the risks increasing with the number of pack-years, years of smoking, and cigarettes per day.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: P.-Y. OuYang, W. Deng, F.-Y. Xie

Development of methodology: P.-Y. OuYang

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): P.-Y. OuYang, Z. Su, Y.-P. Mao

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): P.-Y. OuYang, Z. Su, Y.-P. Mao, Q. Liu, W. Deng, F.-Y. Xie

Writing, review, and/or revision of the manuscript: P.-Y. OuYang, Z. Su, Y.-P. Mao, W. Deng, F.-Y. Xie

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): P.-Y. OuYang, Z. Su, Y.-P. Mao, W. Deng, F.-Y. Xie

Study supervision: P.-Y. OuYang, X.-X. Liang, W. Deng, F.-Y. Xie

Grant Support

This work was supported by grants from the National Natural Science Foundation of China (No. 30930045) and the Science Foundation of Key Hospital Clinical Program of Ministry of Health P.R. China (No. 2010-439).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received May 27, 2013; revised August 30, 2013; accepted September 12, 2013; published OnlineFirst November 19, 2013.

References

- Ouyang PY, Xie C, Mao YP, Zhang Y, Liang XX, Su Z, et al. Significant efficacies of neoadjuvant and adjuvant chemotherapy for nasopharyngeal carcinoma by meta-analysis of published literature-based randomized, controlled trials. *Ann Oncol* 2013;24:2136–46.
- Baujatz B, Audry H, Bourhis J, Chan AT, Onat H, Chua DT, et al. Chemotherapy in locally advanced nasopharyngeal carcinoma: an individual patient data meta-analysis of eight randomized trials and 1753 patients. *Int J Radiat Oncol Biol Phys* 2006;64:47–56.
- Phipps AI, Shi Q, Newcomb PA, Nelson GD, Sargent DJ, Alberts SR, et al. Associations between cigarette smoking status and colon cancer prognosis among participants in North Central Cancer Treatment Group Phase III Trial N0147. *J Clin Oncol* 2013;31:2016–23.
- Kroeger N, Klatte T, Birkhauser FD, Rampersaud EN, Seligson DB, Zomorodian N, et al. Smoking negatively impacts renal cell carcinoma overall and cancer-specific survival. *Cancer* 2012;118:1795–802.

5. Browman GP, Wong G, Hodson I, Sathya J, Russell R, McAlpine L, et al. Influence of cigarette smoking on the efficacy of radiation therapy in head and neck cancer. *N Engl J Med* 1993;328:159-63.
6. Duffy SA, Ronis DL, McLean S, Fowler KE, Gruber SB, Wolf GT, et al. Pretreatment health behaviors predict survival among patients with head and neck squamous cell carcinoma. *J Clin Oncol* 2009;27:1969-75.
7. Gillison ML, Zhang Q, Jordan R, Xiao W, Westra WH, Trotti A, et al. Tobacco smoking and increased risk of death and progression for patients with p16-positive and p16-negative oropharyngeal cancer. *J Clin Oncol* 2012;30:2102-11.
8. Chang ET, Adami HO. The enigmatic epidemiology of nasopharyngeal carcinoma. *Cancer Epidemiol Biomarkers Prev* 2006;15:1765-77.
9. Wei WI, Sham JS. Nasopharyngeal carcinoma. *Lancet* 2005;365:2041-54.
10. Cao SM, Simons MJ, Qian CN. The prevalence and prevention of nasopharyngeal carcinoma in China. *Chin J Cancer* 2011;30:114-9.
11. Ferlay J, Bray F, Pisani P, Parkin DM. Cancer incidence, mortality and prevalence worldwide. 2.0 ed. Lyon: IARC Press; 2004.
12. Zheng X, Luo Y, Christensson B, Drettner B. Induction of nasal and nasopharyngeal tumours in Sprague-Dawley rats fed with Chinese salted fish. *Acta Otolaryngol* 1994;114:98-104.
13. Liebowitz D. Nasopharyngeal carcinoma: the Epstein-Barr virus association. *Semin Oncol* 1994;21:376-81.
14. Gillison ML, Koch WM, Capone RB, Spafford M, Westra WH, Wu L, et al. Evidence for a causal association between human papillomavirus and a subset of head and neck cancers. *J Natl Cancer Inst* 2000;92:709-20.
15. Ji X, Zhang W, Xie C, Wang B, Zhang G, Zhou F. Nasopharyngeal carcinoma risk by histologic type in central China: impact of smoking, alcohol and family history. *Int J Cancer* 2011;129:724-32.
16. Xu FH, Xiong D, Xu YF, Cao SM, Xue WQ, Qin HD, et al. An epidemiological and molecular study of the relationship between smoking, risk of nasopharyngeal carcinoma, and Epstein-Barr virus activation. *J Natl Cancer Inst* 2012;104:1396-410.
17. Hsu WL, Chen JY, Chien YC, Liu MY, You SL, Hsu MM, et al. Independent effect of EBV and cigarette smoking on nasopharyngeal carcinoma: a 20-year follow-up study on 9,622 males without family history in Taiwan. *Cancer Epidemiol Biomarkers Prev* 2009;18:1218-26.
18. Jia WH, Qin HD. Non-viral environmental risk factors for nasopharyngeal carcinoma: a systematic review. *Semin Cancer Biol* 2012;22:117-26.
19. Vaughan TL, Shapiro JA, Burt RD, Swanson GM, Berwick M, Lynch CF, et al. Nasopharyngeal cancer in a low-risk population: defining risk factors by histological type. *Cancer Epidemiol Biomarkers Prev* 1996;5:587-93.
20. Edge SB, Byrd DR, Compton CC, Fritz AG, Greene FL, Trotti A. *AJCC cancer staging manual*. 7th ed. New York: Springer; 2009.
21. Ma J, Liu L, Tang L, Zong J, Lin A, Lu T, et al. Retropharyngeal lymph node metastasis in nasopharyngeal carcinoma: prognostic value and staging categories. *Clin Cancer Res* 2007;13:1445-52.
22. Liang SB, Sun Y, Liu LZ, Chen Y, Chen L, Mao YP, et al. Extension of local disease in nasopharyngeal carcinoma detected by magnetic resonance imaging: improvement of clinical target volume delineation. *Int J Radiat Oncol Biol Phys* 2009;75:742-50.
23. Hanley JA. Receiver operating characteristic (ROC) methodology: the state of the art. *Crit Rev Diagn Imaging* 1989;29:307-35.
24. Shen GP, Xu FH, He F, Ruan HL, Cui C, Chen LZ, et al. Pretreatment lifestyle behaviors as survival predictors for patients with nasopharyngeal carcinoma. *PLoS One* 2012;7:e36515.
25. Warren GW, Kasza KA, Reid ME, Cummings KM, Marshall JR. Smoking at diagnosis and survival in cancer patients. *Int J Cancer* 2013;132:401-10.
26. Mayne ST, Cartmel B, Kirsh V, Goodwin WJ Jr. Alcohol and tobacco use prediagnosis and postdiagnosis, and survival in a cohort of patients with early stage cancers of the oral cavity, pharynx, and larynx. *Cancer Epidemiol Biomarkers Prev* 2009;18:3368-74.
27. Duffy SA, Teknos T, Taylor JM, Fowler KE, Islam M, Wolf GT, et al. Health behaviors predict higher interleukin-6 levels among patients newly diagnosed with head and neck squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 2013;22:374-81.
28. Duffy SA, Taylor JM, Terrell JE, Islam M, Li Y, Fowler KE, et al. Interleukin-6 predicts recurrence and survival among head and neck cancer patients. *Cancer* 2008;113:750-7.
29. Zhang G, Tsang CM, Deng W, Yip YL, Lui VW, Wong SC, et al. Enhanced IL-6/IL-6R signaling promotes growth and malignant properties in EBV-infected premalignant and cancerous nasopharyngeal epithelial cells. *PLoS One* 2013;8:e62284.
30. Chien YC, Chen JY, Liu MY, Yang HI, Hsu MM, Chen CJ, et al. Serologic markers of Epstein-Barr virus infection and nasopharyngeal carcinoma in Taiwanese men. *N Engl J Med* 2001;345:1877-82.
31. Liu MT, Yeh CY. Prognostic value of anti-Epstein-Barr virus antibodies in nasopharyngeal carcinoma (NPC). *Radiat Med* 1998;16:113-7.
32. Wang WY, Twu CW, Chen HH, Jiang RS, Wu CT, Liang KL, et al. Long-term survival analysis of nasopharyngeal carcinoma by plasma Epstein-Barr virus DNA levels. *Cancer* 2013;119:963-70.
33. Jensen JA, Goodson WH, Hopf HW, Hunt TK. Cigarette smoking decreases tissue oxygen. *Arch Surg* 1991;126:1131-4.
34. Grau C, Nordmark M, Khalil AA, Horsman MR, Overgaard J. Effect of carbon monoxide breathing on hypoxia and radiation response in the SCCVII tumor in vivo. *Int J Radiat Oncol Biol Phys* 1994;29:449-54.
35. Hoff CM, Grau C, Overgaard J. Effect of smoking on oxygen delivery and outcome in patients treated with radiotherapy for head and neck squamous cell carcinoma—a prospective study. *Radiother Oncol* 2012;103:38-44.
36. Semenza GL. Hypoxia-inducible factors: mediators of cancer progression and targets for cancer therapy. *Trends Pharmacol Sci* 2012;33:207-14.
37. Perez-Sayans M, Suarez-Penaranda JM, Pilar GD, Barros-Angueira F, Gandara-Rey JM, Garcia-Garcia A. Hypoxia-inducible factors in OSCC. *Cancer Lett* 2011;313:1-8.
38. Brizel DM, Sibley GS, Prosnitz LR, Scher RL, Dewhirst MW. Tumor hypoxia adversely affects the prognosis of carcinoma of the head and neck. *Int J Radiat Oncol Biol Phys* 1997;38:285-9.
39. Cosse JP, Michiels C. Tumor hypoxia affects the responsiveness of cancer cells to chemotherapy and promotes cancer progression. *Anticancer Agents Med Chem* 2008;8:790-7.
40. Nordmark M, Bentzen SM, Rudat V, Brizel D, Lartigau E, Stadler P, et al. Prognostic value of tumor oxygenation in 397 head and neck tumors after primary radiation therapy: an international multi-center study. *Radiother Oncol* 2005;77:18-24.
41. An Y, Kiang A, Lopez JP, Kuo SZ, Yu MA, Abhold EL, et al. Cigarette smoke promotes drug resistance and expansion of cancer stem cell-like side population. *PLoS One* 2012;7:e47919.
42. Simmons VN, Litvin EB, Jacobsen PB, Patel RD, McCaffrey JC, Oliver JA, et al. Predictors of smoking relapse in patients with thoracic cancer or head and neck cancer. *Cancer* 2013;119:1420-7.
43. Zevallos JP, Mallen MJ, Lam CY, Karam-Hage M, Blalock J, Wetter DW, et al. Complications of radiotherapy in laryngopharyngeal cancer: effects of a prospective smoking cessation program. *Cancer* 2009;115:4636-44.
44. Arcavi L, Benowitz NL. Cigarette smoking and infection. *Arch Intern Med* 2004;164:2206-16.
45. Gritz ER. Smoking and smoking cessation in cancer patients. *Br J Addict* 1991;86:549-54.
46. Duffy SA, Terrell JE, Valenstein M, Ronis DL, Copeland LA, Connors M. Effect of smoking, alcohol, and depression on the quality of life of head and neck cancer patients. *Gen Hosp Psychiatry* 2002;24:140-7.
47. Twu CW, Wang WY, Liang WM, Jan JS, Jiang RS, Chao J, et al. Comparison of the prognostic impact of serum anti-EBV antibody and plasma EBV DNA assays in nasopharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 2007;67:130-7.

Cancer Epidemiology, Biomarkers & Prevention

Prognostic Impact of Cigarette Smoking on the Survival of Patients with Established Nasopharyngeal Carcinoma

Pu-Yun OuYang, Zhen Su, Yan-Ping Mao, et al.

Cancer Epidemiol Biomarkers Prev 2013;22:2285-2294. Published OnlineFirst November 19, 2013.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-13-0546](https://doi.org/10.1158/1055-9965.EPI-13-0546)

Supplementary Material Access the most recent supplemental material at:
<http://cebp.aacrjournals.org/content/suppl/2016/04/15/1055-9965.EPI-13-0546.DC1>

Cited articles This article cites 45 articles, 9 of which you can access for free at:
<http://cebp.aacrjournals.org/content/22/12/2285.full#ref-list-1>

Citing articles This article has been cited by 2 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/22/12/2285.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/22/12/2285>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.