
Correction

Correction: Which Risk Model to Use? Clinical Implications of the ACS MRI Screening Guidelines

In this article (Cancer Epidemiol Biomarkers Prev 2013;22:146–9), which was published in the January 2013 issue of *Cancer Epidemiology, Biomarkers & Prevention* (1), the development study for the Tyrer-Cuzick model was incorrectly identified as the Mayo Clinic Benign Breast Disease cohort in the Introduction. The model is correctly identified below. The authors regret the error.

The model is a statistical model based in part on the data obtained from the International Breast Intervention Study (IBIS).

Reference

1. Ozanne EM, Drohan B, Bosinoff P, Semine A, Jellinek M, Cronin C, et al. Which risk model to use? Clinical implications of the ACS MRI screening guidelines. *Cancer Epidemiol Biomarkers Prev* 2013;22:146–9.

Published OnlineFirst May 16, 2013.

doi: 10.1158/1055-9965.EPI-13-0300

©2013 American Association for Cancer Research.

Cancer Epidemiology, Biomarkers & Prevention

Correction: Which Risk Model to Use? Clinical Implications of the ACS MRI Screening Guidelines

Cancer Epidemiol Biomarkers Prev 2013;22:1167. Published OnlineFirst May 16, 2013.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-13-0300](https://doi.org/10.1158/1055-9965.EPI-13-0300)

Cited articles This article cites 1 articles, 1 of which you can access for free at:
<http://cebp.aacrjournals.org/content/22/6/1167.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/22/6/1167>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.