

Hypothesis/Commentary

Follow-up to Abnormal Cancer Screening Tests: Considering the Multilevel Context of Care

Jane M. Zapka¹, Heather M. Edwards², Veronica Chollette³, and Stephen H. Taplin³

Abstract

The call for multilevel interventions to improve the quality of follow-up to abnormal cancer screening has been out for a decade, but published work emphasizes individual approaches, and conceptualizations differ regarding the definition of levels. To investigate the scope and methods being undertaken in this focused area of follow-up to abnormal tests (breast, colon, cervical), we reviewed recent literature and grants (2007–2012) funded by the National Cancer Institute. A structured search yielded 16 grants with varying definitions of "follow-up" (e.g., completion of recommended tests, time to diagnosis); most included minority racial/ethnic group participants. Ten grants concentrated on measurement/intervention development and 13 piloted or tested interventions (categories not mutually exclusive). All studies considered patient-level factors and effects. Although some directed interventions at provider levels, few measured group characteristics and effects of interventions on the providers or levels other than the patient. Multilevel interventions are being proposed, but clarity about endpoints, definition of levels, and measures is needed. The differences in the conceptualization of levels and factors that affect practice need empirical exploration, and we need to measure their salient characteristics to advance our understanding of how context affects cancer care delivery in a changing practice and policy environment. *Cancer Epidemiol Biomarkers Prev*; 23(10); 1965–73. ©2014 AACR.

Introduction

Periodic screening is a well-established strategy for reducing the incidence of cervical and colorectal cancer through the treatment of cancer precursors, and minimizing cancer morbidity and mortality of cervical, breast, and colon cancer (1–4). Although screening rates have generally increased since the 1980s for the three cancers, and cancer morbidity is declining (5–7), challenges remain (5–8). Rates of advanced-disease-at-diagnosis and failure in follow-up of abnormal screening tests persist, particularly in racial minorities and other underserved groups (5–14).

Although there are no national estimates of the proportion of individuals without follow-up after receiving abnormal screening tests, several reviews of studies in specific populations suggest the rates are high, but vary

substantially (15–19). A review by Yabroff and colleagues summarized the prevalence of follow-up after abnormal breast, cervical, and colorectal cancer screening tests (15) and found that the majority of studies showed a failure in follow-up in 25% or more cases; some reported failures in more than 50% of cases. Recent estimates demonstrate a persistent problem such that 17% of abnormal mammographic, 12% of abnormal Pap, and 41% of abnormal fecal occult blood screening tests remain unexamined in specific populations (12, 20, 21). The mammographic study compared follow-up in facilities serving populations with relatively less education and more racial minorities and demonstrated follow-up was lower by 4% to 5% compared with more educated, Caucasian populations.

Considering the importance and challenge of completing follow up to abnormal screening tests, we updated our background literature review and performed an audit of funded NCI proposals to deepen our understanding of relevant work and form the basis for this commentary (22, 23).

Background

To guide intervention strategy development and evaluation, several conceptualizations of levels exist. For example, Mandelblatt and colleagues build on the work by Anderson and Aday to identify patients, primary care providers, and cancer care providers as 3 relevant groups linked by communication and affected by the individual or medical care environment in which they operate (24). Although the research does not explicitly include the idea

¹Department of Public Health Sciences, Medical University of South Carolina, Charleston, South Carolina. ²Clinical Research Directorate/CMRP, Leidos Biomedical Research, Inc., Frederick National Laboratory for Cancer Research, Frederick, Maryland. ³Process of Care Research Branch, Behavioral Research Program, Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, Maryland.

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Jane M. Zapka, Department of Public Health Sciences, Medical University of South Carolina, 135 Cannon Street, Room 302H, Charleston, SC 29425. Phone: 843-876-1604; Fax: 843-876-1126; E-mail: zapka@musc.edu

doi: 10.1158/1055-9965.EPI-14-0454

©2014 American Association for Cancer Research.

of "levels" in its conceptualization, the lead author refers to the patient and provider levels in her conclusion. Bastani and colleagues explicitly call out levels in their conceptualization of factors affecting follow-up and labels them patient, provider, practice, and policy (25). In our conceptualization of the context of care, we build on the ecologic model of Engel and suggest that there is successive nesting of patients within families, within provider teams in organizations, nested within communities that are affected by state- and national-level factors including professional and public policies (26).

Furthermore, we recognize that screening is a process, not a discrete event (27–30), and this provides an opportunity to consider improvements in multiple points along the care process (22, 23, 31). We can achieve higher rates of mortality reduction by assuring that the complete process of additional testing and evaluation subsequent to an abnormal screening test always occurs, particularly among populations with few resources. The question is how to achieve these improvements (22, 32).

Bastani and others have called for considering multilevel interventions (25). The above examples of conceptualizations can all be called multilevel and all assume these levels and their interaction affect the likelihood of follow-up testing after an abnormal screening, albeit in potentially very different ways (23). For example, the age and demographic characteristics of individual patients, how provider teams are organized, whether organizations have direct mail reminders to individuals due for follow-up testing and whether professional policies require direct patient notification of abnormal results are all factors from different levels of our ecologic model of care that may affect follow-up (23). Furthermore, what the follow-up evaluation involves, how the reporting is done (i.e., standard electronic medical records, EMR), and who does the reporting may differ with physician community standards, the reimbursement offered through insurance, and the presence or absence of programs like Medicaid for low-income groups. All these factors affect whether follow-up to abnormal screening occurs, occurs correctly (15) or incompletely or expeditiously (33), and therefore when and whether they reach treatment or are told they do not have cancer.

To address this problem, published intervention research has focused on educational, navigational, and reminder intervention strategies with individual providers or patients. It has been limited in its focus on the effects of such strategies on provider teams or organizations, or the simultaneous effects of organization, provider teams, and individuals on follow-up to abnormal screening (22, 25, 34, 35) despite long-standing calls for such work (25, 36, 37). Published research does not completely reflect the increased awareness in the research community about the multilevel context of care so we also audited funded research to characterize the research progress on multilevel interventions to improve the follow-up of abnormal screening (38).

Identification and Auditing of NCI-Funded Research Related to Follow-up of Abnormal Screening Tests

To assess the state of emerging multilevel research, we conducted a structured review of grant applications funded for at least 1 year by the National Cancer Institute (NCI) from 2007 to 2012. We explored the characteristics of intervention and observational studies. We were particularly interested in whether multilevel interventions were tested, how people defined levels and whether definitions of "follow-up" were clearly specified and measured.

We used NCI's Division of Cancer Control and Population Sciences (DCCPS) Portfolio Management Application (PMA) to identify awarded grants in their first year between October 2007 and April 2012, based on their NIH extramural research funding activity code (i.e., R01, R03, R13, R15, R21, P01, U01; ref. 39). Training grants were excluded. We conducted separate keyword searches of the full text of grant applications to identify projects concerned with follow-up to abnormal cancer screening. Search terms included: (i) cancer screening follow-up, and (ii) follow-up mammogram, or mammography, or digital mammography, or mammogram, or MRI, or magnetic resonance imaging, or Pap and HPV, or human papilloma virus, or Pap, or fecal occult blood test, or colonoscopy, or sigmoidoscopy.

PMA searches yielded 272 funded grants. The grant identification and selection process followed PRISMA guidelines (40) to review abstracts and eliminate duplicates ($n = 57$), and studies that focused on biologic sciences or studies that used the term "follow-up" as a reference to the data collection process of doing baseline and follow-up assessments ($n = 136$) as opposed to our interest in the health care process of follow-up evaluation after an abnormal cancer screening test. The full text of these grant applications were reviewed with 51 rated as ineligible. Sixteen funded proposals were rated eligible for detailed auditing.

A detailed standardized audit form was designed (Supplementary Material). Although this was not a systematic review of findings of research studies, we were aware of the standards for systematic reviews, particularly the standard concerning the itemized steps of developing a protocol (41). The audit form included major design domains for research (later noted in Tables 1 and 2) and each had standardized definition and categories. A pilot coding of 4 proposals by all 4 auditors refined the coding criteria and audit form structure. After this process, the 16 grants were fully independently coded by 2 auditors. Following 4 auditors' independent reviews of their assigned grants, data were entered into an Excel spreadsheet. Any differences in the codes between the 2 assigned authors were noted and resolved by all 4 auditors.

Characteristics of Funded Studies

As reported in Table 1, of the 16 studies, 7 focused on cervical cancer, 4 on breast, and 3 on colon cancers, and 2

Table 1. General study characteristics by screening test type ($n = 16$)^a

	Breast	Colon	Cervical	Multiple	Total
Cancer site	4	3	7	2	16
Mechanism					
R01	1	1	5	1	8
R21	2	2	1	0	5
U01	1	0	0	1	2
R03	0	0	1	0	1
Care type					
Detection	3	3	6	1	13
Diagnosis	4	3	6	2	15
Treatment	0	0	3	0	3
Follow-up criterion					
Time to diagnostic test follow-up	2	0	2	1	5
Completion of recommended additional tests	0	1	3	1	5
Results reported to MD/practice	0	1	0	1	2
Test result reported to patient	0	0	1	1	2
Diagnosis	0	0	1	2	3
Time to diagnosis	0	0	0	1	1
Appropriateness of follow-up	1	1	0	0	2

^aFor some characteristics, the total number may be more than 16 studies, because of multiple responses.

studies considered multiple cancers. The majority of grants with a larger scope (i.e. R01 as compared with R21) focused on cervical cancer tests. The operational definition of "follow-up" was variable, referring to the various steps within the process of care (42). Operational definitions with highest frequency were receipt of recommended additional tests and time to initiation of diagnostic test follow-up. Studies evaluating the appropriateness of follow-up were the minority ($n = 2$), whereas the remainder investigated various steps in the follow-up process (e.g., referral to appropriate subspecialty).

Table 2 summarizes the research methods proposed in the studies. Given that the variable funding mechanisms reflect differing project aims and scopes of work, the research methods were also variable. Considering the research continuum (43, 44), the majority proposed intervention research with 11 testing an intervention strategy. Studies also proposed descriptive epidemiologic research questions about behavioral and organizational factors affecting follow-up, measurement/instrument development and/or tested intervention strategies. With respect to design, the broad categories of study design (45), controlled trials, quasi-experimental designs, and nonexperimental designs were represented. Qualitative, quantitative, and mixed methods of data sources were proposed. Few studies incorporated economic research questions and analyses. A wide variety of models (e.g., the chronic care model) and theories (e.g., social cognitive theory, theory of reasoned action) guided the proposed research. The vast majority of studies focused on impact or outcome measures at the individual patient level, proposing a variety of data sources and statistical methods.

The ages of the populations to be studied reflected the guidelines for a specific screening test. However, there were variations across studies (e.g., ages >18 or 18–64 years for cervical screening). Only one study of colorectal cancer screening included people older than 75 years. Consistent with national goals about health disparities (46), a majority of studies planned to include participants from racial and ethnic populations, as highlighted in Table 2. Interestingly, given the relationship between insurance status and screening behavior (47, 48), many studies ($n = 10$) were unclear whether they planned to investigate the mediating influence of insurance on follow-up of abnormal tests. A variety of settings were proposed, including practice networks, hospital out-patient clinics, community health centers, the Veterans Administration Hospitals, and managed care organizations.

Among the 11 grants that focused on testing the efficacy or effectiveness of interventions, several targeted a practice or practice system level and included strategies such as adding a case manager component to a practice intervention or conducting the trial in a practice-based primary care research network. A small number proposed targeting the community, such as engaging religious congregations to promote the training and use of lay health advisors. Table 3 provides examples of the intervention studies, highlighting the proposed intervention target level, characteristics, strategies, and research methods. The small number and heterogeneity of the studies prevented a simple classification so we have provided examples using broad ecologic model levels of individual, practice, and community.

Table 2. Selected research domains and methods ($n = 16$ studies)^a

Research domains and methods	<i>n</i>
Research continuum	
Prevalence in selected population	7
Behavioral–organizational epidemiology	10
Measurement/instrument development	10
Theory testing/development	6
Intervention design/mapping	6
Piloting of intervention	2
Testing of intervention	11
Dissemination–diffusion planning	3
Economic analyses: (cost of intervention-3, cost of evaluation-1, cost-effectiveness-2)	3
Design	
Experimental	
RCT—Individual level	5
RCT—Group level	3
Quasi-experimental	
Time series	1
Prospective cohort	3
Nonexperimental	
Secondary data analyses	1
Convenience sample descriptive	3
Models	
Chronic care model	3
Ecologic model	2
Precede–Proceed	2
Other—each cited once (e.g., preventive health model, community based approaches model, health belief model, RE-AIM, access to cancer care model)	9
Theories	
Social cognitive/learning model	3
Theory of reasoned action/theory of planned behavior	3
Diffusion theory	2
Other—each cited once (e.g. grounded theory, locus of control, cognitive-social health information-processing model)	8
Data sources	
Qualitative strategies (focus groups and key informant interviews): patients ($n = 10$), clinicians ($n = 5$), and others ($n = 4$).	12
Surveys: patients ($n = 11$), clinicians ($n = 2$)	11
Medical records and summary reports	10
Levels of measures	
Individual patient	16
Lay individual/advisors	2
Health care provider	4
Group (family)	2
Organizations	2
Community	6
Statistical methods	
Qualitative	10

*(Continued in the following column)***Table 2.** Selected research domains and methods ($n = 16$ studies)^a (Cont'd)

Research domains and methods	<i>n</i>
Descriptive	12
Hierarchical	3
Time series	2
Regression	12
Population focus age	
Screening eligible, but variable groups, e.g. 5–79, 16+, depending on screening test	16
Population inclusion—patient race/ethnicity	
Hispanic	10
Asian	6
African American	11
Caucasian	14
American Indian	3
Alaska Native	2
Hawaiian	1
Pacific Asian Islander	2

^aTotal may be >16 studies because of multiple responses.

Observations

Although only a modest number of funded grant applications between 2007 and 2012 addressed issues related to the process of follow-up of abnormal screening tests, the interest in maximizing performance of the steps and interfaces of care to improve outcomes across the continuum of cancer care was evident and growing within the research community (49–53). The studies proposed by the funded grants, however, reflect a traditional focus on the outcomes at the individual level. Although multilevel interventions and measures were proposed, they frequently were operationalized as "process measures" (e.g., clinician referrals to patient navigator) and did not include assessments of the effect of the intervention on providers or provider team functioning. This bi-directional effect is a key concept of multilevel research. Furthermore, few examined characteristics of provider teams or organizations as mediators between the intervention and patient outcome.

We and others argue it is time to conceptualize a new approach to intervention strategy development and measurement (23, 54, 55). This approach would view the problem of follow-up to abnormal screening as a multilevel issue (14, 56) and consider bidirectional interactions. The challenge is to identify potentially modifiable factors among individuals, family members, providers, and/or organizations that can be targeted in interventions that will improve the likelihood of follow-up for the patient (23, 36).

Potential reasons for the limited studies in the research pipeline include several scientific and practical challenges, although we did not evaluate studies that were rejected. Some possible reasons for rejections include the complexity of such interventions (55). Another is that they

Table 3. Nature of intervention ($n = 11$ intervention studies)

Comments	Example
Intervention strategies targeted to the individual level	
Intervention strategies included in-person education ($n = 4$), telephone calls ($n = 4$), mailings ($n = 5$). ^a The majority of interventions ($n = 7$) were delivered or managed by one person, a nurse, or study staff member. Some involved multiple staff, such as a practice manager working with primary care physicians. A few used multiple strategies by nurses, navigators, a lay navigator, social worker, and physician.	One study investigated the performance of follow-up (colposcopy and subsequent medical recommendations) of abnormal Pap and HPV tests using a 3-group study design randomized at the patient level; (i) standard care (SC) plus barriers assessment and counseling delivered via telephone, (ii) SC plus tailored barriers information via home mail, and (iii) SC plus letter notifying patient of abnormal test, the need to undergo colposcopy, an appointment date and telephone appointment confirmation assessment via letter. Numerous patient mediating and moderating measures were included. Patients were 18+ years of age who were tested at a hospital-based women's care center. Intervention was delivered by study staff. The study planned a cost analysis of intervention options.
Interventions strategies targeted at the practice or practice systems level	
Strategies in several studies included information technology innovation. Several considered staff re-design (e.g., a cancer prevention specialist training for interdisciplinary teams, adding a case manager or lay health worker) and staff training (e.g., navigator training women volunteers).	Acknowledging that barriers to screening include patient and provider variables, one study designed practice level changes to be implemented within a primary care practice network. The network used an EMR and established specialist referrals. The intervention was a colorectal cancer screening module using a prospective cohort design. Strategies were designed to supplement the existing medical records; the services of a cancer prevention specialist were added to provide education and facilitate ordering and follow-up (in-person, mailings, and phone calls to home).
Intervention strategies aimed at the community level or community organizations	
No study tried to change community level factors, but a few involved organizations at the community level. Some studies, however, collected measures at the community level for use in the study design.	One study investigated the impact of faith-based lay health advisors on the use of and follow-up to abnormal Pap tests. It used a group randomized (50 religious congregations) experimental design. Formative work included community inventories (e.g., transportation, communication, health resources) to provide information to use in the training of lay advisors who would then talk with women about overcoming barriers.

^aFrequencies may be greater than 16 because some studies studied more than one strategy.

are large and expensive because cluster randomized trials are an appropriate consideration (57). Other potentially appropriate quasi-experimental designs should therefore also be considered (57, 58). Finally, multilevel interventions may be disruptive for teams and organizations being pressed to meet the business demands of medicine.

Because we could not evaluate these potential explanations, we acknowledge the limitations of the structured portfolio audit. We do not have the number (denominator) of all grants that were submitted but not funded, which does not permit comparisons on content and methods of those unfunded proposals. It is also limited in that it is a review of NCI-funded projects so it does not include

the work in other public and private agencies. As always, criteria being audited required interpretation and judgment, some grants may involve multiple research methods, which might be categorized in several ways. We did, however, search not only the abstracts, but the full proposals of the potentially appropriate proposals and also conducted a reliability review for every audit.

Recommendations for Future Research in the Emerging Context of Care

Several priority topics and methodologic recommendations evolve from this review. An important research task is to clearly define "follow-up" to abnormal screening

tests, to allow the comparison of outcomes across studies and refine and test intervention foci and strategies. These definitions could consider multiple measures within the process, for example, time to follow-up test appointment, or completion of all tests as was done in recently reported navigation studies (59). Potential mediating endpoints could be identified and examined discretely such as (i) scheduling the appointment, (ii) initiating the diagnostic work-up, (iii) completing any diagnostic work-up, (iv) completing the recommended diagnostic work-up, (v) reporting results to the referring physician, (vi) reporting results to the patient, and (vii) initiating therapy among those with cancer. The reasons for failures at each of these points could be different and therefore amenable to different intervention strategies.

Explication of the steps and interfaces where breakdowns are most likely to occur, the levels most amenable to intervention, and the most productive interactions among levels will contribute to improving future work to increase follow-up completion. An excellent example is articulated by Sarfaty and colleagues (60). Seven steps were identified in colorectal cancer screening via stool blood test (SBT), including follow-up for positive SBTs. The investigators' work further identified that how the steps occurred varied widely across 15 primary care practices, suggesting need to allow tailoring when testing interventions for practice performance improvement. In addition, different levels and the measures of effect at these levels need explicit operational definition, development of reliable and valid measures, and inclusion in future research and analyses (53, 61).

The majority of the funded studies in the last 5 years proposed familiar intervention strategies. Examples are reminders by various modes such as letters or calls to patients or electronic reminders to clinicians (30), or navigators or case managers to "fix" a challenging process for patients on an individual basis (35, 62). Emerging studies show promise that navigation can improve follow-up of mammography and colorectal cancer tests for the underserved but they have not explored the effect of the navigator on the provider team or organization. Furthermore the effects are not universal, and may occur after 90 days and not change cancer stage (34, 63). For example, providing case management did not promote timely follow-up to abnormal mammography even when the diagnostic care was free (64).

Future research needs to evaluate the effects of such strategies on the provider team functions and organizational systems and structure (57). Multilevel effects may be bi-directional so analyses plans should consider mediators and endpoints at levels other than the person with the abnormal screen. For example, when patients could be referred to a navigator, how many clinicians actually made appropriate referrals? If navigators are added to a practice, how do they interact with providers, and what is their role with patients? Are they incorporated into explicit communication flow? If clinicians are made aware

of community resources to offer, how many patients report they actually discussed and were encouraged to use the resources (65)? End points at the provider level might include acceptance of the navigator, metrics of team functioning, and enthusiasm for the intervention (66). Related considerations include whether multilevel interventions deliver better outcomes than single-level interventions, and whether understanding the multilevel effects will speed adoption (67, 68).

Increasing emphasis has been placed on the need to consider potential adoption and diffusion of efficacious interventions during the early phases of research (69). Despite this emphasis, based on this review of recently funded research, more observational studies (organizational or health systems epidemiology) that clarify the team and organizational and influences that may affect or be affected by follow-up care delivery after an abnormal screening test are needed. The PROSPR initiative is a good example of such work (70).

Given that late-stage diagnoses persist among minority groups, inclusion of them as research participants remains a priority. In addition, concern is emerging about screening appropriateness in the older old age groups, and studies to date rarely include people older than 75 years (71, 72).

Another research need involves cost effectiveness or, at minimum, basic cost analysis. Although studies have assessed the cost effectiveness of screening in reducing mortality, few studies address the cost effectiveness of implementing intervention strategies to improve the screening process (73). Studies are needed to examine which category or combination of multilevel strategies is most effective. A recent analysis found that navigation increased the likelihood of follow-up occurring but did not shorten the time-to completion among those who received it, and cost \$275/completion. Such information will inform realistic planning for subsequent intervention research as well as ultimate diffusion and implementation (58).

The U.S. health system is changing rapidly, bolstered by new legislation such as the Affordable Care Act and the variation in adoption of the Medicaid portion of this act (74). Provisions of the Act that encourage Accountable Care Organizations and Patient Centered Medical Homes prompt further recognition of the need to change practices to promote value and quality care delivery. Although historical reimbursement and professional norms did not encourage integrated care across providers and steps in care, Accountable Care Organizations and Patient Centered Medical Homes offer opportunities—to test follow-up interventions because they incentivize consideration of system designs that affect the processes of care, including organizational relationships, necessary to achieve the diagnostic evaluation and use of interdisciplinary teams (75, 76). The focus on patient-centered care requires competencies at patient, provider, and organizational levels (77). Another important piece of legislation, the Health Information Technology for Economic and Clinical

Health Act (HITECH; ref. 78), turned attention and motivation toward achieving meaningful use of electronic health records including their use in tracking follow-up testing after abnormal screening (79–81) and increased communication between providers caring for a common patient (82). Increased attention has focused on the impact of interventions based on EMRs, for example on reduced time to resolution of abnormal screening tests (83). Simultaneously, however, there are studies that emphasize caution in assuming the validity of EMR measures' data elements given differences in systems and "information overload" for clinicians. These studies stressed the need to account for user behaviors and organizational characteristics (84, 85).

Conclusions

There is a long-standing call to consider the multilevel context of care when considering interventions to improve care quality (25, 37). The changing incentives of Accountable Care Organizations and Patient Center Medical Homes are creating opportunities to undertake such research, the problem of follow-up to abnormal screening tests offers a discrete problem that needs attention. Multilevel interventions are being proposed but clarity about endpoints, definition of levels, and measures is needed. Although several frameworks describe the multilevel context of care they differ with respect to what levels mean. The differences need to be explored empirically and we need to understand the salient characteristics of levels. Therefore, to advance our understanding the effects of context on care delivery, we need more explicit definitions of levels, their measures as covariates, and how they are affected by interventions. Furthermore, the

complexity of the research encourages us to address multilevel problems in discrete areas. Follow-up to abnormal screening is one such area, and we argue here it that it is an underexplored area in which to evaluate the benefits of a multilevel perspective.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Disclaimer

The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. government.

Authors' Contributions

Conception and design: J.M. Zapka, H.M. Edwards, V. Chollette, S.H. Taplin

Development of methodology: J.M. Zapka, H.M. Edwards, S.H. Taplin

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): S.H. Taplin

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J.M. Zapka, H.M. Edwards, V. Chollette, S.H. Taplin

Writing, review, and/or revision of the manuscript: J.M. Zapka, H.M. Edwards, V. Chollette, S.H. Taplin

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): J.M. Zapka, H.M. Edwards, S.H. Taplin

Study supervision: J.M. Zapka

Grant Support

This work was supported in whole or in part with federal funds from the National Cancer Institute, NIH (contract No. HHSN261200800001E, to H.M. Edwards).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received April 24, 2014; revised July 23, 2014; accepted July 23, 2014; published OnlineFirst July 29, 2014.

References

- Smith R, Cokkinides V, Brawley O. Cancer screening in the United States, 2008: a review of current American Cancer Society guidelines and cancer screening issues. *CA Cancer J Clin* 2008;58:161–79.
- U.S. Preventive Services Task Force. Guide to Clinical Preventive Services, 2008: Recommendations of the U.S. Preventive Services Task Force. AHRQ Publication No. 08-05122, September 2008. Agency for Healthcare Research and Quality, Rockville, MD.
- Nicholson FB, Barro JL, Atkin W, Lilford R, Patnick J, Williams CB, et al. Review article: population screening for colorectal cancer. *Aliment Pharmacol Ther* 2005;22:1069–77.
- O'Meara AT. Present standards for cervical cancer screening. *Curr Opin Oncol* 2002;14:505–11.
- Jemal A, Simard EP, Dorell C, Noone AM, Markowitz LE, Kohler B, et al. Annual report to the nation on the status of cancer, 1975–2009, featuring the burden and trends in human papillomavirus (HPV)-associated cancers and HPV vaccination coverage levels. *J Natl Cancer Inst* 2013;105:175–201.
- Centers for Disease Control and Prevention. Cancer screening - United States, 2010. *MMWR Morb Mortal Wkly Rep* 2012;61:41–5.
- Siegel R, DeSantis C, Virgo K, Stein K, Mariotto A, Smith T, et al. Cancer treatment and survivorship statistics, 2012. *CA Cancer J Clin* 2012;62:220–41.
- National Cancer Institute. Cancer Trends Progress Report – 2011/2012 Update; 2012 [cited 2013 Jan 31]. Available from: <http://progressreport.cancer.gov>.
- Abotchie PN, Vernon SW, Du XL. Gender differences in colorectal cancer incidence in the United States, 1975–2006. *J Womens Health (Larchmt)* 2012;21:393–400.
- Fair AM, Wujcik D, Lin JM, Zheng W, Egan KM, Grau AM, et al. Psychosocial determinants of mammography follow-up after receipt of abnormal mammography results in medically underserved women. *J Health Care Poor Underserved* 2010;21:71–94.
- Fiscella K, Humiston S, Hendren S, Winters P, Jean-Pierre P, Idris A, et al. Eliminating disparities in cancer screening and follow-up of abnormal results: what will it take? *J Health Care Poor Underserved* 2011;22:83–100.
- Goldman LE, Walker R, Hubbard R, Kerlikowske K. Timeliness of abnormal screening and diagnostic mammography follow-up at facilities serving vulnerable women. *Med Care* 2013;51:307–14.
- Perez-Stable EJ, Afaible-Munsuz A, Kaplan CP, Pace L, Samayoa C, Somkin C, et al. Factors Influencing time to diagnosis after abnormal mammography results in diverse women. *J Womens Health (Larchmt)* 2013;2:159–166.
- Schoenberg N, Baltisberger J, Bardach S, Dignan M. Perspectives on Pap test follow-up care among rural Appalachian women. *Women Health* 2010;50:580–97.
- Yabroff KR, Washington KS, Leader A, Neilson E, Mandelblatt J. Is the promise of cancer-screening programs being compromised? Quality of follow-up care after abnormal screening results. *Med Care Res Rev* 2003;60:294–331.

16. Lester H, Wilson S. Is default from colposcopy a problem, and if so what can we do? A systematic review of the literature. *Br J Gen Pract* 1999;49:223–9.
17. Marcus AC, Kaplan CP, Crane LA, Berek JS, Bernstein G, Gunning JE, et al. Reducing loss-to-follow-up among women with abnormal Pap smears. Results from a randomized trial testing an intensive follow-up protocol and economic incentives. *Med Care* 1998;36:397–410.
18. Wells KJ, Battaglia TA, Dudley DJ, Garcia R, Greene A, Calhoun E, et al. Patient navigation: state of the art or is it science? *Cancer* 2008; 113:1999–2010.
19. Schofield PE, Butow PN. Towards better communication in cancer care: a framework for developing evidence-based interventions. *Patient Educ Couns* 2004;55:32–9.
20. Eliit L, Krzyzanowska M, Saskin R, Barbera L, Razzaq A, Lofters A, et al. Sociodemographic factors associated with cervical cancer screening and follow-up of abnormal results. *Can Fam Physician* 2012;58:e22–31.
21. Etzioni DA, Yano EM, Rubenstein LV, Lee ML, Ko CY, Brook RH, et al. Measuring the quality of colorectal cancer screening: the importance of follow-up. *Dis Colon Rectum* 2006;49:1002–10.
22. Zapka J, Taplin SH, Anhang Price R, Cranos C, Yabroff R. Factors in quality care—the case of follow-up to abnormal cancer screening tests—problems in the steps and interfaces of care. *J Natl Cancer Inst Monogr* 2010;40:58–71.
23. Taplin SH, Yabroff KR, Zapka J. A multilevel research perspective on cancer care delivery: the example of follow-up to an abnormal mammogram. *Cancer Epidemiol Biomarkers Prev* 2012;21:1709–15.
24. Mandelblatt JS, Yabroff KR, Kerner JF. Equitable access to cancer services: a review of barriers to quality care. *Cancer* 1999;86: 2378–90.
25. Bastani R, Yabroff KR, Myers RE, Glenn B. Interventions to improve follow-up of abnormal findings in cancer screening. *Cancer* 2004;101: 1188–200.
26. Taplin S, Anhang Price R, Edwards H, Foster M, Breslau E, Chollette V, et al. Understanding and influencing multilevel factors across the cancer care continuum. *J Natl Cancer Inst Monogr* 2012;44:2–10.
27. Baron RC, Rimer BK, Coates RJ, Kerner J, Kalra GP, Melillo S, et al. Client-directed interventions to increase community access to breast, cervical, and colorectal cancer screening a systematic review. *Am J Prev Med* 2008;35:S56–66.
28. Baron RC, Rimer BK, Breslow RA, Coates RJ, Kerner J, Melillo S, et al. Client-directed interventions to increase community demand for breast, cervical, and colorectal cancer screening a systematic review. *Am J Prev Med* 2008;35:S34–55.
29. Sabatino SA, Coates RJ, Uhler RJ, Breen N, Tangka F, Shaw KM. Disparities in mammography use among US women aged 40–64 years, by race, ethnicity, income, and health insurance status, 1993 and 2005. *Med Care* 2008;46:692–700.
30. Sabatino SA, Lawrence B, Elder R, Mercer SL, Wilson KM, DeVinney B, et al. Effectiveness of interventions to increase screening for breast, cervical, and colorectal cancers: nine updated systematic reviews for the guide to community preventive services. *Am J Prev Med* 2012;43:97–118.
31. Anhang Price R, Zapka J, Edwards H, Taplin SH. Organizational factors and the cancer screening process. *J Natl Cancer Inst Monogr* 2010;40:38–57.
32. Zapka J, Taplin S, Solberg L, Manos M. A framework for improving the quality of cancer care: the case of breast and cervical cancer screening. *Cancer Epidemiol Biomarkers Prev* 2003;12:4–13.
33. Battaglia TA, Santana MC, Bak S, Gokhale M, Lash TL, Ash AS, et al. Predictors of timely follow-up after abnormal cancer screening among women seeking care at urban community health centers. *Cancer* 2010;116:913–21.
34. Raich PC, Whitley EM, Thorland W, Valverde P, Fairclough D, Program DPNR. Patient navigation improves cancer diagnostic resolution: an individually randomized clinical trial in an underserved population. *Cancer Epidemiol Biomarkers Prev* 2012;21:1629–38.
35. Maxwell AE, Jo AM, Crespi CM, Sudan M, Bastani R. Peer navigation improves diagnostic follow-up after breast cancer screening among Korean American women: results of a randomized trial. *Cancer Causes Control* 2010;21:1931–40.
36. National Research Council. Building a Better Delivery System: A New Engineering/Health Care Partnership. Washington, DC: The National Academies Press; 2005.
37. Ferlie EB, Shortell SM. Improving the quality of health care in the United Kingdom and the United States: a framework for change. *Milbank Q* 2001;79:281–315.
38. Zapka J, Taplin S, Ganz P, Grunfeld E, Sterba K. Multilevel factors affecting quality: examples from the cancer care continuum. *J Natl Cancer Inst Monogr* 2012;44:11–19.
39. National Institutes of Health. Types of grant programs; 2012 [cited 2013 Mar 27]. Available from: http://grants.nih.gov/grants/funding/funding_program.htm.
40. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535.
41. Institutes of Medicine. Finding what works in health care: standards for systematic reviews. Washington, DC: The National Academies Press; 2011.
42. Taplin SH, Clauser S, Rodgers AB, Breslau E, Rayson D. Interfaces across the cancer continuum offer opportunities to improve the process of care. *J Natl Cancer Inst Monogr* 2010;40:104–110.
43. Greenwald P, Cullen JW, McKenna JW, Hiatt RA, Rimer BK. Cancer prevention and control: from research through applications; A new strategy for cancer control research. *J Natl Cancer Inst* 1987;79: 389–400.
44. Hiatt RA, Rimer BK. A new strategy for cancer control research. *Cancer Epidemiol Biomarkers Prev* 1999;8:957–64.
45. Shadish WR, Cook TD, Campbell DT. Experimental and quasi-experimental designs for generalized causal inference. Boston, MA: Houghton Mifflin; 2002.
46. United States Department of Health and Human Services. Healthy people; 2012 [cited 2013 Mar 11]. Available from: <http://www.healthypeople.gov>. Accessed March 11, 2013.
47. ACSAN (American Cancer Society Cancer Action Network). 2009. Cancer disparities: a chartbook. Washington, DC: American Cancer Society Cancer Action Network.
48. Halpern MT, Ward EM, Pavluck AL, Schrag NM, Bian J, Chen AY. Association of insurance status and ethnicity with cancer stage at diagnosis for 12 cancer sites: a retrospective analysis. *Lancet Oncol* 2008;9:222–31.
49. Grunfeld E, Earle C. The interface between primary and oncology specialty care: treatment through survivorship. *J Natl Cancer Inst Monogr* 2010;40:25–30.
50. Han PKJ, Rayson D. The interface between primary and oncology specialty care at the end of life. *J Natl Cancer Inst Monogr* 2010; 40:31–7.
51. Nekhlyudov L, Latosinsky S. The interface between primary oncology specialty care: from symptoms to diagnosis. *J Natl Cancer Inst Monogr* 2010;40:11–7.
52. Sussman J, Baldwin LM. The interface between primary and oncology specialty care: from diagnosis through primary treatment. *J Natl Cancer Inst Monogr* 2010;40:18–24.
53. Taplin SH, Rodgers AB. Toward improving the quality of cancer care: addressing the interfaces of primary and oncology-related subspecialty care. *J Natl Cancer Inst Monogr* 2010;40:3–10.
54. Clauser S, Taplin S, Foster M, Fagan P, Kaluzny A. multilevel intervention research: lessons learned and pathways forward. *J Natl Cancer Inst Monogr* 2012;44:127–133.
55. Weiner B, Lewis M, Clauser S, Stitzenberg K. In search of synergy: strategies for combining interventions at multiple levels. *J Natl Cancer Inst Monogr* 2012;44:34–41.
56. Weingart SN, Saadeh MG, Simchowit B, Gandhi TK, Nekhlyudov L, Studdert DM, et al. Process of care failures in breast cancer diagnosis. *J Gen Intern Med* 2009;24:702–9.
57. Cleary P, Gross C, Zaslavsky A, Taplin S. Multilevel interventions: study design and analysis issues. *J Natl Cancer Inst Monogr* 2012; 44:49–55.
58. Stange K, Breslau E, Dietrich A, Glasgow R. State-of-the-art and future directions in multilevel interventions across the cancer control continuum. *J Natl Cancer Inst Monogr* 2012;44:20–31.

59. Paskett ED, Katz ML, Post DM, Pennell ML, Young GS, Seiber EE, et al. The Ohio Patient Navigation Research Program: does the American Cancer Society patient navigation model improve time to resolution in patients with abnormal screening tests? *Cancer Epidemiol Biomarkers Prev* 2012;16:20–8.
60. Sarfaty M, Myers RE, Harris DM, Borsky AE, Sifri R, Cocroft J, et al. Variation in colorectal cancer screening steps in primary care: basis for practice improvement. *Am J Med Qual* 2012;27:458–66.
61. National Cancer Institute. Grid Enabled Measures Database. n.d. [cited 2013 Feb 12]. Available from: <https://www.gem-beta.org/Public/Home.aspx>.
62. Markossian TW, Darnell JS, Calhoun EA. Follow-up and timeliness after an abnormal cancer screening among underserved, urban women in a patient navigation program. *Cancer Epidemiol Biomarkers Prev* 2012;16:91–700.
63. Freund KM, Battaglia TA, Calhoun E, Darnell JS, Dudley DJ, Fiscella K, et al. Impact of patient navigation on timely cancer care: the patient navigation research program. *J Natl Cancer Inst* 2014;106:dju115.
64. Lobb R, Allen JD, Emmons KM, Ayanian JZ. Timely care after an abnormal mammogram among low-income women in a public breast cancer screening program. *Arch Intern Med* 2010;170:521–8.
65. Koh HK, Brach C, Harris LM, Parchman ML. A proposed 'health literate care model' would constitute a systems approach to improving patients' engagement in care. *Health Aff (Millwood)* 2013;35:357–367.
66. Havyer RD, Wingo MT, Comfere NI, Nelson DR, Halvorsen AJ, McDonald FS, et al. Teamwork assessment in internal medicine: a systematic review of validity evidence and outcomes. *J Gen Intern Med* 2014;29:894–910.
67. Glasgow RE, Marcus AC, Bull SS, Wilson KM. Disseminating effective cancer screening interventions. *Cancer* 2004;101:1239–50.
68. Yano E, Green L, Glanz K, Ayanian J, Mittman B, Chollette V, et al. Implementation and spread of interventions into the multilevel context of routine practice and policy: implications for the cancer care continuum. *J Natl Cancer Inst Monogr* 2012;44:86–99.
69. National Cancer Institute. Process of Care Research Branch (PCRB); 2012 [cited 2013 Feb 12]. Available from: <http://cancercontrol.cancer.gov/brp/pcrb>.
70. Population-based Research Optimizing Screening through Personalized Regimens (PROSPR); 2011 [cited 2013 Jun 5]. Available from: <http://appliedresearch.cancer.gov/networks/prospr>.
71. Goodwin JS, Singh A, Reddy N, Riall TS, Kuo YF. Overuse of screening colonoscopy in the Medicare population. *Arch Intern Med* 2011;171:1335–43.
72. Schenck AP, Peacock SC, Klabunde CN, Lapin P, Coan JF, Brown ML. Trends in colorectal cancer test use in the medicare population, 1998–2005. *Am J Prev Med* 2009;37:1–7.
73. Yabroff KR. Interventions to improve cancer screening: commentary from a health services research perspective. *Am J Prev Med* 2008;35:S6–9.
74. Public Law 111 - 148: Patient Protection and Affordable Care Act.
75. Fennell ML, Prabhu Das I, Clauser S, Petrelli N, Salner A. The organization of multidisciplinary care teams: modeling internal and external influences on cancer care quality. *J Natl Cancer Inst Monogr* 2010;40:72–80.
76. Nutting PA, Crabtree BF, Miller WL, Stange KC, Stewart E, Jaen C. Transforming physician practices to patient-centered medical homes: lessons from the national demonstration project. *Health Aff (Millwood)* 2011;30:439–45.
77. Bernabeo E, Holmboe ES. Patients, providers, and systems need to acquire a specific set of competencies to achieve truly patient-centered care. *Health Aff (Millwood)* 2013;32:250–8.
78. U.S. Department of Health and Human Services. HITECH Act. n.d. [cited 2013 March 27]. Available from: <http://www.healthit.gov/policy-researchers-implementers/hitech-act-0>.
79. Bates DW, Bitton A. The future of health information technology in the patient-centered medical home. *Health Aff (Millwood)* 2010;29:614–21.
80. Hesse BW, Hanna C, Massett HA, Hesse NK. Outside the box: will information technology be a viable intervention to improve the quality of cancer care? *J Natl Cancer Inst Monogr* 2010;40:81–89.
81. Singh H, Wilson L, Petersen LA, Sawhney MK, Reis B, Espadas D, et al. Improving follow-up of abnormal cancer screens using electronic health records: trust but verify test result communication. *BMC Med Inform Decis Mak* 2009;9:49.
82. Centers for Medicare and Medicaid Services. Stage 1 vs. Stage 2 comparison table for eligible professionals; 2012 [cited 2013 Mar 11]. Available from: <https://www.cms.gov/Regulations-and-Guidance/Legislation/EHRIncentivePrograms/Downloads/Stage1vsStage2-CompTablesforEP.pdf>.
83. Dupuis EA, White HF, Newman D, Sobieraj JE, Gokhale M, Freund KM. Tracking abnormal cervical cancer screening: evaluation of an EMR-based intervention. *J Gen Intern Med* 2010;25:575–580.
84. Kern LM, Malhotra S, Barrón Y, Quaresimo J, Dhopeswarkar R, Pichardo M, et al. Accuracy of electronically reported "meaningful use" clinical quality measures: a cross-sectional study. *Ann Intern Med* 2013;158:77–8.
85. Singh H, Spitzmueller C, Petersen NJ, Sawhney MK, Sittig DF. Information overload and missed test results in electronic health record-based settings. *JAMA Intern Med* 2013;173:702–4.

Cancer Epidemiology, Biomarkers & Prevention

Follow-up to Abnormal Cancer Screening Tests: Considering the Multilevel Context of Care

Jane M. Zapka, Heather M. Edwards, Veronica Chollette, et al.

Cancer Epidemiol Biomarkers Prev 2014;23:1965-1973. Published OnlineFirst July 29, 2014.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-14-0454](https://doi.org/10.1158/1055-9965.EPI-14-0454)

**Supplementary
Material** Access the most recent supplemental material at:
<http://cebp.aacrjournals.org/content/suppl/2014/07/31/1055-9965.EPI-14-0454.DC1>

Cited articles This article cites 68 articles, 10 of which you can access for free at:
<http://cebp.aacrjournals.org/content/23/10/1965.full#ref-list-1>

Citing articles This article has been cited by 3 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/23/10/1965.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and
Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department
at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/23/10/1965>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC)
Rightslink site.