

Cancer Research in Asian American, Native Hawaiian, and Pacific Islander Populations: Accelerating Cancer Knowledge by Acknowledging and Leveraging Heterogeneity

Scarlett Lin Gomez^{1,2}, Sally L. Glaser^{1,2}, Pamela L. Horn-Ross^{1,2}, Iona Cheng^{1,2}, Thu Quach^{1,2}, Christina A. Clarke^{1,2}, Peggy Reynolds^{1,2}, Salma Shariff-Marco^{1,2}, Juan Yang¹, Marion M. Lee³, William A. Satariano⁴, and Ann W. Hsing^{1,2}

Abstract

The Asian American, Native Hawaiian, and Pacific Islander population is large, growing, and extremely heterogeneous. Not only do they bear unique burdens of incidence and outcomes for certain cancer types, they exhibit substantial variability in cancer incidence and survival patterns across the ethnic groups. By acknowledging and leveraging this heterogeneity through investing in cancer research within these populations, we have a unique opportunity to accelerate the availability of useful and impactful cancer knowledge.

See all the articles in this CEBP Focus section, "Cancer in Asian and Pacific Islander Populations."

Cancer Epidemiol Biomarkers Prev; 23(11); 2202–5. ©2014 AACR.

Asian Americans, Native Hawaiians, and Pacific Islanders (AANHPI) are collectively the most rapidly growing racial/ethnic group in the United States, recently surpassing Hispanics in rates of population growth (1). In the 2000 Census, AANHPIs were enumerated at 15.8 million individuals (19 million including AANHPIs of multiple race), representing 5.1% or 1 in 20 persons in the United States (2), with the largest populations in California [6.0 million (15.8% of total population)] and New York [1.7 million (8.7% of total population)]. By Census estimates, the number of AANHPIs in the United States will exceed 40 million in 2050 (3). Reflecting recent immigration trends, the proportion of Asian Americans who are foreign born ranges from 28% among Japanese to 70% among Asian Indians and Koreans; 25% of all Asian Americans immigrated within the past decade (4).

AANHPIs include populations from more than 30 different countries, and from strikingly different social and economic backgrounds, speaking more than 100 languages. This diversity is based, in part, on broad historical immigration patterns: Chinese laborers arrived in the United States in the 19th century, Vietnamese refugees in the 1970s, Filipino health professionals starting in the 1970s, and South Asian technology professionals in the

2000s. Furthermore, the many cultures and lifestyles AANHPIs bring from their native countries have been modified to varying degrees through acculturation in the United States. The net result is an aggregate AANHPI population that is highly heterogeneous, varying profoundly in characteristics such as English language proficiency, socioeconomic status, insurance coverage, health beliefs, use of health services, diets, body size, and lifestyles (5, 6). For example, the median household income in 2000 across California AANHPI groups ranged almost 3-fold, from \$24,337 among the Hmong population to \$68,935 among the Asian Indian population (7). The percentage of AANHPIs with limited English proficiency ranged from 12% among Native Hawaiian and Pacific Islanders (NHPI) to 54% among Vietnamese (3). The percentage of AANHPIs with less than high school education ranged from 7% among Japanese to 56% to 68% among Southeast Asians (Cambodians/Kampuchians, Laotians, and Hmong; ref. 3). Within each AANHPI population, access to health care and cancer behavioral risk factors also differ by immigration, acculturation, and socioeconomic status (8, 9). In addition, differences in residential neighborhoods and types of occupation affect environmental exposures (e.g., air pollution, workplace chemicals, and built environment). All these factors affect cancer risk and associated outcomes, and thus have important implications for health.

Despite this diversity, most cancer research still considers AANHPI populations in the aggregate. Aggregation can mask important differences across specific AANHPI populations that largely have been overlooked (10), such as disparities in cancer incidence/risk and outcomes. This practice of aggregation has persisted, in part, because data (e.g., from the Census) for detailed ethnicities are often not readily available. The lack of

¹Cancer Prevention Institute of California, Fremont, California. ²Division of Epidemiology, Department of Health Research and Policy, Stanford University School of Medicine, Stanford, California. ³University of California, San Francisco, San Francisco, California. ⁴University of California, Berkeley, Berkeley, California.

Corresponding Author: Scarlett Lin Gomez, Cancer Prevention Institute of California, 2201 Walnut Avenue, Suite 300, Fremont, CA 94538. Phone: 510-608-5041; Fax: 510-608-5085; E-mail: scarlett@cpic.org

doi: 10.1158/1055-9965.EPI-14-0624

©2014 American Association for Cancer Research.

disaggregated cancer data for specific AANHPI populations also may be a consequence of the persistent stereotype of attributing positive health profiles to the aggregate group as a single "model minority" (10), and thus of the limited awareness of the need and value of examining specific ethnic groups. Early studies that did focus on specific AANHPI populations and their migration patterns revealed clear differences in cancer incidence between Asians living in their native countries and those living in the United States, as well as changes in rates associated with length of residence and number of generations in the United States. These classic studies are often cited together as evidence for the role of environmental factors in cancer etiology (11–15). However, consistent consideration of specific ethnic populations individually has not typified the AANHPI cancer epidemiology literature within the United States. This failure to disaggregate groups is a disservice to public health by allowing potential disparities and vulnerable populations to go undetected (16). It also has resulted in lost opportunities to leverage the tremendous heterogeneity across these populations in the United States for the purpose of uncovering potential group-specific cancer risk and prognostic factors.

In two recently published articles using national SEER data to examine cancer rates over nearly 20 years (1990 through 2008), we demonstrated substantial variation in incidence time trends across eight Asian American ethnic populations and three Native Hawaiian and Pacific Islander populations (17, 18). Such patterns had not been examined previously due to lack of availability of population estimates for specific AANHPI ethnic groups. Using California SEER data, we also have examined cancer incidence and survival patterns among Asian American populations by nativity, or immigration status, revealing previously unidentified disparities, and uncovering patterns pointing to new directions for inquiries into etiology and survivorship (11, 16, 19–22). Such data on contemporary trends and patterns in cancer incidence and outcomes, together with the diversity across these ethnic groups, their exposures, and their immigration patterns, provide significant but underutilized opportunities to make discoveries about the relative contributions of environmental and genetic influences on cancer etiology and outcomes. Moreover, emerging research shows that aspects of the biology of cancers may be unique to some AANHPI ethnic groups; examples include the observed higher proportions of HER2⁺ breast cancers among Filipinas, Koreans, and Vietnamese (23); the higher prevalence of EGFR mutations in lung cancer among East Asians (24); the higher Gleason grade prostate tumors among Chinese, Japanese, foreign-born Filipinos, foreign-born Koreans, and foreign-born Vietnamese (22); and the higher incidence of premenopausal breast cancer among U.S.-born Chinese and Filipino women relative to non-Hispanic whites (11). The incidence patterns and variations noted in these descriptive cancer registry-based studies provide a launching point for deeper inquiries into

factors underlying potentially unique etiology and prognosis profiles in these groups.

This issue of *Cancer Epidemiology, Biomarkers & Prevention* features two editorials and six original articles on AANHPI populations. Together, they (i) provide insights into cancer etiology and cancer outcomes by studying exposures, exposure levels, and gene–environment interactions for specific AANHPI populations; (ii) expose and debunk the "model minority" myth of positive cancer health profiles among AANHPIs as an aggregate by demonstrating variations both in exposures and in diseases/outcomes across the specific groups; (iii) feature methodologically novel ways of studying AANHPIs; and (iv) focus on cancers of unique burden to AANHPI groups.

The article by Thompson and colleagues demonstrates the power of existing data from patients' electronic health records (EHR) for studying patient-, provider-, and system-level factors associated with cancer screening in an insured population of seven AANHPI ethnic groups (25). This article illustrates the heterogeneity in cancer screening adherence and associated factors across multiple AANHPI groups even after accounting for health care access. EHRs will be ever more useful as they become the norm for medical record collection and reporting, although they must be designed to capture patient socio-demographic indicators that are meaningful for planning, surveillance, and research (e.g., detailed race/ethnicity based on self-report, birthplace, language, ancestry, and socioeconomic status). Quach and colleagues combined demographics data from the U.S. Census and air toxics data from the U.S. Environmental Protection Agency to examine average concentrations of hazardous air pollutants in residentially concentrated AANHPI census tracts across California (26). By showing higher air pollutant levels for some tracts with heavily concentrated AANHPI ethnic groups, they highlight potential environmental exposure disparities in ethnic minority neighborhoods.

Pollack and colleagues focus on hepatitis B infection, which in most Asian American groups is a predominant risk factor for hepatocellular carcinoma, a cancer of high burden among many AANHPI populations (27). This article shows that seroprevalence of hepatitis B surface antigen determined in a large New York City screening program varied across foreign-born AANHPIs by ethnic group, ranging from 3% to 15%, as well as by gender, age, and specific place of birth. Park and colleagues, using data from the Multiethnic Cohort (MEC), compared levels of urinary metabolites of 1,3-butadiene among Japanese Americans, Native Hawaiians, and non-Hispanic whites to elucidate differential risks of smoking-associated lung cancers (28). Cheng and colleagues, taking advantage of more than 20 years of cancer incidence data for 10 AANHPI ethnic groups, showed that incidence of adenocarcinoma of the lung has been increasing among Filipino and Korean females and among Chinese men (29); this pattern among Filipino and Korean females

are of particular concern, given the low prevalence of smoking in these populations. These findings indicate that research into risk factors other than smoking is warranted and that AANHPI women are an ideal population in which to conduct such research.

Finally, Nguyen and colleagues offer a commentary on the "model minority" myth and why it persists; a multilevel, historical, and life course context for sociocultural factors that shape health and disease for AANHPI populations; and a forward-looking perspective on the approaches and strategies to advance impactful research and public health policy for these populations while addressing methodologic concerns such as small sample sizes (30). They also present a portfolio analysis of grants from the National Cancer Institute's Division of Cancer Control and Population Sciences (DCCPS), demonstrating the paucity of favorably reviewed and funded grants that focus on AANHPIs. Notably, of the few grants funded, none is focused on cancer etiology.

Not only is federally funded cancer research on AANHPIs sparse, but these groups also are currently underrepresented within the Cancer Epidemiology Cohorts (CEC) of the Epidemiology and Genomics Research Program of the DCCPS. Although Chinese in Shanghai and in Singapore are well represented in the CECs, only in the MEC are AANHPIs in the United States included in high numbers, and even in this cohort, the only specific AANHPI ethnic groups of sufficient size for subgroup-specific analyses are Japanese Americans and Native Hawaiians. The CECs, many with rich biospecimen resources, offer a powerful resource for studying the independent and joint effects on cancer incidence and outcomes of factors over the life course and at multiple levels, from molecular and genetic characteristics to contextual influences. However, without

appropriate representation of specific AANHPI ethnic groups in the CECs, these data are unable to provide information that is relevant and targeted to the growing populations of AANHPIs, and also prevents researchers from capitalizing on the heterogeneity in the AANHPI populations to draw insights into cancer etiology and prognosis/survivorship that will benefit cancer prevention and control efforts in all segments of the population.

The AANHPI populations are diverse, dynamic, and growing in numbers. By acknowledging and leveraging this heterogeneity, we have a unique opportunity to accelerate the availability of useful and impactful cancer knowledge.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: S.L. Gomez, S.L. Glaser, I. Cheng, T. Quach, C.A. Clarke, S. Shariff-Marco, M.M. Lee, W.A. Satariano, A.W. Hsing
Development of methodology: S.L. Gomez, S.L. Glaser, T. Quach, J. Yang, M.M. Lee

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): S.L. Gomez, J. Yang

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): S.L. Gomez, S.L. Glaser, I. Cheng, T. Quach, J. Yang, M.M. Lee

Writing, review, and/or revision of the manuscript: S.L. Gomez, S.L. Glaser, P.L. Horn-Ross, I. Cheng, T. Quach, C.A. Clarke, P. Reynolds, S. Shariff-Marco, J. Yang, M.M. Lee, W.A. Satariano, A.W. Hsing

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): S.L. Gomez, T. Quach

Study supervision: S.L. Gomez

Grant Support

This work was supported by the Stanford Cancer Institute (S.L. Glaser).

Received June 6, 2014; revised August 21, 2014; accepted August 22, 2014; published online November 3, 2014.

References

- Asian American Center for Advancing Justice. A Community of Contrasts - Asian Americans in the United States: 2011. Washington, DC: Asian American Center for Advancing Justice; 2013.
- Annual Estimates of the Resident Population by Sex, Race, and Hispanic Origin for the United States: April 1, 2010 to July 1, 2011 (NC-EST2011-03). National Characteristics: Vintage 2011. Population Estimates. [database on the Internet]. [cited May 6, 2014]. Available from: <http://www.census.gov/popest/data/national/asrh/2011/index.html>.
- United States Census Bureau, U.S. Department of Commerce. Asian/Pacific American Heritage Month: May 2014. Profile American Facts for Features. Washington, DC. April 2014.
- American Community Survey. 2007-2009 3-Year Estimates, S0201: Selected Population Profile in the United States 2011.
- California Health Interview Survey. CHIS 2007 Adult Public Use File [computer file]. Los Angeles, CA: UCLA Center for Health Policy Research; 2009.
- Tseng W, McDonnell DD, Ho W, Lee C, Wong S. Ethnic Health Assessment for Asian Americans, Native Hawaiians and Pacific Islanders in California. Prepared for the California Program on Access to Care (CPAC). UC Berkeley School of Public Health; 2010.
- Asian Pacific American Legal Center of Southern California, Asian Law Caucus, National Asian Pacific American Legal Consortium. The Diverse Face of Asians and Pacific Islanders in California. Asian & Pacific Islander Demographic Profile. Los Angeles, CA; 2005.
- Frisbie WP, Cho Y, Hummer RA. Immigration and the health of Asian and Pacific Islander adults in the United States. *Am J Epidemiol* 2001;153:372-80.
- Gomez SL, Glaser SL, Kelsey JL, Lee MM, Sidney S. Immigration and acculturation in relation to health and health-related risk factors among specific Asian subgroups in a health maintenance organization. *Am J Public Health* 2004;94:1977-84.
- Pew Research Center. The Rise of Asian Americans. Washington, DC; 2012.
- Gomez SL, Quach T, Horn-Ross PL, Pham JT, Cockburn M, Chang ET, et al. Hidden breast cancer disparities in Asian women: disaggregating incidence rates by ethnicity and migrant status. *Am J Public Health* 2010;100 Suppl 1:S125-31.
- Prehn A, Lin S, Clarke C, Packer L, Lum R, Lui S, et al. Cancer incidence in Chinese, Japanese and Filipinos in the US and Asia 1988-1992. Union City, CA: Northern California Cancer Center; 1999.
- Stanford JL, Herrinton LJ, Schwartz SM, Weiss NS. Breast cancer incidence in Asian migrants to the United States and their descendants. *Epidemiology* 1995;6:181-3.
- Ziegler RG, Hoover RN, Pike MC, Hildesheim A, Nomura AM, West DW, et al. Migration patterns and breast cancer risk in Asian-American women. *J Natl Cancer Inst* 1993;85:1819-27.
- Wu AH, Ziegler RG, Pike MC, Nomura AM, West DW, Kolonel LN, et al. Menstrual and reproductive factors and risk of breast cancer in Asian-Americans. *Br J Cancer* 1996;73:680-6.

16. Gomez SL, Clarke CA, Shema SJ, Chang ET, Keegan THM, Glaser SL. Disparities in breast cancer survival among Asian women by ethnicity and immigrant status: a population-based study. *Am J Public Health* 2010;100:861–9.
17. Gomez SL, Noone AM, Lichtensztajn DY, Scoppa S, Gibson JT, Liu L, et al. Cancer incidence trends among Asian American populations in the United States, 1990–2008. *J Natl Cancer Inst* 2013;105:1096–110.
18. Liu L, Noone AM, Gomez SL, Scoppa S, Gibson JT, Lichtensztajn D, et al. Cancer incidence trends among native Hawaiians and other Pacific Islanders in the United States, 1990–2008. *J Natl Cancer Inst* 2013;105:1086–95.
19. Chang ET, Yang J, Alfaro-Velcamp T, So SK, Glaser SL, Gomez SL. Disparities in liver cancer incidence by nativity, acculturation, and socioeconomic status in California Hispanics and Asians. *Cancer Epidemiol Biomarkers Prev* 2010;19:3106–18.
20. Clarke CA, Glaser SL, Gomez SL, Wang SS, Keegan TH, Yang J, et al. Lymphoid malignancies in U.S. Asians: incidence rate differences by birthplace and acculturation. *Cancer Epidemiol Biomarkers Prev* 2011;20:1064–77.
21. Horn-Ross PL, McClure LA, Chang ET, Clarke CA, Keegan TH, Rull RP, et al. Papillary thyroid cancer incidence rates vary significantly by birthplace in Asian American women. *Cancer Causes Control* 2011;22:479–85.
22. Lichtensztajn DY, Gomez SL, Sieh W, Chung BI, Cheng I, Brooks JD. Prostate cancer risk profiles of Asian-American men: disentangling the effects of immigration status and race/ethnicity. *J Urol* 2014;191:952–6.
23. Telli ML, Chang ET, Kurian AW, Keegan TH, McClure LA, Lichtensztajn D, et al. Asian ethnicity and breast cancer subtypes: a study from the California Cancer Registry. *Breast Cancer Res Treat* 2011;127:471–8.
24. Subramanian J, Govindan R. Lung cancer in never smokers: a review. *J Clin Oncol* 2007;25:561–70.
25. Thompson CA, Gomez SL, Chan A, Chan JK, McClellan SR, Chung S, et al. Patient and provider characteristics associated with colorectal, breast, and cervical cancer screening among Asian Americans. *Cancer Epidemiol Biomarkers Prev* 2014;23:2208–17.
26. Quach T, Liu R, Nelson DO, Hurley S, Von Behren J, Hertz A, Reynolds P. Disaggregating data on Asian American and Pacific Islander women to provide new insights on potential exposures to hazardous air pollutants in California. *Cancer Epidemiol Biomarkers Prev* 2014;23:2218–28.
27. Pollack HJ, Kwon SC, Wang SH, Wyatt LC, Trinh-Shevrin C, on behalf of the AAHBP Coalition. Chronic hepatitis B and liver cancer risks among Asian immigrants in New York City: results from a large, community-based screening, evaluation, and treatment program. *Cancer Epidemiol Biomarkers Prev* 2014;23:2229–39.
28. Park CL, Kotapati S, Wilkens LR, Tiirikainen M, Murphy SE, Tretyakova N, Le Marchand L. 1,3-Butadiene exposure and metabolism among Japanese American, Native Hawaiian, and white smokers. *Cancer Epidemiol Biomarkers Prev* 2014;23:2240–9.
29. Cheng I, Le GM, Noone A-M, Gali K, Patel M, Haile RW, et al. Lung cancer incidence trends by histology type among Asian American, Native Hawaiian, and Pacific Islander Populations in the United States, 1990–2010. *Cancer Epidemiol Biomarkers Prev* 2014;23:2250–65.
30. Nguyen AB, Chawla N, Noone A-M, Srinivasan S. Disaggregated data and beyond: future queries in cancer control research. *Cancer Epidemiol Biomarkers Prev* 2014;23:2266–72.

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Cancer Research in Asian American, Native Hawaiian, and Pacific Islander Populations: Accelerating Cancer Knowledge by Acknowledging and Leveraging Heterogeneity

Scarlett Lin Gomez, Sally L. Glaser, Pamela L. Horn-Ross, et al.

Cancer Epidemiol Biomarkers Prev 2014;23:2202-2205.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/23/11/2202>

Cited articles This article cites 21 articles, 9 of which you can access for free at:
<http://cebp.aacrjournals.org/content/23/11/2202.full#ref-list-1>

Citing articles This article has been cited by 2 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/23/11/2202.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/23/11/2202>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.