

Research Article

**Maternal Smoking and DNA Methylation in Newborns:
In Utero Effect or Epigenetic Inheritance?**Bonnie R. Joubert¹, Siri E. Håberg³, Douglas A. Bell¹, Roy M. Nilsen^{4,5}, Stein Emil Vollset^{3,5}, Øivind Midttun⁶, Per Magne Ueland⁵, Michael C. Wu², Wenche Nystad³, Shyamal D. Peddada¹, and Stephanie J. London¹**Abstract**

Background: Maternal smoking in pregnancy is associated with adverse health outcomes in children, including cancers; underlying mechanisms may include epigenetic modifications. Using Illumina's 450K array, we previously identified differential DNA methylation related to maternal smoking during pregnancy at 26 CpG sites (CpGs) in 10 genes in newborn cord bloods from the Norwegian Mother and Child Cohort Study (MoBa). Whether these methylation signals in newborns reflect *in utero* exposure only or possibly epigenetic inheritance of smoking-related modifications is unclear.

Methods: We therefore evaluated the impact of the timing of mother's smoking (before or during pregnancy using cotinine measured at 18 weeks gestation), the father's smoking before conception, and the grandmother's smoking during her pregnancy with the mother on methylation at these 26 CpGs in 1,042 MoBa newborns. We used robust linear regression, adjusting for covariates, applying Bonferroni correction.

Results: The strongest and only statistically significant associations were observed for sustained smoking by the mother during pregnancy through at least gestational week 18 ($P < 1.6 \times 10^{-5}$ for all 26 CpGs). We observed no statistically significant differential methylation due to smoking by the mother before pregnancy or that ceased by week 18, father's smoking before conception, or grandmother's smoking while pregnant with the mother.

Conclusions: Differential methylation at these CpGs in newborns seems to reflect sustained *in utero* exposure rather than epigenetic inheritance.

Impact: Smoking cessation in early pregnancy may negate effects on methylation. Analyses of maternal smoking during pregnancy and offspring health outcomes, including cancer, limited to ever smoking might miss true associations. *Cancer Epidemiol Biomarkers Prev*; 23(6); 1007–17. ©2014 AACR.

Introduction

Maternal smoking during pregnancy is associated with many adverse health outcomes in children including certain cancers such as childhood leukemia, lymphoma, and others (1). Recent evidence suggests that the underlying mechanisms behind detrimental effects of maternal smoking may involve epigenetic modifications such as DNA methylation (2–6). We previously reported associa-

tions between maternal smoking during pregnancy (measured objectively using cotinine in maternal plasma samples taken at gestational week 18) and differential DNA methylation in cord blood from newborns in the Norwegian Mother and Child Cohort Study (MoBa; ref. 3). Using the Illumina Infinium HumanMethylation450 Beadchip (450K), we identified epigenome-wide statistically significant associations at 26 CpGs mapping to 10 genes. The genes included *AHRR* and *CYP1A1* that are key members of the aryl hydrocarbon receptor pathway well known to be involved in biologic response to polycyclic aromatic hydrocarbons in tobacco smoke. We also identified novel genes not previously recognized as playing a role in the response to tobacco smoke including genes involved in development (*GFI1*, *MYO1G*, *CNTNAP2*, and *RUNX1*) and other processes (*HLA-DPB2*, *ENSG00000225718*, *EXT1*, and *TTC7B*). We replicated our findings in an independent U.S. birth cohort. The methylation differences at these 26 CpGs were not seen in the range of cotinine consistent with secondhand smoke exposure of the mother. Of note, differential methylation in *AHRR*, *GFI1*, *MYO1G*, and *CNTNAP2* has also been associated with smoking in adults (7–10). Methylation differences in

Authors' Affiliations: ¹Division of Intramural Research, National Institute of Environmental Health Sciences (NIEHS), National Institutes of Health, Department of Health and Human Services, Research Triangle Park, North Carolina; ²Fred Hutchinson Cancer Research Center, Seattle, Washington; ³Norwegian Institute of Public Health, Oslo; ⁴Haukeland University Hospital; ⁵University of Bergen; and ⁶Bevital A/S, Laboratoriebygget, Bergen, Norway

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Stephanie J. London, NIEHS, PO Box 12233, 111 T.W. Alexander Drive, Building 101, Research Triangle Park, NC 27709. Phone: 919-541-5772; Fax: 919-541-2511; E-mail: london2@niehs.nih.gov

doi: 10.1158/1055-9965.EPI-13-1256

©2014 American Association for Cancer Research.

AHRR related to smoking in adults have been observed in lung as well as blood (9), confirming that these findings do not reflect shifts in cell types due to smoking.

Multigenerational health effects from *in utero* exposure to smoking and other toxicants have been proposed in a few epidemiologic studies (11, 12). Our observation that DNA methylation at birth at these 26 CpGs is related to having a mother who smoked during pregnancy raises several questions about when and how these changes might occur. There is considerable interest in the possibility that environmental exposures such as smoking result in epigenetic effects that can be transmitted from one generation to the next but there is no direct evidence in humans (13). This mechanism has been referred to as transgenerational epigenetic inheritance (via the gametes; ref. 14) and implies epigenetic alterations to gametes that escape reprogramming after fertilization.

One possible scenario for the inheritance of smoking-related methylation is that the mother's smoking before becoming pregnant affects the epigenome of the ovum that gives rise to the child. If this occurs, we might observe that offspring methylation is associated with the mother's smoking before pregnancy, even if the mother stopped smoking before becoming pregnant. In a similar way, the father's smoking before conception could affect the epigenome of the sperm that gave rise to the study child. If so, we might observe that offspring methylation is associated with smoking by the father before conception.

Another scenario for the inheritance of the smoking-related methylation changes could be that smoking by the study child's grandmother, when she was pregnant with the mother, affects the epigenome of the developing ovum that gave rise to the study child. If so, we would expect the grandmother's smoking while pregnant with the mother to be associated with the study child's methylation at birth, independently of the mother's smoking during her pregnancy.

Alternatively, it is possible that the methylation differences at birth related to maternal smoking primarily reflect the *in utero* exposure. If so, it is relevant to ask whether early exposure (smoking very early in pregnancy followed by cessation) is sufficient or whether sustained exposure through pregnancy is needed.

To address these questions, we performed new statistical analyses for the 26 CpGs in which we observed methylation differences at birth related to maternal smoking during pregnancy. We examined the impact of the timing of the mother's smoking (before or during pregnancy), the father's smoking before conception, and the grandmother's smoking in her pregnancy with the mother on methylation at these 26 CpGs.

Materials and Methods

Study population

Participants in the current analysis were selected from a substudy of the Norwegian MoBa (15, 16) that evaluated

the association between maternal plasma folate during pregnancy and childhood asthma status at 3 years of age (17). Umbilical cord blood samples were collected and frozen at birth at -80°C and maternal plasma samples were collected at approximately gestational week 18. All biologic material was obtained from the biobank of the MoBa study (16). DNA methylation in cord blood was measured in 1,068 singleton births using the Illumina Infinium HumanMethylation450 Beadchip (450K). We previously analyzed 1,062 of these participants who had complete data for maternal plasma cotinine and covariates to evaluate the association between maternal smoking during pregnancy and DNA methylation in cord blood (3). In the current study, we analyzed the 1,042 participants who had DNA methylation measurements and data for maternal cotinine, self-reported smoking behavior before and during pregnancy, and covariates. Smoking information for the grandmother was reported as unknown or missing for 114 participants, leaving 928 of 1,042 for analyses with that variable. Analyses of father's smoking information included 1,035 participants with data for that variable (7 missing). The MoBa study has been approved by the Regional Committee for Ethics in Medical Research, the Norwegian Data Inspectorate, and the Institutional Review Board of the National Institute of Environmental Health Sciences, North Carolina, and written informed consent was provided by all participants.

Methylation measurements

Details of the 450K methylation measurements and quality control were previously published (3) and are described in detail in the Supplementary Material of that article. Briefly, bisulfite conversion was performed using the EZ-96 DNA Methylation Kit (Zymo Research Corporation) and DNA methylation was measured at 485,577 CpGs in cord blood using Illumina's Infinium HumanMethylation450 BeadChip (18, 19). Illumina's GenomeStudio Methylation module version 1.0 (Illumina Inc.) was used to calculate the methylation level at each CpG as the β value [$\beta = \text{intensity of the methylated allele (M)} / (\text{intensity of the unmethylated allele, U} + \text{intensity of the methylated allele, M} + 100)$; ref. 18]. The laboratory analysis was designed to minimize potential batch effects; the bisulfite conversion and methylation measurements including reruns were completed in less than 1 month on a single machine. Variables representing chip (12 samples), chip set (four contiguous chips or half of a plate), and plate (96 samples) included as covariates in statistical models did not influence the results and thus were not included in the final models.

For the current analysis, we present results accounting for the two different probe designs by applying the intra-array normalization strategy Beta Mixture Quantile dilation (BMIQ; ref. 20). The 26 CpGs evaluated in this analysis did not include any underlying single-nucleotide polymorphisms in the probe sequence as detailed previously in the Supplementary Material (3).

Timing of mother's smoking

Information about smoking by the mother, father, and grandmother was reported by the mother on questionnaires completed at different time points in pregnancy (15, 16; Supplementary Fig. S1). For the mother, cotinine, a biomarker of smoking, was measured by liquid chromatography–tandem mass spectrometry (LC/MS-MS; ref. 21) in plasma collected at approximately gestational week 18. Cotinine values above 56.8 nmol/L were used to indicate that a mother was smoking at this time point (22). We classified mother's smoking into four categories using her report of smoking during pregnancy, and cotinine values: never smoked, quit before pregnancy, smoked during pregnancy but quit by 18 weeks, and smoked through gestational week 18. Quitting by 18 weeks was defined by mother's report plus having a cotinine value below 56.8 nmol/L.

Father's smoking

Father's smoking is an important source of second-hand tobacco smoke exposure to the mother. However, in our previous analysis, maternal cotinine levels consistent with secondhand smoke exposure alone (0 to 56.8 nmol/L) were not associated with differential methylation at the 26 CpGs we evaluate here (3). Nonetheless, the father's smoking before pregnancy could possibly influence methylation in the sperm that could be passed to the offspring. We classified the father's smoking before pregnancy using the mother's response to the question, "Did the baby's father smoke before you became pregnant?"

Grandmother's smoking and combined mother's and grandmother's smoking

The grandmother's smoking was determined by the mother's response to the following question on a questionnaire administered in early pregnancy, "Did your mother smoke when she was pregnant with you?" The response choices were "Yes," "No," or "Don't know." Mother's report of the grandmother's smoking during pregnancy has been validated in a previous publication reporting an association between the grandmother smoking in her pregnancy with the mother and a lower birth weight of the mother (23). We created a categorical variable to jointly classify mother and grandmother smoking during pregnancy into four groups: neither smoked, only grandmother smoked in her pregnancy, only mother smoked in her pregnancy, and both smoked in their pregnancies. For this variable, the mother's smoking in pregnancy was determined solely based on a cotinine value above 56.8 nmol/L.

Statistical analysis

As in our previous publication (3), we used robust linear regression to account for potential outliers or heteroskedasticity (24). However, in the current analysis, rather than logratios, we used the actual methylation β values because the results were nearly identical and

coefficients are easily interpretable as the incremental change in methylation at each probe between the categories compared.

We ran four separate models to examine the association between each smoking variable and methylation in newborns for the 26 CpGs we previously found to be associated with maternal smoking in pregnancy at Bonferroni-corrected epigenome-wide statistical significance (480,000 tests P value $< 1 \times 10^{-7}$). The smoking variables evaluated were the timing of the mother's smoking (quit before pregnancy, quit by 18 weeks, or smoked through 18 weeks relative to never smoked), father smoking before the pregnancy (yes compared with no), and the combined mother and grandmother smoking (mother only, grandmother only, or both smoked relative to neither smoking). To further tease apart the effects of grandmother and mother smoking, we also compared smoking by both grandmother and mother in pregnancy with smoking only by the mother.

All models were adjusted for maternal age, maternal education, and parity. For the analysis of father's smoking before the mother's pregnancy, we additionally adjusted for whether the mother was smoking during pregnancy. The number of cigarettes per day that the mother reported smoking during pregnancy was not a confounder and not included in the final models. Sex of the child would not be expected to be associated with smoking behavior of the mother, father, or grandmother before the birth of the child and also did not affect results so was not included. After our previous publication, a method was published to evaluate potential confounding by differential cell counts in whole blood (25). A reference dataset of cord blood, which would be most applicable to our data, is not available. Therefore, we used the reference dataset of 6 adult men (26) to implement this method (25) and adjustment for estimated cell counts did not alter our results so we present results without this adjustment. We applied Bonferroni correction for 26 tests (26 CpGs evaluated) adjusting the level of significance from 0.05 to 0.0019. We also note the CpGs considered statistically significant after additional Bonferroni correction for the four models run, adjusting the level of significance to $0.05/(26 \times 4) = 0.00048$. These statistical analyses were performed using R (27).

Results

Study population

Of the 1,042 mothers, 50% had never smoked, 22% quit before pregnancy, 15% quit early in pregnancy, and 13% smoked through gestational week 18 (Table 1). Among the mothers who quit by 18 weeks, 80% reported quitting by 6 weeks or earlier and 95% reported that they quit by 10 weeks or earlier. Father's smoking before pregnancy was reported for 31% of study newborns (Table 1). The frequencies for the combined classification of grandmother and mother smoking are also reported in Table 1.

Table 1. Descriptive characteristics of the study population^a

Variable	Category	N (%)
Timing of mother's smoking during pregnancy ^b	Never	520 (49.9)
	Quit before pregnancy	230 (22.1)
	Quit during pregnancy by 18 weeks	156 (15.0)
	Smoked through gestational week 18	136 (13.1)
Combined grandmother's and mother's smoking during their pregnancies ^c (N = 928)	Neither grandmother nor mother smoked	607 (65.4)
	Only grandmother smoked	204 (22.0)
	Only mother smoked	57 (6.1)
	Both grandmother and mother smoked	60 (6.5)
Father's smoking before the mother's pregnancy (N = 1,035)	Yes	317 (30.6)
	No	718 (69.4)
Sex of the child	Male	556 (53.4)
	Female	486 (46.6)
Maternal age	<25	124 (11.9)
	25–30	504 (48.4)
	>30	414 (39.7)
Maternal education	Less than high school	75 (7.2)
	High school degree	336 (32.2)
	Some college	464 (44.5)
	4 years of college or more	167 (16.0)
Parity	0	439 (42.1)
	1	425 (40.8)
	2	135 (13.0)
	3+	43 (4.1)

^aN = 1,042 individuals in the study population. Grandmother's smoking missing for 114 and father's smoking missing for 7 individuals.

^bDetermined by mother's self-report and mother's plasma cotinine measured during pregnancy at approximately gestational week 18, where cotinine values above 56.8 nmol/L indicate smoking.

^c"Their pregnancies" reflects the grandmother's pregnancy with the study mother and the mother's pregnancy with the study newborn whose cord blood DNA methylation we measured.

The results for models unadjusted and adjusted for covariates were very similar. Thus, we present only the adjusted model results.

Timing of the mother's smoking

Figure 1 shows graphically that the largest differences in mean methylation level corresponded to having a mother who smoked through gestational week 18, relative to never smoking, whereas the mean methylation differed very little between never, quit before, and quit during pregnancy smoking categories.

Table 2 and the Supplementary volcano plot (Supplementary Fig. S2) provide the model results for the analysis of methylation differences across the timing of mother smoking categories. Relative to never smoking, former smoking by the mother (quitting before pregnancy) and smoking in pregnancy that stopped by 18 weeks had minimal effects on cord blood methylation that were not statistically significant for any of the 26 CpGs evaluated (Table 2). In contrast, smoking by the mother through at least gestational week 18 had a much stronger association with methylation, relative to never smoking, with regression coefficients ranging from

–0.149 to 0.084. Relative to never smoking, the median regression coefficient (across all 26 CpGs) for smoking through gestational week 18 was 4-fold higher than the median regression coefficient for smoking that stopped by week 18 and 8-fold higher than the median regression coefficient for stopping before pregnancy. Only smoking through 18 weeks in pregnancy was statistically significantly associated with differential methylation in cord blood (P values $< 1.64 \times 10^{-4}$ for all 26 CpGs, Table 2, Supplementary Fig. S2). These associations retain statistical significance if additional Bonferroni correction for four models run is applied (P value $< 4.8 \times 10^{-4}$). As expected, our most statistically significant finding from our epigenome-wide analysis (3), *AHRR* cg05575921, was one of the most statistically significant findings for smoking through gestational week 18 relative to never smoking (regression coefficient = –0.074, SE = 0.008, P value = 9.70×10^{-22}). All of the methylation differences due to smoking are in the same direction as in our original report (3). Models including an additional adjustment for the amount of cigarettes smoked per day reported by the mother in pregnancy (17-week questionnaire) gave similar results.




Figure 1. Mean DNA methylation level for each probe by timing of mother smoking (before and during pregnancy) category. For clarity, only CpGs in genes with multiple CpGs showing consistent direction of effect are displayed (*AHRR* cg23067299 is not displayed).

Father's smoking

We did not observe any statistically significant differences in DNA methylation related to smoking by the father (Supplementary Table S1). Effect estimates were small (coefficients ranged from -0.011 to 0.009).

Grandmother's smoking during pregnancy with the mother alone and in combination with the mother's smoking during pregnancy

Figure 2 shows graphically that the largest differences in the mean methylation level occurred when only the mother smoked, relative to either only the grandmother smoked or neither the mother nor grandmother smoked. In comparison, the mean methylation levels differed very little between neither and grandmother only, and between mother only and both grandmother and mother smoking during pregnancy categories.

Table 3 and the Supplementary volcano plot (Supplementary Fig. S3) display the model results for the analysis of methylation differences across the combined mother and grandmother smoking categories. We observed no statistically significant association for grandmother's smoking alone, relative to no smoking by either the grandmother or the mother, with methylation at any

of the 26 CpGs (Table 3, Supplementary Fig. S3) and the effect sizes were small (ranging from -0.007 to 0.004 ; Table 3). Much larger effect sizes (regression coefficients ranging from -0.137 to 0.075) were observed for only the mother smoking relative to no smoking by either the grandmother or the mother (Table 3). Relative to no smoking by either the grandmother or the mother, the median regression coefficient (across all 26 CpGs) for smoking by the mother only was 12-fold higher than the median regression coefficient for smoking by the grandmother only. The associations remain statistically significant for 23 of the 26 CpGs if additional Bonferroni correction for four models run is applied (P value $< 4.8 \times 10^{-4}$). For the combined grandmother and mother smoking analyses, additional adjustment for the amount of cigarettes smoked per day reported by the mother in pregnancy (17-week questionnaire) gave similar results.

We also compared the effects of both the grandmother and mother smoking with only the mother smoking. If the mother smoked, the additional effect of grandmother smoking in her pregnancy with the mother was minimal and not statistically significant (Supplementary Table S2 and Supplementary Fig. S4).

Table 2. Differential methylation in cord blood DNA in relation to the timing of the mother's smoking^a

Chr ^b	Gene	Distance to gene ^c	CpG	Position ^d	Quit before pregnancy vs. never smokers			Quit during pregnancy by 18 weeks vs. never smokers			Smoked through gestational week 18 vs. never smokers		
					Coef ^e	SE ^f	P	Coef	SE	P	Coef	SE	P
1	GFI1	3,688	cg10399789	92945668	0.001	0.004	0.867	-0.008	0.005	0.091	-0.041	0.006	2.19E-10
1	GFI1	3,224	cg09662411	92946132	0.002	0.005	0.654	-0.017	0.007	0.016	-0.082	0.009	7.47E-19
1	GFI1	3,169	cg06338710	92946187	-0.002	0.004	0.581	-0.011	0.006	0.046	-0.063	0.009	2.02E-13
1	GFI1	2,656	cg18146737	92946700	0.005	0.006	0.439	-0.021	0.008	0.011	-0.117	0.012	4.21E-23
1	GFI1	2,531	cg12876356	92946825	0.006	0.007	0.415	-0.020	0.010	0.046	-0.134	0.012	1.62E-27
1	GFI1	2,321	cg18316974	92947035	0.001	0.003	0.810	-0.010	0.004	0.016	-0.071	0.008	2.93E-18
1	GFI1	1,768	cg09935388	92947588	0.010	0.008	0.227	-0.025	0.010	0.014	-0.149	0.013	7.74E-32
1	GFI1	1,395	cg14179389	92947961	0.004	0.007	0.568	-0.011	0.008	0.155	-0.087	0.007	2.07E-35
5	AHRR	19,617	cg23067299	323907	-0.002	0.004	0.569	0.000	0.004	1.000	0.030	0.005	1.77E-09
5	AHRR	64,157	cg03991871	368447	-0.005	0.002	0.035	0.000	0.003	0.906	-0.021	0.003	4.23E-10
5	AHRR	69,088	cg05575921	373378	-0.003	0.003	0.343	-0.004	0.004	0.302	-0.074	0.008	9.70E-22
5	AHRR	95,070	cg21161138	399360	-0.006	0.003	0.033	-0.004	0.004	0.320	-0.031	0.004	9.79E-13
6	HLA-DPB2	11,549	cg11715943	33091841	-0.003	0.003	0.411	0.003	0.004	0.364	-0.019	0.004	2.99E-06
7	MYO1G	16,417	cg19089201	45002287	-0.002	0.002	0.473	0.001	0.002	0.562	0.013	0.002	4.08E-08
7	MYO1G	16,218	cg22132788	45002486	-0.002	0.003	0.509	0.002	0.003	0.589	0.023	0.003	3.80E-14
7	MYO1G	15,968	cg04180046	45002736	-0.001	0.004	0.747	0.001	0.005	0.896	0.061	0.007	1.39E-19
7	MYO1G	15,785	cg12803068	45002919	-0.002	0.008	0.768	0.007	0.008	0.364	0.084	0.009	2.09E-20
7	ENSG00000225718	198,306	cg04598670	68697651	-0.007	0.006	0.236	-0.014	0.008	0.076	-0.045	0.007	8.88E-10
7	CNTNAP2	854	cg25949550	145814306	0.001	0.001	0.327	-0.002	0.001	0.129	-0.016	0.001	1.01E-39
8	EXT1	-33,821	cg03346806	119157879	-0.004	0.003	0.214	0.004	0.004	0.322	-0.014	0.004	1.64E-04
14	TTC7B	274,756	cg18655025	91008005	-0.002	0.002	0.266	0.002	0.002	0.297	-0.011	0.003	5.31E-05
15	CYP1A1	-1,266	cg05549655	75019143	-0.004	0.004	0.272	-0.004	0.004	0.338	0.036	0.006	2.63E-10
15	CYP1A1	-1,374	cg22549041	75019251	-0.012	0.008	0.156	-0.011	0.010	0.278	0.065	0.011	1.57E-08
15	CYP1A1	-1,406	cg11924019	75019283	-0.004	0.005	0.440	-0.005	0.006	0.433	0.039	0.006	1.43E-09
15	CYP1A1	-1,425	cg18092474	75019302	-0.007	0.007	0.336	-0.011	0.009	0.238	0.060	0.010	2.05E-09
21	RUNX1	1,746	cg12477880	36259241	-0.001	0.004	0.838	0.004	0.006	0.528	0.038	0.008	9.25E-07

^aN = 1,042 individuals analyzed. Categorized using a combination of mother's self-report and mother's plasma cotinine measured during pregnancy at approximately gestational week 18, where cotinine values above 56.8 nmol/L indicate smoking. All analyses adjusted for maternal age, maternal education, and parity.

^bChromosome.

^cDistance (nucleotides) from CpG to transcription start site of the nearest gene.

^dChromosomal position based on National Center for Biotechnology Information human reference genome assembly Build 37.3.

^eRegression coefficient.

^fStandard error for regression coefficient. CpGs with *P* values reaching a Bonferroni-corrected statistical significance threshold of 0.05/26 = 0.0019 are noted in bold.




Figure 2. Mean DNA methylation level for each probe by the combined grandmother and mother smoking category. For clarity, only CpGs in genes with multiple CpGs showing consistent direction of effect are displayed (*AHRR* cg23067299 is not displayed).

Discussion

We recently reported effects of maternal smoking during pregnancy on DNA methylation in newborn cord blood at epigenome-wide statistical significance (P value $< 1 \times 10^{-7}$) for 26 CpGs across 10 loci using the Illumina 450K array (3). In the current article, we extend our analysis to investigate fundamental questions in epigenetics: inheritance and persistence of exposure effects. We looked for evidence of epigenetic inheritance by evaluating the impact of the mother's and father's smoking before pregnancy and the maternal grandmother's smoking in her pregnancy with the mother and on DNA methylation in newborn cord blood. Each of these exposure conditions might potentially alter methylation status of ova or sperm and if effects persisted through fertilization and embryonic development, these might be detected in the cord blood. Our findings do not support epigenetic inheritance. Rather, the methylation differences we observed at birth in relation to maternal smoking seem to reflect *in utero* exposure.

We found that the effects of *in utero* exposure on newborn methylation at these 26 loci were much stronger when the mother smoked past 18 weeks in pregnancy

than when she quit earlier in pregnancy. Methylation at these loci in newborns with mothers who quit smoking earlier in pregnancy was nearly indistinguishable from those whose mothers never smoked. Thus, sustained exposure through at least 18 weeks in pregnancy seems to be required to observe the effects on DNA methylation at birth. Our findings in newborns are in agreement with results from studies of adults, suggesting that smoking-related epigenetic effects are stronger for current than former smoking (9, 10).

The prevalence of smoking in our study population is comparable with the larger MoBa population. Kvalvik and colleagues measured maternal plasma cotinine concentrations at approximately gestational week 18 in a larger sample of 2,997 women in the MoBa study which partially overlaps with the current sample (28). The prevalence of daily or occasional smoking during pregnancy in this larger sample based on self-report and cotinine concentrations > 30 nmol/L is 13%, the same as in our smaller population. A ban on smoking in restaurants and bars went into effect in Norway in June 2004, resulting in decreased prevalence of smoking (29). Among all 19,140 MoBa women delivering singleton births between

Table 3. Differential methylation in cord blood DNA in relation to the combined grandmother's and mother's smoking during their pregnancies^a

Chr ^b	Gene	CpG	Only grandmother smoked in her pregnancy vs. neither smoked			Only mother smoked in her pregnancy vs. neither smoked			Both grandmother and mother smoked in their pregnancies vs. neither smoked		
			Coef ^c	SE ^d	P	Coef	SE	P	Coef	SE	P
1	<i>GFI1</i>	cg10399789	-0.004	0.004	0.339	-0.042	0.009	5.47E-06	-0.039	0.010	1.14E-04
1	<i>GFI1</i>	cg09662411	-0.001	0.006	0.824	-0.075	0.012	6.89E-10	-0.082	0.013	5.01E-10
1	<i>GFI1</i>	cg06338710	-0.005	0.004	0.260	-0.056	0.010	7.61E-08	-0.065	0.013	6.84E-07
1	<i>GFI1</i>	cg18146737	-0.003	0.007	0.624	-0.104	0.017	4.70E-10	-0.117	0.017	1.50E-11
1	<i>GFI1</i>	cg12876356	-0.006	0.008	0.438	-0.127	0.017	1.45E-14	-0.133	0.019	2.90E-12
1	<i>GFI1</i>	cg18316974	-0.003	0.003	0.444	-0.067	0.010	1.66E-10	-0.070	0.014	2.91E-07
1	<i>GFI1</i>	cg09935388	-0.003	0.008	0.692	-0.137	0.016	1.79E-18	-0.147	0.019	2.48E-14
1	<i>GFI1</i>	cg14179389	0.004	0.006	0.495	-0.083	0.010	3.29E-18	-0.079	0.010	1.45E-16
5	<i>AHRR</i>	cg23067299	-0.003	0.004	0.450	0.025	0.007	2.94E-04	0.025	0.006	6.61E-05
5	<i>AHRR</i>	cg03991871	-0.004	0.002	0.116	-0.021	0.005	2.18E-05	-0.020	0.005	3.71E-05
5	<i>AHRR</i>	cg05575921	-0.002	0.003	0.514	-0.057	0.011	2.67E-07	-0.074	0.009	1.18E-16
5	<i>AHRR</i>	cg21161138	-0.001	0.003	0.777	-0.026	0.005	1.70E-06	-0.027	0.007	1.29E-04
6	<i>HLA-DPB2</i>	cg11715943	-0.007	0.003	0.028	-0.019	0.006	9.76E-04	-0.018	0.005	4.85E-04
7	<i>MYO1G</i>	cg19089201	-0.002	0.002	0.337	0.010	0.003	6.78E-04	0.015	0.003	1.00E-06
7	<i>MYO1G</i>	cg22132788	-0.0002	0.003	0.935	0.021	0.004	2.70E-08	0.023	0.004	4.17E-09
7	<i>MYO1G</i>	cg04180046	0.001	0.004	0.882	0.050	0.009	2.74E-08	0.062	0.008	4.61E-14
7	<i>MYO1G</i>	cg12803068	0.003	0.007	0.685	0.071	0.012	7.89E-09	0.086	0.012	4.63E-13
7	<i>ENSG00000225718</i>	cg04598670	-0.00004	0.006	0.995	-0.038	0.009	4.05E-05	-0.036	0.011	9.81E-04
7	<i>CNTNAP2</i>	cg25949550	-0.001	0.001	0.600	-0.014	0.002	1.31E-17	-0.016	0.002	4.30E-23
8	<i>EXT1</i>	cg03346806	-0.001	0.003	0.732	-0.022	0.005	6.19E-06	-0.007	0.006	1.95E-01
14	<i>TTC7B</i>	cg18655025	-0.002	0.002	0.388	-0.014	0.004	2.83E-04	-0.007	0.004	8.20E-02
15	<i>CYP1A1</i>	cg05549655	0.001	0.004	0.838	0.043	0.008	3.41E-08	0.034	0.009	1.08E-04
15	<i>CYP1A1</i>	cg22549041	-0.007	0.008	0.436	0.075	0.014	1.57E-07	0.057	0.018	1.50E-03
15	<i>CYP1A1</i>	cg11924019	0.001	0.005	0.789	0.043	0.007	4.37E-09	0.040	0.011	1.39E-04
15	<i>CYP1A1</i>	cg18092474	0.001	0.007	0.877	0.068	0.012	4.79E-09	0.059	0.017	4.51E-04
21	<i>RUNX1</i>	cg12477880	-0.002	0.004	0.700	0.026	0.010	1.17E-02	0.054	0.011	1.16E-06

^aN = 928 individuals analyzed. "Their pregnancies" indicates the grandmother's pregnancy with the study mother and the mother's pregnancy with the study newborn whose cord blood DNA methylation we measured. All analyses adjusted for maternal age, maternal education, and parity.

^bChromosome.

^cRegression coefficient.

^dStandard error for regression coefficient. CpGs with P values reaching a Bonferroni-corrected statistical significance threshold of 0.05/26 = 0.0019 are noted in bold.

2002 and 2004 (the date of birth range for our sample), smoking in pregnancy was reported by approximately 11% of mothers. However, cotinine was not available for most participants and Kvalvik and colleagues (30) found that the smoking proportion incorporating cotinine is slightly higher than that obtained from self-report in MoBa. The prevalence of grandmaternal smoking (23% grandmother only) and paternal smoking before conception (31%) in these 19,140 MoBa subjects was also similar to the prevalence in our smaller sample (22% and 31%, respectively). The percentage of women that stopped smoking during pregnancy was 16% among the 19,140 MoBa subjects and 15% in our sample. Among mothers

who reported that they stopped smoking during pregnancy, the median gestational week of quitting was 5.0 [interquartile range (IQR) 4-9] in both the 19,140 MoBa subjects and our sample.

Cotinine measurements were available only at approximately gestational week 18, so we could not objectively confirm reports on the 30-week questionnaire of quitting later in pregnancy. Among the 136 women smoking at gestational week 18 based on cotinine, only 8 (6%) reported that they quit smoking later in pregnancy; excluding these 8 women yielded similar results.

This study cannot address mechanisms for the apparent need for sustained exposure (defined in this study as

through at least 18 weeks gestation) *in utero* to identify methylation differences at birth. However, one can speculate that methylation differences reflect a cellular response to exposure that enables fetal cells, such as hematopoietic progenitors, to be more resistant to toxins in tobacco smoke. If this occurs, these cells may have a selective advantage for survival in the presence of exposure to Ah receptor ligands such as polycyclic aromatic hydrocarbons. However, when exposure is not persistent, proliferation of progenitor cells without these epigenetic changes might be favored.

If mothers who were able to quit smoking early in pregnancy smoke less than mothers who continued to smoke, it is possible that this lower amount of smoking accounts for why we see no significant effect of quitting early in pregnancy. However, adjustment for the number of cigarettes smoked per day reported by the mother early in pregnancy did not materially alter the associations. The 136 women who smoked through at least gestational week 18 were disproportionately light smokers (median cigarettes per day = 4; IQR = 2–10) and thus we had little power to observe a dose response. However, we observed a statistically significant association between cigarettes smoked per day and methylation at 5 of the 8 *GFI1* CpGs and 2 of the 4 *AHRR* CpGs, adjusting for maternal age, parity, and education and applying Bonferroni correction ($P < 0.0019$).

Our previous publication included the Newborn Epigenetics Study as a replication dataset to confirm our findings (3). We were not able to use those data for replication of the current analyses because information on the grandmother's smoking during pregnancy with the mother, the mother's past smoking, mother cotinine data, and the father's past smoking was not available.

Subsequent to our original article, a wide range of preprocessing methods for the Illumina 450K data, including background correction, normalization, and transformation, have been published. In addition to the BMIQ performed for this analysis, we evaluated various normalization methods in this dataset and observed little impact on the results (31).

Environmental stimuli, including smoking, can affect the cell type composition of blood. In our previous report, we had evaluated possible confounding of our findings by smoking-related shifts in cell type by evaluating methylation at the top CpGs in two major cell pools—mononuclear cells (mostly lymphocytes) and granulocytes (mostly polymorphonuclear leukocytes; ref. 3). The methylation differences by smoking were in all cases larger than methylation differences between these two major pools, suggesting that the results were not due to confounding by smoking-related shifts in differential cell counts. Subsequent to submission of our publication (3), a method for statistical adjustment for cell type differences was published (25) and implementing this did not alter our results for the 26 CpGs. Of note, Shenker and colleagues confirmed findings in blood for *AHRR* by showing differential methylation at *AHRR* by smoking in the lung. Shenker

and colleagues also reported consistent methylation levels for *AHRR* cg05775921 across 6 different cell type pools (9). These various lines of evidence make it unlikely that our previously published findings reflect smoking-related cell type differences.

There is little information about the functional impact of the methylation differences that we observed, many of which have been replicated in studies of adult smokers (7–10). However, for one of the top hits in our analysis and several studies of adult smoking, *AHRR* cg05575921, Zeilinger and colleagues found methylation-specific protein-binding patterns (10). In addition, Shenker and colleagues (9) observed decreases in *AHRR* expression related to *AHRR* methylation.

We use the term epigenetic inheritance to indicate epigenetic modifications in the germline that escape meiotic resetting during gametogenesis (32), where meiotic resetting is the DNA methylation erasure that takes place just after fertilization (epigenetic reprogramming; ref. 33). Other than imprinting, the data on epigenetic inheritance in humans are sparse. The epigenetic inheritance effects that we evaluated here might be more precisely determined by comparing DNA methylation status in relation to smoking across the three generations—grandparents, parents, and the study child. Unfortunately, we do not have these extensive data. However, even with such direct cross-generation data, interpretation could be challenging because of known methylation changes with age (34) as well as distinguishing between the effects of personal smoking in adults and their *in utero* exposure. We do believe that our data in newborns at different levels of exposure bring relevant evidence to bear on this difficult issue. Our findings do not support the inheritance of these epigenetic marks across generations or from embryonic to somatic cells.

Our results may have relevance to the design of epidemiologic studies of maternal smoking and childhood leukemia or other cancers. Recent studies of childhood leukemia have not identified associations between maternal smoking during pregnancy and childhood leukemia (35–38). However, maternal smoking was classified in these studies based on any smoking during pregnancy. In our study, more than half of the women who reported smoking in the early part of pregnancy had quit by week 18, confirmed by cotinine measurements. Although we do not know that the methylation changes that we observed predispose to the development of childhood leukemia, our results indicate that a potential biomarker of *in utero* exposure to maternal smoking in pregnancy is only seen with sustained exposure (through at least 18 weeks gestation). We note that several of the genes differentially methylated in relation to smoking in our data are plausibly involved in normal or disordered hematopoiesis, including *RUNX1* (39), *MYO1G* (40), *GFI1* (41), and *AHRR* (42). The fact that methylation changes that we observed require sustained exposure suggests that epidemiologic studies of leukemia or other childhood cancers or health outcomes may miss true associations if data on maternal

smoking are limited to ever smoking during the pregnancy.

Disclosure of Potential Conflicts of Interest

Ø. Midttun is a board member of Bevital A/S, Norway, which promotes research into vitamin B12 deficiency. No potential conflicts of interest were disclosed by the other authors.

Authors' Contributions

Conception and design: B.R. Joubert, S.E. Håberg, W. Nystad, S.J. London
Development of methodology: B.R. Joubert, Ø. Midttun, P.M. Ueland, S.D. Peddada, S.J. London

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): B.R. Joubert, S.E. Håberg, R.M. Nilsen, S.E. Vollset, Ø. Midttun, W. Nystad

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): B.R. Joubert, S.E. Håberg, D.A. Bell, R.M. Nilsen, S.E. Vollset, M.C. Wu, W. Nystad, S.D. Peddada, S.J. London
Writing, review, and/or revision of the manuscript: B.R. Joubert, S.E. Håberg, D.A. Bell, R.M. Nilsen, S.E. Vollset, Ø. Midttun, P.M. Ueland, W. Nystad, S.J. London

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): B.R. Joubert, S.E. Håberg, W. Nystad, S.J. London

Study supervision: B.R. Joubert, W. Nystad, S.J. London

Acknowledgments

The authors thank the families in Norway for taking part in this ongoing cohort study; Rolf Erik Kolstad, Arild Sunde, and Kari Harbak from the MoBa Biobank, and Huiling Li, Gary Pittman, Laura Wharey, and Kevin Gerrish from the NIEHS for expert technical assistance. They also thank Shuangshuang Dai of SRA International Inc., Jianping Jin and David Shore of Westat, and Huiling Li of the NIEHS for computing support.

Grant Support

B.R. Joubert, S.J. London, D.A. Bell, and S.D. Peddada are supported by the Intramural Research Program of the NIH, NIEHS (Z01-ES-49019 and Z01-ES-046008). S.E. Håberg and W. Nystad are supported by the Norwegian Ministry of Health and the Ministry of Education and Research, NIH/NIEHS (contract no. NO-ES-75558), NIH/NINDS (grant no.1 UO1 NS 047537-01), and the Norwegian Research Council/FUGE (grant no. 151918/S10). R.M. Nilsen is supported by the Faculty of Medicine and Dentistry, University of Bergen, Norway.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received December 3, 2013; revised March 8, 2014; accepted March 26, 2014; published OnlineFirst April 16, 2014.

References

- Office of the Surgeon General. The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General. Rockville, MD: U.S. Dept. of Health and Human Services, Public Health Service, Office of the Surgeon General; 2006.
- Breton CV, Byun HM, Wenten M, Pan F, Yang A, Gilliland FD. Prenatal tobacco smoke exposure affects global and gene-specific DNA methylation. *Am J Respir Crit Care Med* 2009;180:462–7.
- Joubert BR, Haberg SE, Nilsen RM, Wang X, Vollset SE, Murphy SK, et al. 450K epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy. *Environ Health Perspect* 2012;120:1425–31.
- Suter M, Ma J, Harris AS, Patterson L, Brown KA, Shope C, et al. Maternal tobacco use modestly alters correlated epigenome-wide placental DNA methylation and gene expression. *Epigenetics* 2011;6:1284–94.
- Lee KW, Pausova Z. Cigarette smoking and DNA methylation. *Front Genet* 2013;4:132.
- Flom JD, Ferris JS, Liao Y, Tehranifar P, Richards CB, Cho YH, et al. Prenatal smoke exposure and genomic DNA methylation in a multi-ethnic birth cohort. *Cancer Epidemiol Biomarkers Prev* 2011;20:2518–23.
- Monick MM, Beach SR, Plume J, Sears R, Gerrard M, Brody GH, et al. Coordinated changes in AHRR methylation in lymphoblasts and pulmonary macrophages from smokers. *Am J Med Genet B Neuropsychiatr Genet* 2012;159B:141–51.
- Philibert RA, Beach SR, Brody GH. Demethylation of the aryl hydrocarbon receptor repressor as a biomarker for nascent smokers. *Epigenetics* 2012;7:1331–8.
- Shenker NS, Polidoro S, van Veldhoven K, Sacerdote C, Ricceri F, Birrell MA, et al. Epigenome-wide association study in the European Prospective Investigation into Cancer and Nutrition (EPIC-Turin) identifies novel genetic loci associated with smoking. *Hum Mol Genet* 2013;22:843–51.
- Zeilinger S, Kuhnel B, Klopp N, Baurecht H, Kleinschmidt A, Gieger C, et al. Tobacco smoking leads to extensive genome-wide changes in DNA methylation. *PLoS ONE* 2013;8:e63812.
- Miller LL, Henderson J, Northstone K, Pembrey M, Golding J. Do grandmaternal smoking patterns influence the aetiology of childhood asthma? *Chest* 2013 Oct 24. [Epub ahead of print].
- Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K, Sjöström M, et al. Sex-specific, male-line transgenerational responses in humans. *Eur J Hum Genet* 2006;14:159–66.
- Grossniklaus U, Kelly B, Ferguson-Smith AC, Pembrey M, Lindquist S. Transgenerational epigenetic inheritance: how important is it? *Nat Rev Genet* 2013;14:228–35.
- Daxinger L, Whitelaw E. Understanding transgenerational epigenetic inheritance via the gametes in mammals. *Nat Rev Genet* 2012;13:153–62.
- Magnus P, Irgens LM, Haug K, Nystad W, Skjaerven R, Stoltenberg C. Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa). *Int J Epidemiol* 2006;35:1146–50.
- Ronningen KS, Paltiel L, Meltzer HM, Nordhagen R, Lie KK, Hovengen R, et al. The biobank of the Norwegian Mother and Child Cohort Study: a resource for the next 100 years. *Eur J Epidemiol* 2006;21:619–25.
- Håberg SE, London SJ, Nafstad P, Nilsen RM, Ueland PM, Vollset SE, et al. Maternal folate levels in pregnancy and asthma in children at age 3 years. *J Allergy Clin Immunol* 2011;127:262–4, 4 e1.
- Bibikova M, Barnes B, Tsan C, Ho V, Klotzle B, Le JM, et al. High density DNA methylation array with single CpG site resolution. *Genomics* 2011;98:288–95.
- Sandoval J, Heyn HA, Moran S, Serra-Musach J, Pujana MA, Bibikova M, et al. Validation of a DNA methylation microarray for 450,000 CpG sites in the human genome. *Epigenetics* 2011;6:692–702.
- Teschendorff AE, Marabita F, Lechner M, Bartlett T, Tegner J, Gomez-Cabrero D, et al. A beta-mixture quantile normalization method for correcting probe design bias in Illumina Infinium 450 k DNA methylation data. *Bioinformatics* 2013;29:189–96.
- Midttun O, Hustad S, Ueland PM. Quantitative profiling of biomarkers related to B-vitamin status, tryptophan metabolism and inflammation in human plasma by liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom* 2009;23:1371–9.
- Shaw GM, Carmichael SL, Vollset SE, Yang W, Finnell RH, Blom H, et al. Mid-pregnancy cotinine and risks of orofacial clefts and neural tube defects. *J Pediatr* 2009;154:17–9.
- Cupul-Uicab LA, Baird DD, Skjaerven R, Saha-Chaudhuri P, Haug K, Longnecker MP. In utero exposure to maternal smoking and women's risk of fetal loss in the Norwegian Mother and Child Cohort (MoBa). *Hum Reprod* 2011;26:458–65.
- Fox J, Weisberg S. *An R companion to applied regression*. 2nd ed. Thousand Oaks, CA: SAGE Publications; 2011.
- Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, et al. DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC Bioinformatics* 2012;13:86.

26. Reinius LE, Acevedo N, Joerink M, Pershagen G, Dahlen SE, Greco D, et al. Differential DNA methylation in purified human blood cells: implications for cell lineage and studies on disease susceptibility. *PLoS ONE* 2012;7:e41361.
27. R Development Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2010.
28. Kvalvik LG, Nilsen RM, Skjaerven R, Vollset SE, Midttun O, Ueland PM, et al. Self-reported smoking status and plasma cotinine concentrations among pregnant women in the Norwegian Mother and Child Cohort Study. *Pediatr Res* 2012;72:101–7.
29. Braverman MT, Aaro LE, Hetland J. Changes in smoking among restaurant and bar employees following Norway's comprehensive smoking ban. *Health Promot Int* 2008;23:5–15.
30. Kvalvik LG, Nilsen RM, Skjaerven R, Vollset SE, Midttun O, Ueland PM, et al. Self-reported smoking status and plasma cotinine concentrations among pregnant women in the Norwegian Mother and Child Cohort Study. *Pediatr Res* 2012;72:101–7.
31. Wu MC, Joubert BR, Kuan PF, Haberg SE, Nystad W, Peddada SD, et al. A systematic assessment of normalization approaches for the Infinium 450k methylation platform. *Epigenetics* 2013;9:318–29.
32. Gomez-Diaz E, Jorda M, Peinado MA, Rivero A. Epigenetics of host-pathogen interactions: the road ahead and the road behind. *PLoS Pathog* 2012;8:e1003007.
33. Monk M, Boubelik M, Lehnert S. Temporal and regional changes in DNA methylation in the embryonic, extraembryonic and germ cell lineages during mouse embryo development. *Development* 1987;99:371–82.
34. Florath I, Butterbach K, Muller H, Bewerunge-Hudler M, Brenner H. Cross-sectional and longitudinal changes in DNA methylation with age: an epigenome-wide analysis revealing over 60 novel age-associated CpG sites. *Hum Mol Genet* 2014;23:1186–201.
35. Klimentopoulou A, Antonopoulos CN, Papadopoulou C, Kanavidis P, Tourvas AD, Polychronopoulou S, et al. Maternal smoking during pregnancy and risk for childhood leukemia: a nationwide case-control study in Greece and meta-analysis. *Pediatr Blood Cancer* 2012;58:344–51.
36. Metayer C, Zhang L, Wiemels JL, Bartley K, Schiffman J, Ma X, et al. Tobacco smoke exposure and the risk of childhood acute lymphoblastic and myeloid leukemias by cytogenetic subtype. *Cancer Epidemiol Biomarkers Prev* 2013;22:1600–11.
37. Milne E, Greenop KR, Scott RJ, Bailey HD, Attia J, Dalla-Pozza L, et al. Parental prenatal smoking and risk of childhood acute lymphoblastic leukemia. *Am J Epidemiol* 2012;175:43–53.
38. Slater ME, Linabery AM, Blair CK, Spector LG, Heerema NA, Robison LL, et al. Maternal prenatal cigarette, alcohol and illicit drug use and risk of infant leukaemia: a report from the Children's Oncology Group. *Paediatr Perinat Epidemiol* 2011;25:559–65.
39. Tsuzuki S, Seto M. Expansion of functionally defined mouse hematopoietic stem and progenitor cells by a short isoform of RUNX1/AML1. *Blood* 2012;119:727–35.
40. Pierce RA, Field ED, Mutis T, Golovina TN, Von Kap-Herr C, Wilke M, et al. The HA-2 minor histocompatibility antigen is derived from a diallelic gene encoding a novel human class I myosin protein. *J Immunol* 2001;167:3223–30.
41. van der Meer LT, Jansen JH, van der Reijden BA. Gfi1 and Gfi1b: key regulators of hematopoiesis. *Leukemia* 2010;24:1834–43.
42. Zudaire E, Cuesta N, Murty V, Woodson K, Adams L, Gonzalez N, et al. The aryl hydrocarbon receptor repressor is a putative tumor suppressor gene in multiple human cancers. *J Clin Invest* 2008;118:640–50.

Cancer Epidemiology, Biomarkers & Prevention

Maternal Smoking and DNA Methylation in Newborns: *In Utero* Effect or Epigenetic Inheritance?

Bonnie R. Joubert, Siri E. Håberg, Douglas A. Bell, et al.

Cancer Epidemiol Biomarkers Prev 2014;23:1007-1017. Published OnlineFirst April 16, 2014.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-13-1256](https://doi.org/10.1158/1055-9965.EPI-13-1256)

Supplementary Material Access the most recent supplemental material at:
<http://cebp.aacrjournals.org/content/suppl/2014/04/17/1055-9965.EPI-13-1256.DC1>

Cited articles This article cites 38 articles, 5 of which you can access for free at:
<http://cebp.aacrjournals.org/content/23/6/1007.full#ref-list-1>

Citing articles This article has been cited by 7 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/23/6/1007.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/23/6/1007>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.