

Long-term Particulate Matter Exposures during Adulthood and Risk of Breast Cancer Incidence in the Nurses' Health Study II Prospective Cohort

Jaime E. Hart^{1,2}, Kimberly A. Bertrand³, Natalie DuPre⁴, Peter James^{2,4}, Verónica M. Vieira⁵, Rulla M. Tamimi^{1,4}, and Francine Laden^{1,2,4}

Abstract

Background: There is increasing concern that environmental exposures, such as air pollution, may be related to increasing rates of breast cancer; however, results from cohort studies have been mixed. We examined the association between particulate matter (PM) and measures of distance to roadway with the risk of incident breast cancer in the prospective nationwide Nurses' Health Study II (NHSII) cohort.

Methods: Incident invasive breast cancer from 1993 to 2011 ($N = 3,416$) was assessed among 115,921 women in the NHSII cohort. Time-varying Cox proportional hazards models were used to calculate HRs and 95% confidence intervals (95% CI) for increases in ambient exposures to PM_{10} , $PM_{2.5-10}$, and $PM_{2.5}$ and residential roadway proximity categories.

Results: In multivariable adjusted models, there was little evidence of an increased risk of breast cancer (or any of the

receptor-specific subtypes) overall or by menopausal status with PM exposure. There was, however, a suggestion of increased risks among women living <50 m of the largest road type (HR = 1.60; 95% CI, 0.80–3.21) or within <50 m of the two largest road types (1.14; 95% CI, 0.84–1.54) compared with women living farther (≥ 200 m) away.

Conclusions: Among women in the NHSII, we found no statistically significant associations between particulate matter exposures and incidence of breast cancer overall, by menopausal status, or by hormone receptor subtype. There was, however, a suggestion that residential proximity to major roadways may be associated with increased risk.

Impact: These results suggest no elevated breast cancer risk with increasing exposures to particulate matter air pollution, but that other traffic-related exposures may be important. *Cancer Epidemiol Biomarkers Prev*; 25(8); 1274–6. ©2016 AACR.

Introduction

The International Agency for Research on Cancer monograph on the carcinogenicity of ambient air pollution presented mixed evidence for air pollution and breast cancer risk (1). However, some recent studies have suggested associations (2–7). Studies with large numbers of breast cancer cases, information on specific breast cancer subtypes, and the ability to control for other breast cancer risk factors are needed. Therefore, we examined the association of particulate matter (PM) exposures and roadway proximity on risk of overall,

hormone receptor-specific, and menopausal status-specific breast cancer risk among women in the nationwide, prospective Nurses' Health Study II (NHSII) cohort.

Materials and Methods

The NHSII is a cohort of 116,430 female nurses with no previous history of cancer (except nonmelanoma skin cancer) enrolled in 1989 when they were aged 25 to 42 years. Mailed biennial follow-up questionnaires update information on risk factors and medical history. Self-reports of incident cancers are confirmed by medical record review.

Biennial residential addresses were geocoded, and roadway proximity was assigned as a proxy for traffic-related exposures. Specifically, we calculated the distance from each address to the nearest street segments in the ESRI StreetPro 2007 data layer listed as U.S. Census Feature class codes A1, A2, or A3. On the basis of case distributions and exposure studies, we categorized distance to road as 0 to 49 m, 50 to 199 m, and ≥ 200 m, for all three road types together, for the two largest road types (A1, A2) and for the largest road type (A1). Forty-eight-month moving average and cumulative average exposures to three size fractions of particulate matter, PM_{10} , $PM_{2.5-10}$, and $PM_{2.5}$, were assigned to each participant based on monthly spatiotemporal prediction models applied to the geocoded address history (8). For participants who moved, we assumed they changed addresses at the beginning of the biennial cycle.

¹Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts. ²Exposure, Epidemiology, and Risk Program, Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, Massachusetts. ³Slone Epidemiology Center at Boston University, Boston, Massachusetts. ⁴Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts. ⁵Program in Public Health, University of California, Irvine, California.

R.M. Tamimi and F. Laden share senior authorship of this article.

Corresponding Author: Jaime E. Hart, Brigham and Women's Hospital and Harvard Medical School, 401 Park Drive, Landmark Center, 3rd Floor West, Boston, MA 02215. Phone: 617-525-2289; Fax: 617-525-2578; E-mail: jaime.hart@channing.harvard.edu

doi: 10.1158/1055-9965.EPI-16-0246

©2016 American Association for Cancer Research.

Time-varying Cox proportional hazards models on a biennial time scale were used to calculate HRs and 95% confidence intervals (95% CI) for each roadway proximity category compared with the farthest, and for each 10 $\mu\text{g}/\text{m}^3$ increase in each of the PM size fractions, after assessing linearity using splines. All models were adjusted for age, race, and calendar year; multivariable models were additionally adjusted for established breast cancer risk factors. Separate models assessed risk of overall breast cancer, and hormone receptor-specific subtypes (ER^+/PR^+ and ER^-/PR^-). We also stratified models by menopausal status.

Results

Eligible participants ($N = 115,921$) were 47 years old on average during follow-up, with an average body mass index (BMI) of 26 kg/m^2 . Sixteen percent were nulliparous, 10% were current users of postmenopausal hormone therapy, and 64% were never smokers. There was no association between exposures to 48-month PM with breast cancer overall or for specific hormone receptor subtypes, or by menopausal status (Tables 1 and 2); estimates were similar for cumulative average exposures (not shown). Although case numbers were small, there was a suggestion of increasing risk with residential proximity to the two largest road types, with a multivariable adjusted HR of 1.60 (95% CI, 0.80–3.21) for women living <50 m of the largest road type and 1.14 (95% CI, 0.84–1.54) for women closest to the two largest road types compared with those living ≥ 200 m away. These risks were consistent across the various breast cancer subtypes

Discussion

We did not observe an association between PM exposures and breast cancer incidence in this prospective cohort study. This is in contrast to an earlier case-control study in Western New York, where total suspended particulate exposures were associated with increased odds of postmenopausal breast cancer, but not premenopausal breast cancer (detailed in ref. 1). Our results are similar to a recent analysis in the Sister Study, where no elevations in overall or hormone receptor-specific breast cancer risk were observed with increasing PM_{10} or $\text{PM}_{2.5}$ exposures (2).

Our suggestive findings of an elevated risk with roadway proximity, although not statistically significant and based on small numbers of exposed cases, are in contrast to several studies that have found no association (1, 3) but do agree with some studies using other measures of traffic exposure (traffic density, motor vehicle density; refs. 1, 7). Our findings also agree with numerous studies that have observed positive associations between exposures to nitrogen oxides or other traffic-related pollutants and breast cancer (1–3, 5–7).

Our analyses in a large, nation-wide, prospective cohort has many strengths, including fine control for potential confounders, high spatial and temporal resolution estimates of PM exposure, and large numbers of cases in many subcategories of interest (e.g., premenopausal, ER^-/PR^-). However, we only had information on adult exposures, which may not be an important etiologic period. The findings in this cohort may not be generalizable to populations with more racial/ethnic diversity or a broader range of socioeconomic status. Future studies

Table 1. Risk of overall and hormone receptor subtype-invasive breast cancer with exposures to particulate matter and roadway proximity among 115,921 women in the NHSII followed 1993–2011

Exposure	Invasive breast cancer			ER^+/PR^+ Invasive breast cancer			ER^-/PR^- Invasive breast cancer			Multivariable ^b HR (95% CI)		
	Cases	Basic ^a HR (95% CI)	Multivariable ^b HR (95% CI)	Cases	Basic ^a HR (95% CI)	Multivariable ^b HR (95% CI)	Cases	Basic ^a HR (95% CI)	Multivariable ^b HR (95% CI)	Cases	Basic ^a HR (95% CI)	Multivariable ^b HR (95% CI)
48-month average												
PM_{10} (per 10 $\mu\text{g}/\text{m}^3$)	3,416	0.98 (0.92–1.04)	1.00 (0.93–1.07)	1,853	1.02 (0.94–1.11)	1.05 (0.95–1.15)	492	0.94 (0.79–1.11)	0.97 (0.80–1.18)			
$\text{PM}_{2.5-10}$ (per 10 $\mu\text{g}/\text{m}^3$)	3,416	1.01 (0.93–1.09)	1.06 (0.96–1.17)	1,853	1.07 (0.96–1.19)	1.13 (0.99–1.29)	492	0.91 (0.73–1.13)	0.96 (0.73–1.26)			
$\text{PM}_{2.5}$ (per 10 $\mu\text{g}/\text{m}^3$)	3,416	0.89 (0.78–1.01)	0.90 (0.79–1.03)	1,853	0.92 (0.77–1.10)	0.95 (0.79–1.14)	492	0.99 (0.70–1.39)	0.97 (0.68–1.40)			
Proximity to A1–A3 roads (m)												
0–49	341	0.92 (0.82–1.03)	0.95 (0.85–1.07)	194	0.98 (0.84–1.14)	1.01 (0.86–1.18)	52	0.95 (0.71–1.28)	1.02 (0.75–1.37)			
50–199	777	1.00 (0.92–1.09)	1.00 (0.92–1.09)	431	1.05 (0.93–1.17)	1.04 (0.93–1.16)	100	0.87 (0.69–1.09)	0.91 (0.72–1.14)			
≥ 200	1,954	Reference	Reference	1,050	Reference	Reference	289	Reference	Reference			
Proximity to A1–A2 roads (m)												
0–49	43	1.06 (0.78–1.43)	1.14 (0.84–1.54)	19	0.87 (0.55–1.36)	0.93 (0.59–1.47)	8	1.36 (0.67–2.73)	1.44 (0.71–2.92)			
50–199	177	1.14 (0.97–1.32)	1.16 (1.00–1.35)	102	1.21 (0.99–1.48)	1.23 (1.00–1.50)	25	1.12 (0.75–1.69)	1.18 (0.79–1.78)			
≥ 200	2,852	Reference	Reference	1,554	Reference	Reference	408	Reference	Reference			
Proximity to A1 roads (m)												
0–49	8	1.52 (0.76–3.05)	1.60 (0.80–3.21)	4	1.40 (0.52–3.75)	1.48 (0.55–3.97)	14	1.41 (0.83–2.41) ^c	1.52 (0.89–2.60) ^c			
50–199	76	1.10 (0.87–1.38)	1.11 (0.89–1.40)	40	1.07 (0.78–1.47)	1.08 (0.79–1.48)		Reference	Reference			
≥ 200	2,988	Reference	Reference	1,631	Reference	Reference	427	Reference	Reference			

NOTE: Distance to road measures were only calculated for women with a street level geocode or better ($N = 111,545$).

Abbreviations: ER, estrogen receptor; PR, progesterone receptor.

^aAdjusted for current age, race, and calendar period.

^bAdditionally adjusted for history of benign breast disease, family history, age at menarche, parity, age at first birth, height, BMI at age 18, current BMI, alcohol consumption at ages 15 to 17 and 18 to 22, overall diet quality (AHEI-2010), oral contraceptive use, menopausal status and hormone use, smoking status, physical activity, individual-level socioeconomic status (marital status, living arrangements, household income) and area-level socioeconomic status (census tract level median income and median home value), and region of residence.

^cThe two closest proximity categories were consolidated due to small case numbers.

Hart et al.

Table 2. Risk of invasive breast cancer by menopausal status with exposures to particulate matter and roadway proximity among women in the NHSII

Exposure	Premenopausal			Postmenopausal		
	Cases	Basic ^a HR (95% CI)	Multivariable ^b HR (95% CI)	Cases	Basic ^a HR (95% CI)	Multivariable ^b HR (95% CI)
48-month average						
PM ₁₀ (per 10 µg/m ³)	1,966	0.98 (0.90–1.07)	1.03 (0.93–1.13)	1,296	0.99 (0.89–1.10)	0.97 (0.86–1.09)
PM _{2.5–10} (per 10 µg/m ³)	1,966	0.98 (0.87–1.09)	1.07 (0.93–1.22)	1,296	1.08 (0.96–1.22)	1.07 (0.92–1.25)
PM _{2.5} (per 10 µg/m ³)	1,966	0.98 (0.83–1.16)	0.99 (0.83–1.18)	1,296	0.75 (0.60–0.93)	0.76 (0.61–0.95)
Proximity to A1–A3 roads (m)						
0–49	194	0.98 (0.84–1.14)	1.01 (0.86–1.17)	130	0.83 (0.69–1.00)	0.87 (0.72–1.05)
50–199	471	1.09 (0.98–1.22)	1.08 (0.97–1.21)	273	0.88 (0.76–1.01)	0.88 (0.77–1.01)
≥200	1,082	Reference	Reference	782	Reference	Reference
Proximity to A1–A2 roads (m)						
0–49	25	1.18 (0.79–1.75)	1.23 (0.83–1.83)	17	1.03 (0.64–1.67)	1.14 (0.70–1.84)
50–199	106	1.20 (0.99–1.47)	1.20 (0.98–1.46)	62	1.01 (0.78–1.30)	1.06 (0.82–1.38)
≥200	1,616	Reference	Reference	1,106	Reference	Reference
Proximity to A1 roads (m)						
0–49	5	1.80 (0.74–4.35)	1.74 (0.72–4.21)	3	1.34 (0.43–4.17)	1.48 (0.47–4.62)
50–199	49	1.27 (0.95–1.69)	1.26 (0.94–1.67)	25	0.93 (0.62–1.38)	0.97 (0.65–1.45)
≥200	1,693	Reference	Reference	1,157	Reference	Reference

NOTE: Distance to road measures were only calculated for the 115,921 women with a street level geocode or better ($N = 111,545$), and women can contribute person-time to both pre- and postmenopausal models.

^aAdjusted for current age, race, and calendar period.

^bAdditionally adjusted for history of benign breast disease, family history, age at menarche, parity, age at first birth, height, BMI at age 18, current BMI, alcohol consumption at ages 15 to 17 and 18 to 22, overall diet quality (AHEI-2010), oral contraceptive use, menopausal hormone use, smoking status, physical activity, individual-level socioeconomic status (marital status, living arrangements, household income) and area-level socioeconomic status (census tract level median income and median home value), and region of residence.

should examine the association of traffic-related exposures in these groups.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: J.E. Hart, K.A. Bertrand, V.M. Vieira, R.M. Tamimi, F. Laden

Development of methodology: J.E. Hart, R.M. Tamimi, F. Laden

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): J.E. Hart, F. Laden

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J.E. Hart, K.A. Bertrand, R.M. Tamimi, F. Laden

Writing, review, and/or revision of the manuscript: J.E. Hart, K.A. Bertrand, N. DuPre, P. James, V.M. Vieira, R.M. Tamimi, F. Laden

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): R.M. Tamimi

Study supervision: R.M. Tamimi, F. Laden

Acknowledgments

The authors thank the participants and staff of the Nurses' Health Study II and following state cancer registries: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, WY.

Grant Support

All authors were funded by Susan G. Komen for the Cure grant IIR13264020. The NHSII cohort is funded by NIH grant UM1 CA176726. J.E. Hart and F. Laden received R01 ES017017, J.E. Hart received P30 ES000002, and P. James and N. DuPre were supported by T32 CA090001.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received March 24, 2016; accepted May 10, 2016; published OnlineFirst June 2, 2016.

References

- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Outdoor air pollution. Lyon, France: International Agency for Research on Cancer; 2015.
- Reding KW, Young MT, Szpiro AA, Han CJ, DeRoo LA, Weinberg C, et al. Breast cancer risk in relation to ambient air pollution exposure at residences in the sister study cohort. *Cancer Epidemiol Biomarkers Prev* 2015;24:1907–9.
- Hystad P, Villeneuve PJ, Goldberg MS, Crouse DL, Johnson K, Canadian Cancer Registries Epidemiology Research Group. Exposure to traffic-related air pollution and the risk of developing breast cancer among women in eight Canadian provinces: a case-control study. *Environ Int* 2015;74:240–8.
- Shekarrizfard M, Valois MF, Goldberg MS, Crouse D, Ross N, Parent ME, et al. Investigating the role of transportation models in epidemiologic studies of traffic related air pollution and health effects. *Environ Res* 2015;140:282–91.
- Mordukhovich I, Beyea J, Herring AH, Hatch M, Stellman SD, Teitelbaum SL, et al. Vehicular traffic-related polycyclic aromatic hydrocarbon exposure and breast cancer incidence: The Long Island Breast Cancer Study Project (LIBCSP). *Environ Health Perspect* 2016;124:30–8.
- Al-Ahmadi K, Al-Zahrani A. NO(2) and cancer incidence in Saudi Arabia. *Int J Environ Res Public Health* 2013;10:5844–62.
- Chen F, Bina WF. Correlation of white female breast cancer incidence trends with nitrogen dioxide emission levels and motor vehicle density patterns. *Breast Cancer Res Treat* 2012;132:327–33.
- Yanosky JD, Paciorek CJ, Laden F, Hart JE, Puett RC, Liao D, et al. Spatio-temporal modeling of particulate air pollution in the conterminous United States using geographic and meteorological predictors. *Environ Health* 2014;13:63.

Cancer Epidemiology, Biomarkers & Prevention

Long-term Particulate Matter Exposures during Adulthood and Risk of Breast Cancer Incidence in the Nurses' Health Study II Prospective Cohort

Jaime E. Hart, Kimberly A. Bertrand, Natalie DuPre, et al.

Cancer Epidemiol Biomarkers Prev 2016;25:1274-1276. Published OnlineFirst June 2, 2016.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-16-0246](https://doi.org/10.1158/1055-9965.EPI-16-0246)

Cited articles This article cites 7 articles, 1 of which you can access for free at:
<http://cebp.aacrjournals.org/content/25/8/1274.full#ref-list-1>

Citing articles This article has been cited by 3 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/25/8/1274.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/25/8/1274>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.