

Genetic Variants in Immune-Related Pathways and Breast Cancer Risk in African American Women in the AMBER Consortium



Chi-Chen Hong¹, Lara E. Sucheston-Campbell², Song Liu³, Qiang Hu³, Song Yao¹, Kathryn L. Lunetta⁴, Stephen A. Haddad⁵, Edward A. Ruiz-Narváez⁶, Jeannette T. Bensen⁷, Ting-Yuan David Cheng⁸, Elisa V. Bandera⁹, Lynn A. Rosenberg⁵, Christopher A. Haiman¹⁰, Kelvin Lee¹¹, Sharon S. Evans¹¹, Scott I. Abrams¹¹, Elizabeth A. Repasky¹¹, Andrew F. Olshan⁷, Julie R. Palmer⁵, and Christine B. Ambrosone¹

Abstract

Background: Constitutional immunity shaped by exposure to endemic infectious diseases and parasitic worms in Sub-Saharan Africa may play a role in the etiology of breast cancer among African American (AA) women.

Methods: A total of 149,514 gene variants in 433 genes across 45 immune pathways were analyzed in the AMBER consortium among 3,663 breast cancer cases and 4,687 controls. Gene-based pathway analyses were conducted using the adaptive rank truncated product statistic for overall breast cancer risk, and risk by estrogen receptor (ER) status. Unconditional logistic regression analysis was used to estimate ORs and 95% confidence intervals (CIs) for single variants.

Results: The top pathways were Interleukin binding ($P = 0.01$), Biocarta TNFR2 ($P = 0.005$), and positive regulation of cytokine production ($P = 0.024$) for overall, ER⁺, and ER⁻ cancers, respectively. The most significant gene was *IL2RB*

($P = 0.001$) for overall cancer, with rs228952 being the top variant identified (OR = 0.85; 95% CI, 0.79–0.92). Only *BCL3* contained a significant variant for ER⁺ breast cancer. Variants in *IL2RB*, *TLR6*, *IL8*, *PRKDC*, and *MAP3K1* were associated with ER⁻ disease. The only genes showing heterogeneity between ER⁻ and ER⁺ cancers were *TRAF1*, *MAP3K1*, and *MAPK3* ($P \leq 0.02$). We also noted genes associated with autoimmune and atopic disorders.

Conclusions: Findings from this study suggest that genetic variants in immune pathways are relevant to breast cancer susceptibility among AA women, both for ER⁺ and ER⁻ breast cancers.

Impact: Results from this study extend our understanding of how inherited genetic variation in immune pathways is relevant to breast cancer susceptibility. *Cancer Epidemiol Biomarkers Prev*; 27(3): 321–30. ©2018 AACR.

¹Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, New York. ²College of Pharmacy, The Ohio State University, Columbus, Ohio. ³Department of Biostatistics and Bioinformatics, Roswell Park Cancer Institute, Buffalo, New York. ⁴Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts. ⁵Slope Epidemiology Center at Boston University, Boston, Massachusetts. ⁶University of Michigan School of Public Health, Ann Arbor, Michigan. ⁷Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina. ⁸Department of Epidemiology, College of Public Health and Health Professions & College of Medicine, University of Florida, Gainesville, Florida. ⁹Cancer Prevention and Control, Rutgers Cancer Institute of New Jersey, New Brunswick, New Jersey. ¹⁰Department of Preventive Medicine, Keck School of Medicine, University of Southern California/Norris Comprehensive Cancer Center, Los Angeles, California. ¹¹Department of Immunology, Roswell Park Cancer Institute, Buffalo, New York.

Note: Supplementary data for this article are available at *Cancer Epidemiology, Biomarkers & Prevention* Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Chi-Chen Hong, Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY 14263. Phone: 716-845-7785; Fax: 716-845-3200; E-mail: Chi-Chen.Hong@RoswellPark.org

doi: 10.1158/1055-9965.EPI-17-0434

©2018 American Association for Cancer Research.

Introduction

Endemic infectious diseases fundamentally shape the immune systems of populations at the genetic level, selecting for immunity that preserves well-being through the reproductive years (1). The immune system is organized into two central arms of defense: the nonspecific innate arm that provides protection against a wide variety of pathogens by activating the inflammatory response, and the more specific adaptive arm that targets specific antigens of pathogens (2). Individuals of African ancestry (AA) show inherited differences in constitutional immune function from European Americans (EA), partly due to evolutionary responses to infection and endemic helminth exposure in Sub-Saharan Africa, causing both a stronger inflammatory immune response as well as a type II immunosuppressive phenotype (1, 3).

As reviewed by Palmer and colleagues (4), AA women are more likely than EAs to be diagnosed with breast cancer before age 50, and to have more aggressive disease of higher grade, lacking receptors for estrogen (ER), progesterone (PR), and HER2, which precludes use of targeted therapies, often resulting in poorer survival. Although it is becoming clearer that some reproductive and hormonal factors may, in part, account for some of the disparities in breast cancer aggressiveness (5, 6), it is possible that constitutional immune phenotype could play a role in breast

Hong et al.

cancer etiology, particularly for ER⁻ disease (7). In the AMBER Consortium, we hypothesized that the higher prevalence of ER⁻ tumors in AAs could arise, in part, from differences in immunologic profiles such as those that typically define inflammation and/or suppressive phenotypes. Thus, we examined genetic variation in immune response pathways involved in host defense and/or self-tolerance in relation to overall breast cancer risk and risk of ER⁻ compared with ER⁺ cancers. Because ER⁻ breast cancers are most prevalent among premenopausal AAs, and immune biomarkers were previously observed to be more strongly associated with being diagnosed with ER⁻ compared with ER⁺ breast cancer among premenopausal women (8), we also examined the role of genetic variation in immune pathways and breast cancer risk by menopausal status.

Materials and Methods

Study population

These analyses were conducted in the context of the African American Breast Cancer Epidemiology and Risk (AMBER) Consortium designed to investigate the etiology of breast cancer subtypes among AA women (4). Studies forming the basis for AMBER were the Women's Circle of Health Study (WCHS), the Carolina Breast Cancer Study (CBCS), the Black Women's Health Study (BWHS), and the Multi-Ethnic Cohort Study (MEC; 9–13), which are described in Supplementary Methods S1. Research protocols were conducted in accordance with the ethical standards of the Declaration of Helsinki, and were approved by the Institutional Review Board at each participating institution. Informed consent was provided by all study participants. Table 1 shows the total number of cases and controls included in the analysis by age, menopausal status, and ER status.

Gene and SNP selection

A total of 433 genes in 45 curated pathways involved in immune response were selected from the Molecular Signatures Database (MSigDB) and included BIOCARTA, KEGG, REACTOME, and Gene Ontology (GO) pathways (14). Pathways considered for inclusion were those related to cytokines, Th1- and Th2-related immunity, inflammation, T-cell activation, and immune response regulation. Priority was given to those containing the highest proportion of genes identified in GWAS (full

catalog downloaded from the National Genome Research Institute on March 8, 2012) found to be associated with breast cancer, or disease conditions and/or phenotypes associated with dysregulated immune function, hormonal or reproductive factors, obesity-related traits, or circulating vitamin D levels as risk factors for breast cancer. A full list of traits used to generate the list of GWAS genes used to help prioritize immune pathways is provided in Supplementary Table S1. The list of GWAS genes used for prioritization is provided in Supplementary Table S2, with $N = 74$ found within the 45 immune pathways prioritized. Tag single-nucleotide polymorphisms (SNP) were selected for the 433 unique genes ± 10 kb to capture SNPs with minor allele frequency $\geq 10\%$ (at $r^2 \geq 0.8$) based on the haplotype structure of the Yoruban population (YRI) in 1000 Genomes Phase I reference panel (<http://www.1000genomes.org>).

Genotyping and quality control protocol

Genotyping was performed by the Center for Inherited Disease (CIDR) using the Illumina Human Exome Beadchip v1.1 with additional custom content SNPs, as described previously (15). A total of 13,235 SNPs in the 433 immune-related genes examined passed quality control (QC). Imputation was performed using IMPUTE2 software (16) and the 1000 Genomes Phase I reference panel (May 21, 2011 1000G data, December 2013 haplotype release in IMPUTE2 site). Genetic data from 533 cases and 989 controls were available in the MEC, from a previous GWAS on the Illumina Human 1 M-Duo chip, with SNPs imputed to the same release of 1000 Genomes. Imputed genotypes were combined into a final dataset after omitting variants with mismatching alleles in AMBER and MEC, allele frequencies that were different by more than 0.15, and had imputation INFO scores < 0.5 in either study. A total of 149,514 SNPs were included in these analyses. Genotype principal components were computed using the smartpca program in the EIGENSOFT package to examine population structure.

Protein–protein interaction network construction

The Search Tool for the Retrieval of Interacting Genes (STRING, V.10) database was used to construct a protein–protein interaction network to identify potentially relevant immune pathways. The STRING database provides integrated knowledge on the known and predicted functional relationships between proteins

Table 1. Descriptive characteristics of participants in the AMBER consortium by study

	BWHS		CBCS		WCHS		MEC		AMBER	
	Ca	Co	Ca	Co	Ca	Co	Ca	Co	Ca	Co
Total, <i>N</i>	901	2,249	1,408	615	821	834	533	989	3,663	4,687
Age, years, <i>n</i> (%)										
<40	47 (5.2%)	217 (9.6%)	204 (14.5%)	87 (14.1%)	85 (10.4%)	116 (13.9%)	0 (0.0%)	0 (0.0%)	336 (9.2%)	420 (9.0%)
40–49	262 (29.1%)	652 (29.0%)	459 (32.6%)	211 (34.3%)	215 (26.2%)	228 (27.3%)	9 (1.7%)	16 (1.6%)	945 (25.8%)	1,107 (23.6%)
50–59	302 (33.5%)	770 (34.2%)	381 (27.1%)	150 (24.4%)	292 (35.6%)	319 (38.2%)	112 (21.0%)	222 (22.4%)	1,087 (29.7%)	1,461 (31.2%)
60–69	204 (22.6%)	442 (19.7%)	267 (19.0%)	114 (18.5%)	173 (21.1%)	142 (17.0%)	175 (32.8%)	288 (29.1%)	819 (22.4%)	986 (21.0%)
70+	86 (9.5%)	168 (7.5%)	97 (6.9%)	53 (8.6%)	56 (6.8%)	29 (3.5%)	237 (44.5%)	463 (46.8%)	476 (13.0%)	713 (15.2%)
Menopausal status, <i>n</i> (%)										
Premenopausal	310 (34.4%)	834 (37.1%)	526 (37.4%)	234 (38.0%)	363 (44.2%)	394 (47.2%)	0 (0.0%)	0 (0.0%)	1,199 (32.7%)	1,462 (31.2%)
Postmenopausal	482 (53.5%)	1,178 (52.4%)	745 (52.9%)	308 (50.1%)	361 (44.0%)	386 (46.3%)	533 (100.0%)	989 (100.0%)	2,121 (57.9%)	2,861 (61.0%)
Unknown	109 (12.1%)	237 (10.5%)	137 (9.7%)	73 (11.9%)	97 (11.8%)	54 (6.5%)	0 (0.0%)	0 (0.0%)	343 (9.4%)	364 (7.8%)
ER status, <i>n</i> (%)										
ER ⁺	498 (55.3%)		741 (52.6%)		435 (53.0%)		309 (58.0%)		1,983 (54.1%)	
ER ⁻	233 (25.9%)		565 (40.1%)		165 (20.1%)		135 (25.3%)		1,098 (30.0%)	
Unknown	170 (18.9%)		102 (7.2%)		221 (26.9%)		89 (16.7%)		582 (15.9%)	

Abbreviations: AMBER, African American Breast Cancer Epidemiology and Risk; BWHS, Black Women's Health Study; Ca, cases; CBCS, Carolina Breast Cancer Study; Co, controls; ER, estrogen receptor; MEC, Multi-Ethnic Cohort; WCHS, Women's Circle of Health Study.

(17). Only associations with high confidence (score ≥ 0.7) are presented. Networks were generated for significant genes identified for breast cancer overall, and by ER and menopausal status.

Statistical analysis

As described previously (15), gene-based pathway analyses were conducted using the adaptive rank truncated product (ARTP) statistic as implemented in the R package PIGE, which allowed us to optimize the number of SNP P values combined in each pathway and in each gene-level test. To avoid capturing highly correlated signals within a gene, a subset of "pruned-in" SNPs were selected to ensure that all SNP pairs had linkage disequilibrium (LD) $r^2 < 0.8$, using the R2 filter option in the R package AdaJoint. The ARTP gene-level tests combined the optimal number of the most significant SNP P values from among the top 10 pruned-in SNPs for each gene. Single variant tests were only conducted within genes reaching a nominal significance level of $P = 0.05$. Unconditional logistic regression analysis was performed using imputed dosage genotype data. A Bonferroni correction was applied for the number of pruned-in SNPs tested within each gene to identify SNPs with gene-wide significance. We performed case–case analyses to compare odds of being diagnosed with ER⁻ to ER⁺ disease to identify pathways, genes, and SNPs that may differentially impact ER status. Although none of the genotype principal components tested were strongly associated with risk, we included three principal components that were associated at $P < 0.10$ in all regression models to control for any potential confounding by population structure. Models were also adjusted for study, age, geographic location, and DNA source (saliva, blood, mouthwash). Lifestyle factors and comorbid conditions that affect immune function and breast cancer risk were not treated as potential confounders in our study because they could be on the causal pathway between genetic variation and breast cancer risk, and adjustments may inappropriately attenuate risk estimates. Functional follow-up was performed in RegulomeDB, PolyPhen-2, HaploReg v4.1, and GTEx databases (18–21).

Results

Pathways associated with breast cancer risk

Pathway analyses yielded nominally or borderline significant associations for overall breast cancer with cytokine-related pathways (IL receptor activity, $P = 0.05$; IL binding, $P = 0.01$; cytokine and chemokine mediated signaling, $P = 0.04$) as well as the immune effector process pathway ($P = 0.04$; Supplementary Table S3). When stratifying by ER status, only the Biocarta TNFR2 pathway achieved nominal significance for ER⁺ cancer ($P = 0.005$), and the positive regulation of cytokine production pathway was associated with ER⁻ disease ($P = 0.024$).

When considering menopausal status, pathways most significant for premenopausal breast cancer (Supplementary Table S4) were those associated with regulation of immune system processes (positive regulation of immune system process, $P = 0.005$; regulation of immune system process, $P = 0.01$; immune effector process, $P = 0.02$) and lymphocyte and T-cell activation (regulation of lymphocyte activation, $P = 0.03$; regulation of T-cell activation, $P = 0.06$). Premenopausal ER⁺ cancers were most strongly related to programmed death-1 (PD-1) signaling ($P = 0.04$), while ER⁻ cancers were associated with the adaptive immune response pathway ($P = 0.03$), and cytokine and inflammation-related pathways (Biocarta cytokine pathway, $P = 0.004$;

Biocarta inflammation pathway, $P = 0.03$). Immune activation pathways were also important among postmenopausal women (Supplementary Table S5; regulation of lymphocyte activation, $P = 0.06$; T-cell activation, $P = 0.06$), but cytokine-related pathways appeared to play a more prominent role (IL receptor activity, $P = 0.01$; IL binding, $P = 0.02$; cytokine and chemokine-mediated signaling pathway, $P = 0.03$), involved in both ER⁺ (cytokine binding, $P = 0.03$; Biocarta IL5 pathway, $P = 0.05$), and ER⁻ cancers (cytokine and chemokine mediated signaling pathway, $P = 0.04$).

Genes and SNPs associated with breast cancer risk

In gene-based testing, a number of genes were nominally related to overall breast cancer risk, and by ER status at $P < 0.01$ (Table 2; Supplementary Table S3). Gene functions are summarized in Supplementary Table S6. No genes remained significant after correcting for the overall number of genes tested. For overall breast cancer, the most notable genes were those involved in cytokine-related pathways, which included *IL2RB* ($P = 0.001$), *CADM1* ($P = 0.003$), *ATP6AP2* ($P = 0.006$), and *TLR6* ($P = 0.006$). Subsequent SNP testing of these genes showed five variants significant at the gene-wide level (Table 3). The top variant was rs228952 in *IL2RB* ($P_{\text{adj}} = 0.007$). In SNP-based analyses, several intronic variants in *POU2AF1* were inversely associated with risk, including rs145624147 ($P_{\text{adj}} = 0.008$).

Among premenopausal women (Table 2; Supplementary Table S4), genes most strongly related to risk were *IL21*, *SLA2*, and *CFHR1* ($P \leq 0.004$), involved in regulation of immune system processes. An intronic SNP in *SLA2* (rs221310; $P_{\text{adj}} = 0.003$) was associated with approximately 30% increased risk for each copy of the G allele. Two intergenic SNPs associated with *IL21*, rs115698762 ($P_{\text{adj}} = 0.008$), and rs143266239 ($P_{\text{adj}} = 0.01$), increased risk by approximately two-fold. Among postmenopausal women, genes associated with overall risk included *IL2RB*, *CYP4F11*, and *POUF2AF1* ($P < 0.007$), as well as *IL21* ($P = 0.004$). Two genes, *CASP8* ($P = 0.006$) and *IRAK2* ($P = 0.008$) were specific to overall breast cancer risk in postmenopausal women. When SNPs were examined, several in genes involved in interleukin and inflammatory response pathways, that is, *IL2RB*, *CYP4F11*, *IRAK2*, and *IL21* were associated with risk. Of these, the most significant were in *IL21* (rs2390350, $P_{\text{adj}} = 0.008$; rs17005895, $P_{\text{adj}} = 0.01$).

Genes and SNPs associated with risk or ER⁺ and ER⁻ cancers

None of the genes associated with overall breast cancer risk were significantly associated with ER⁺ cancers at $P \leq 0.01$. Only *BCL3*, a putative proto-oncogene important for the development, survival, and activity of adaptive immune cells (22) contained a variant (rs34698726) associated with overall breast cancer risk ($P_{\text{adj}} = 0.01$) and the risk of ER⁺ cancers ($P_{\text{adj}} = 0.01$). Except for *CADM1* ($P = 0.006$), none of the genes associated with overall breast cancer risk were associated with ER⁻ cancers. Instead, genes involved in TNFR1 (*PRKDC*, $P = 0.002$) and inflammatory response pathways (*IL8*, $P = 0.003$; *NFATC3*, $P = 0.008$), regulation of cytokine production (*CARD11*, $P = 0.007$), and *MAPK* signaling pathways (*MAPK3*, $P = 0.006$) were related to risk of ER⁻ breast cancers. Several SNPs in *IL8* (rs188246983, rs113973067), *PRKDC* (rs148411126, rs8178033, rs56411879), and *MAP3K1* (rs863839, rs191188130) were associated with risk of ER⁻ breast cancer.

Hong et al.

Table 2. Nominally significant genes associated with overall, ER⁺, ER⁻ breast cancer risk in the AMBER Consortium

Gene	Chromosomal location	Pathway ^a	N SNPs		P			
			Total	Pruned In	All	ER ⁺	ER ⁻	ER vs. ER ⁺
All women								
IL2RB	22q13	INTERLEUKIN BINDING ^b	322	220	0.001	0.074	0.018	0.485
CADM1	11q23.2	CYTOKINE PRODUCTION	1,605	638	0.003	0.031	0.006	0.957
ATP6AP2	Xp11.4	POSITIVE REGULATION OF CYTOKINE PRODUCTION ^d	283	96	0.006	0.130	0.091	0.689
TLR6	4p16.1	CYTOKINE METABOLIC PROCESS	494	141	0.006	0.142	0.024	0.147
HLA-DQA1	6p21.3	REACTOME PD1 SIGNALING [Programmed death-1 (PD-1) is a cell surface molecule that regulates the adaptive immune response]	1,986	188	0.007	0.043	0.079	0.361
POU2AF1	11q23.1	HUMORAL IMMUNE RESPONSE	244	94	0.009	0.033	0.338	0.881
TLR7	Xp22.3	BIOCARTA DC PATHWAY	251	159	0.009	0.330	0.563	0.949
CYP4F11	19p13.1	INFLAMMATORY RESPONSE	336	122	0.010	0.068	0.231	0.910
UBE2N	12q22	POSITIVE REGULATION OF IMMUNE RESPONSE	227	66	0.098	0.001	0.378	0.345
IFNA1	9p22	BIOCARTA INFLAM PATHWAY	143	77	0.014	0.002	0.146	0.770
TRAF1	9q33-q34	BIOCARTA TNFR2 PATHWAY ^c	258	62	0.377	0.003	0.725	0.001
CALCA	11p15.2	CYTOKINE PRODUCTION	129	49	0.034	0.005	0.015	0.124
MAP3K1	5q11.2	BIOCARTA TNFR1 PATHWAY	544	141	0.096	0.007	0.015	0.007
BCL3	19q13.32	CYTOKINE PRODUCTION	214	131	0.038	0.007	0.205	0.422
ICOSLG	21q22.3	REGULATION OF LYMPHOCYTE ACTIVATION	264	148	0.018	0.009	0.574	0.225
PRKDC	8q11	BIOCARTA TNFR1 PATHWAY	444	109	0.131	0.488	0.002	0.052
IL8	4q13-q21	BIOCARTA INFLAM PATHWAY	122	49	0.395	0.644	0.003	0.570
MAPK3	16p11.2	KEGG NOD LIKE RECEPTOR SIGNALING PATHWAY	24	19	0.688	0.953	0.006	0.019
CARD11	7p22	REGULATION OF CYTOKINE PRODUCTION ^d	1,471	828	0.069	0.665	0.007	0.099
NFATC3	16q22	INFLAMMATORY RESPONSE	660	117	0.898	0.948	0.008	0.147
Premenopausal women								
IL21	4q26-q27	POSITIVE REGULATION OF IMMUNE SYSTEM PROCESS ^b	75	32	0.001	0.019	0.448	0.954
SLA2	20q11.23	POSITIVE REGULATION OF IMMUNE SYSTEM PROCESS ^b	220	80	0.001	0.076	0.122	0.704
TLR7	Xp22.3	CYTOKINE BIOSYNTHETIC PROCESS	251	159	0.003	0.038	0.311	0.500
CFHR1	1q32	REGULATION OF IMMUNE SYSTEM PROCESS ^b	500	123	0.004	0.034	0.210	0.621
PRG3	11q12.1	CYTOKINE BIOSYNTHETIC PROCESS	93	39	0.006	0.004	0.272	0.211
CXCR4	2q21	INFLAMMATORY RESPONSE	90	54	0.008	0.024	0.228	0.060
HLA-DMA	6p21.3	REACTOME PD1 SIGNALING [Programmed death-1 (PD-1) is a cell surface molecule that regulates the adaptive immune response] ^c	135	40	0.009	0.002	0.623	0.455
MAP3K1	5q11.2	BIOCARTA TNFR2 PATHWAY	544	141	0.187	0.003	0.595	0.025
LBP	20q11.23	INFLAMMATORY RESPONSE	389	196	0.012	0.006	0.608	0.613
HLA-DOA	6p21.3	REACTOME PD1 SIGNALING [Programmed death-1 (PD-1) is a cell surface molecule that regulates the adaptive immune response] ^c	303	136	0.060	0.008	0.261	0.474
IL10	1q31-q32	BIOCARTA CYTOKINE PATHWAY ^d	114	67	0.442	0.498	0.001	0.021
IL8	4q13-q21	KEGG NOD LIKE RECEPTOR SIGNALING PATHWAY	122	49	0.327	0.732	0.001	0.110
HELLS	10q24.2	LYMPHOCYTE ACTIVATION	344	77	0.863	0.982	0.008	0.098
Postmenopausal women								
IL2RB	22q13	INTERLEUKIN RECEPTOR ACTIVITY ^b	322	220	0.001	0.023	0.008	0.137
CYP4F11	19p13.1	INFLAMMATORY RESPONSE	336	122	0.001	0.005	0.213	0.985
IL21	4q26-q27	REGULATION OF LYMPHOCYTE ACTIVATION	75	32	0.004	0.225	0.237	0.912
CASP8	2q33-q34	KEGG NOD LIKE RECEPTOR SIGNALING PATHWAY	259	122	0.006	0.014	0.100	0.572
POU2AF1	11q23.1	HUMORAL IMMUNE RESPONSE	244	94	0.007	0.012	0.335	0.787
IRAK2	3p25.2	INFLAMMATORY RESPONSE	645	317	0.008	0.478	0.001	0.631
XCR1	3p21.3-p21.1	CYTOKINE BINDING	95	31	0.086	0.002	0.296	0.988
CD274	9p24.1	REACTOME PD1 SIGNALING [Programmed death-1 (PD-1) is a cell surface molecule that regulates the adaptive immune response]	130	73	0.113	0.005	0.524	0.532
UBE2N	12q22	POSITIVE REGULATION OF IMMUNE RESPONSE	227	66	0.013	0.006	0.455	0.200
AOAH	7p14.2	INFLAMMATORY RESPONSE	1,608	770	0.030	0.006	0.465	0.911
IL1R1	2q12	INTERLEUKIN RECEPTOR ACTIVITY ^b	858	334	0.022	0.009	0.630	0.874
CARD11	7p22	REGULATION OF CYTOKINE PRODUCTION	1,471	828	0.554	0.911	0.004	0.011
SMAD3	15q21-q22	CYTOKINE PRODUCTION	837	448	0.094	0.116	0.006	0.051
CD247	1q24.2	REACTOME PD1 SIGNALING [Programmed death-1 (PD-1) is a cell surface molecule that regulates the adaptive immune response]	682	367	0.284	0.242	0.008	0.238

NOTE: P-values for genes significant at $P \leq 0.01$ are indicated in bold text.^aPathway associated with the greatest statistical significance for gene as shown in Supplementary Tables S3-S5.^bPathway associated with overall breast cancer risk within study population specified (i.e., all women combined, premenopausal women, or postmenopausal women), as shown in Supplementary Tables S3-S5 ($P < 0.05$).^cPathway associated with risk of ER⁺ breast cancer risk within study population specified, as shown in Supplementary Tables S3-S5 ($P < 0.05$).^dPathway associated with ER⁻ breast cancer within study population specified, as shown in Supplementary Tables S3-S5 ($P < 0.05$).**Genes and SNPs associated with risk of ER⁺ and ER⁻ cancers by menopausal status**

Among premenopausal women, genes associated with overall breast cancer risk were also associated with ER⁺ cancers, but not

ER⁻ disease (Table 2; Supplementary Table S4). None of the genes associated with overall risk of ER⁻ breast cancer were observed when limited to premenopausal women, except for *IL8*, with two significant variants (rs113976067, rs188246983). The genes *IL10*

Table 3. Associations between SNPs reaching gene-wide significance with overall, ER⁺, and ER⁻ breast cancer risk in the AMBER consortium

Gene	SNP	Function	Ref/variant allele	MAF	PLINK INFO score	All cases		ER ⁺		ER ⁻		
						Per allele OR (95% CI)	Gene-wide P _{adj}	Per allele OR (95% CI)	Gene-wide P _{adj}	Per allele OR (95% CI)	Gene-wide P _{adj}	
All women	rs228952	Intronic	G/T	0.28	0.97	0.85 (0.79-0.92)	0.007	0.88 (0.80-0.96)	0.967	0.80 (0.71-0.90)	0.051	
	rs145624147	Intronic	CAG/C	0.24	1.01	0.85 (0.79-0.92)	0.008	0.88 (0.80-0.96)	0.549	0.82 (0.73-0.93)	0.149	
	rs1815948	Intronic	G/C	0.14	1.01	0.83 (0.75-0.92)	0.023	0.85 (0.76-0.96)	0.868	0.83 (0.71-0.93)	1.000	
	rs7698807	Intronic	T/C	0.08	1.00	0.80 (0.70-0.90)	0.035	0.79 (0.67-0.91)	0.180	0.86 (0.71-1.03)	1.000	
	rs12722043	Exonic syn SNV	C/T	0.25	0.98	0.85 (0.79-0.92)	0.010	0.86 (0.79-0.94)	0.287	0.83 (0.74-0.94)	0.486	
	rs15976249	Intronic	G/A	0.07	0.94	0.76 (0.66-0.87)	0.012	0.77 (0.65-0.91)	0.369	0.75 (0.61-0.92)	1.000	
	rs141273518	Intronic	C/T	0.05	0.86	1.39 (1.18-1.65)	0.017	1.33 (1.08-1.63)	1.000	1.63 (1.27-2.08)	0.015	
	rs141628846	Upstream	G/A	0.02	0.97	1.68 (1.28-2.20)	0.024	1.61 (1.16-2.22)	0.556	2.02 (1.39-2.96)	0.037	
	rs115047524	Intronic	G/A	0.09	0.98	0.80 (0.72-0.90)	0.025	0.82 (0.71-0.95)	0.639	0.80 (0.67-0.96)	1.000	
	rs176526843	Intronic	G/T	0.02	0.90	0.64 (0.49-0.83)	0.028	0.74 (0.54-1.00)	1.000	0.65 (0.42-0.98)	1.000	
CADMI	rs73570052	Intronic	A/C	0.04	0.97	1.43 (1.20-1.70)	0.052	1.41 (1.14-1.74)	0.843	1.58 (1.23-2.05)	0.274	
	rs34698726	Intergenic	A/T	0.32	0.91	1.16 (1.08-1.25)	0.010	1.19 (1.09-1.30)	0.013	1.11 (0.99-1.25)	1.000	
	rs113976067	Intergenic	T/C	0.07	0.99	1.17 (1.03-1.34)	0.914	1.14 (0.97-1.34)	1.000	1.49 (1.23-1.81)	0.002	
	rs863839	Intronic	A/G	0.08	1.00	1.10 (1.01-1.19)	1.000	1.08 (0.98-1.19)	1.000	1.23 (1.09-1.39)	0.047	
	rs191188130	Intronic	G/T	0.09	0.87	0.96 (0.84-1.09)	1.000	1.08 (0.93-1.25)	1.000	0.65 (0.53-0.81)	0.003	
	rs14841126	Intronic	C/CT	0.08	0.98	1.03 (0.91-1.17)	1.000	0.95 (0.81-1.11)	1.000	1.32 (1.10-1.58)	0.052	
	rs8178033	Exonic nonsyn SNV	G/C	0.08	1.00	1.16 (1.03-1.32)	1.000	1.03 (0.89-1.19)	1.000	1.45 (1.22-1.72)	0.003	
	rs56411879	Intronic	T/C	0.02	0.86	1.15 (1.02-1.30)	1.000	1.02 (0.88-1.19)	1.000	1.41 (1.19-1.68)	0.011	
Premenopausal	rs221310	Intronic	A/G	0.73	gtyped	1.32 (1.16-1.51)	0.003	1.26 (1.08-1.48)	0.354	1.28 (1.06-1.56)	0.949	
	rs17848049	Intergenic	G/C	0.09	1.00	0.65 (0.52-0.80)	0.004	0.59 (0.45-0.78)	0.009	0.74 (0.54-1.01)	1.000	
	rs4411290	Intergenic	C/G	0.47	1.00	0.79 (0.70-0.89)	0.007	0.76 (0.66-0.89)	0.016	0.84 (0.71-1.00)	1.000	
	rs1867128	Intergenic	A/T	0.51	1.00	0.81 (0.72-0.92)	0.039	0.78 (0.67-0.91)	0.043	0.86 (0.72-1.03)	1.000	
	rs115698762	Intergenic	C/T	0.02	0.94	2.22 (1.45-3.41)	0.008	2.19 (1.33-3.60)	0.064	2.02 (1.07-3.79)	0.934	
	rs143266239	Intergenic	G/A	0.04	0.97	1.77 (1.29-2.43)	0.013	1.80 (1.25-2.60)	0.053	1.67 (1.06-2.65)	0.914	
	rs580962	Intergenic	C/T	0.78	gtyped	1.29 (1.12-1.49)	0.016	1.42 (1.19-1.70)	0.004	1.19 (0.96-1.46)	1.000	
	rs2232587	Intronic	T/C	0.11	0.96	0.68 (0.56-0.83)	0.025	0.64 (0.50-0.82)	0.079	0.73 (0.55-0.98)	1.000	
	rs252911	Intergenic	A/G	0.83	0.99	1.22 (1.04-1.43)	1.000	1.45 (1.18-1.77)	0.043	1.03 (0.82-1.29)	1.000	
	chr1:206953202:1	Intergenic	T/TC	0.11	0.92	1.21 (1.00-1.48)	1.000	1.06 (0.83-1.35)	1.000	1.73 (1.32-2.26)	0.005	
IL8	rs140929284	Intergenic	TC/T	0.06	0.88	1.27 (0.98-1.63)	1.000	1.10 (0.80-1.52)	1.000	1.88 (1.35-2.63)	0.015	
	rs74148793	Intergenic	C/T	0.15	0.99	1.21 (1.02-1.42)	0.990	1.12 (0.92-1.37)	1.000	1.50 (1.19-1.88)	0.035	
	rs113976067	Intergenic	T/C	0.21	1.00	1.20 (1.04-1.38)	0.660	1.15 (0.97-1.37)	1.000	1.50 (1.21-1.85)	0.008	
	rs188246983	Intergenic	T/C	0.07	0.99	1.25 (0.99-1.58)	1.000	1.20 (0.91-1.60)	1.000	1.78 (1.29-2.45)	0.022	
	rs143193835	Intronic	G/C	0.03	0.99	1.74 (1.22-2.47)	0.091	1.45 (0.94-2.23)	1.000	2.65 (1.69-4.17)	0.016	
	rs200175744	Intergenic	ACT/A	0.32	0.96	1.11 (0.97-1.26)	1.000	1.03 (0.88-1.20)	1.000	1.43 (1.18-1.72)	0.017	
	rs10882476	Intronic	T/G	0.09	0.99	1.05 (0.85-1.30)	0.990	0.86 (0.66-1.13)	1.000	1.64 (1.23-2.17)	0.049	
	rs11188009	Intergenic	A/G	0.47	0.98	1.08 (0.96-1.22)	1.000	1.00 (0.87-1.16)	1.000	1.38 (1.16-1.66)	0.032	
Postmenopausal	rs75716067	Intronic	A/G	0.03	1.02	0.53 (0.40-0.71)	0.002	0.46 (0.32-0.66)	0.003	0.60 (0.38-0.93)	1.000	
	rs15976249	Intronic	G/A	0.07	0.94	0.66 (0.54-0.80)	0.004	0.70 (0.56-0.88)	0.359	0.61 (0.45-0.83)	0.332	
	rs2390350	Intronic	A/G	0.51	1.00	1.19 (1.08-1.30)	0.008	1.16 (1.04-1.29)	0.236	1.16 (1.00-1.33)	1.000	
	rs17005895	Intergenic	A/T	0.15	0.97	1.27 (1.11-1.44)	0.014	1.21 (1.03-1.41)	0.363	1.32 (1.08-1.61)	0.160	
	rs4572524	Intergenic	A/G	0.65	0.87	0.82 (0.73-0.90)	0.015	0.84 (0.74-0.95)	0.560	0.80 (0.68-0.93)	0.659	
	rs73406995	Intergenic	C/G	0.16	0.97	0.78 (0.68-0.88)	0.021	0.77 (0.66-0.90)	0.186	0.71 (0.58-0.87)	0.207	

(Continued on the following page)

Table 3. Associations between SNPs reaching gene-wide significance with overall, ER⁺, and ER⁻ breast cancer risk in the AMBER consortium (Cont'd)

Gene	SNP	Function	Ref/variant allele	MAF	PLINK INFO score	All cases			ER ⁺			ER ⁻		
						Per allele OR (95% CI)	Gene-wide P _{adj}	Per allele OR (95% CI)	Gene-wide P _{adj}	Per allele OR (95% CI)	Gene-wide P _{adj}	Per allele OR (95% CI)	Gene-wide P _{adj}	
UBE2N	rs76506230	Intergenic	T/C	0.02	0.77	2.12 (1.40–3.22)	0.026	2.52 (1.57–4.03)	0.008	2.13 (1.18–3.86)	0.814			
CASP8	rs55637196	Intronic	G/A	0.20	0.99	0.81 (0.72–0.91)	0.034	0.78 (0.68–0.90)	0.065	0.81 (0.67–0.97)	1.000			
IRAK2	rs149858020	Intronic	C/T	0.13	0.99	1.31 (1.14–1.50)	0.038	1.23 (1.05–1.45)	1.000	1.51 (1.24–1.85)	0.017			
CD274	rs2890657	Intronic	G/C	0.04	0.97	0.75 (0.59–0.95)	1.000	0.56 (0.42–0.76)	0.016	0.84 (0.59–1.20)	1.000			
CALCA	rs10481593	Intronic	G/A	0.24	gtyped	0.87 (0.78–0.97)	0.655	0.80 (0.70–0.91)	0.040	0.91 (0.77–1.07)	1.000			
XCR1	rs34587547	Exonic nonsyn SNV	C/G	0.01	0.99	2.23 (1.26–4.29)	0.130	3.20 (1.67–6.16)	0.024	1.50 (0.54–4.13)	1.000			
	rs2373148	Intergenic	T/C	0.85	1.01	0.85 (0.75–0.96)	0.361	0.78 (0.67–0.90)	0.026	0.89 (0.73–1.07)	1.000			
	rs2371	Upstream	A/G	0.91	0.98	0.82 (0.70–0.96)	0.382	0.73 (0.61–0.88)	0.033	0.85 (0.67–1.08)	1.000			
PRKDC	rs8178153	Intronic	C/T	0.08	0.96	1.26 (1.06–1.50)	1.000	1.09 (0.88–1.34)	1.000	1.60 (1.24–2.06)	0.028			
CD247	rs12066323	Intronic	G/A	0.13	0.98	1.12 (0.97–1.28)	1.000	0.98 (0.83–1.15)	1.000	1.49 (1.22–1.82)	0.037			
SMAD3	rs2289259	Intronic	G/A	0.34	1.00	1.08 (0.98–1.19)	1.000	1.00 (0.90–1.12)	1.000	1.34 (1.15–1.55)	0.049			

NOTE: P-values for SNPs with corrected gene-wide significance at $P \leq 0.05$ are indicated in bold text.

Abbreviations: gtyped: genotyped; MAF, minor allele frequency; Ref, referent; nonsyn SNV, nonsynonymous single nucleotide variant; syn SNV, synonymous single nucleotide variant.

($P = 0.001$) and *HELLS* ($P = 0.008$) were also associated with risk of premenopausal ER⁻ cancer, with ORs for SNPs associated with *IL10* ranging from 1.5 to 1.8 for each additional copy of the variant allele. Several SNPs within the *HELLS* gene were significant (rs200175744, $P_{\text{adj}} = 0.02$; rs10882476; $P_{\text{adj}} = 0.05$; rs11188009, $P_{\text{adj}} = 0.03$).

Similar to the pattern observed in premenopausal women, genes associated with overall breast cancer risk in postmenopausal women were also associated with ER⁺ disease (*IL2RB*, *CYP4F11*, *CASP8*, *POU2AF1*, $P \leq 0.01$; Table 2). Several SNP variants in *POU2AF1*, *UBE2N*, *CD274*, and *XCR1* were associated with postmenopausal ER⁺ risk, including rs75716067 in *POU2AF1* ($P_{\text{adj}} = 0.003$). The genes *IL2RB* ($P = 0.008$) and *IRAK2* ($P = 0.001$), associated with overall breast cancer risk in postmenopausal women, were also associated with ER⁻ cancers. Generally, few SNPs were significantly associated with postmenopausal ER⁻ breast cancer. The most significant were intronic SNPs in *IRAK2* (rs149858020, $P_{\text{adj}} = 0.02$) and *PRKDC* (rs8178153, $P_{\text{adj}} = 0.03$).

Case–case analyses


In case–case analyses comparing odds of being diagnosed with ER⁻ versus ER⁺ breast cancers, the only genes showing heterogeneity between ER⁻ and ER⁺ cancers for all women combined were *TRAF1* ($P = 0.001$) and *MAP3K1* ($P = 0.007$). There was also some indication that *MAPK3* ($P = 0.02$) plays a more pronounced role in ER⁻ cancers. No heterogeneity were observed in pre- or postmenopausal women at the $P = 0.01$ level. Of the above genes, only rs863839 in *MAP3K1* was significantly different in ER⁻ versus ER⁺ disease among all women combined (OR = 0.66; 95% CI: 0.52–0.82, $P_{\text{adj}} = 0.04$), with the study having 80% power to detect this level of difference.

PPI networks

In the PPI network for all significant immunity genes identified at $P \leq 0.01$ ($N = 38$; Supplementary Table S6), there were a total of 22 genes with proteins (nodes) that were connected to at least one other protein (Fig. 1). The greatest number of protein–protein interactions were observed with *IL2RB* (node degree = 5), *MAPK3* (node degree = 4), *MAP3K1* (node degree = 6), *IL8* (node degree = 5), and *IL10* (node degree = 5). For ER⁺ cancers ($N = 29$ genes), *MAP3K1* showing the most protein–protein interactions (node degree = 4), while *IL2RB* interacted with the most proteins (node degree = 4) for ER⁻ cancers ($N = 26$ genes).

Discussion

The immune system plays a critical role in preventing and inhibiting tumor development, but may also act to promote tumor growth and progression (23). In these analyses, we used a comprehensive approach to investigate the role of hereditary immunity on breast cancer risk in AA women by assessing multiple immune response pathways, individual genes within these pathways, and the contribution of specific gene variants. Pathways and genes associated with regulation of immune system processes, immune activation, and inflammation were associated with overall breast cancer, with genes and SNPs in the NF κ B pathway associated with innate immunity and activation of the inflammatory response playing a role in ER⁺ breast cancers and pathways associated with *MAP3K1* activation playing a role in ER⁻ cancers. Of the genes identified by these analyses, only *MAP3K1* was previously identified as a breast cancer GWAS locus, and

**Figure 1.**

Protein network for all genes associated with breast cancer risk overall (N genes = 38), and for ER⁺ (N genes = 29) and ER⁻ (N genes = 26) breast cancers, in either all women combined, or in pre- or postmenopausal women only. Gene-gene interactions are represented by interconnecting lines, with edge combined scores ≥ 0.7 . Disconnected nodes are not shown.

previously reported to be related to ER⁻ cancer in the AMBER consortium (see Haddad and colleagues for discussion; ref. 15). We also noted that a number of genes related to dysregulation of humoral immunity, and involved in autoimmune and atopic disorders were associated with risk.

Th2-related and PD-1 immunosuppressive pathways relevant to both ER⁺ and ER⁻ cancers

Individuals and populations vary in their resistance to infectious disease and pathogens. Much of this variation in phenotype is genetic, with pathogens acting as a selective force on genetic diversity (1, 24). Studies in West Africa show inheritance of immune phenotypes (3). The higher prevalence of bias toward a stronger T-helper type 2 (Th2) immune response among AAs is due in part to evolutionary responses to endemic helminth exposure in Sub-Saharan Africa, and is traditionally viewed as immunosuppressive, favoring tumor growth by inhibiting cell-mediated immunity and promoting angiogenesis. This immune bias contributes to higher incidence of atopic conditions such as asthma, which is characterized by chronic activation of humoral immunity pathways (25) and can lead to allergy-driven inflammatory cytokines (25, 26). In this study, a number of genes identified in relation to breast cancer risk were previously associated with asthma in GWAS studies, including *IL2RB*, *HLA-DQ*,

and *SMAD3*, the main mediator of TGF β signaling, which is important in T-cell activation and immune tolerance (27, 28), and promotes a Th2 immune phenotype (29). The variant rs2289259 in *SMAD3* was strongly related to increased risk of ER⁻ cancer among postmenopausal women and is in high LD with rs2033784 ($r^2 = 0.81$), which affects *SMAD3* expression in thyroid tissue and blood (18).

IL5 was the Th2-related pathway most strongly related to breast cancer risk in these analyses, and is the most important driver of eosinophil production and host defense against helminth parasite infections. Excessive production of IL5 and eosinophilia would be expected to dampen antitumor immune responses and increase allergy-related inflammation. Among postmenopausal women, the Biocarta IL5 pathway was significantly related to ER⁺ breast cancer, with significant associations observed for *IL5*, *IL5RA*, and *CCR3* in gene-level analyses ($P < 0.05$). No significant variants, however, were found in any of these genes. These findings were similar to those in a recent pooled analysis of approximately 40,000 cases and 40,000 controls in the Breast Cancer Association Consortium, which found an association with *IL5* in gene-level analyses, but did not identify individual SNPs associated with risk (30). *PRG3*, coding for proteoglycan 3, an eosinophil protein that is involved in the positive regulation of IL8, histamine, and

Hong et al.

leukotriene C4 release was also associated with breast cancer risk, but only among premenopausal women. The *PRG3* variant rs4411290 and rs1867128 are eQTL sites and are associated with increased *PRG2* expression in adipose tissue (31). These findings support the original hypothesis that a type 2, immunosuppressive phenotype plays a role in breast cancer risk among AA women, although it appears that this phenotype may be relevant for both ER⁺ and ER⁻ cancers.

In further support of a potential role for immunosuppression and breast cancer risk was the identification of several genes involved in PD-1 signaling, including among postmenopausal women an association between ER⁺ cancers and CD274, the primary ligand for the PD-1 checkpoint molecule on T cells, as well as an association between ER⁻ cancers and CD247, which is involved in the downregulation of T-lymphocytes due to chronic inflammation, and has been identified as a susceptibility locus for systemic sclerosis and rheumatoid arthritis (32, 33).

Association with genes linked to autoimmune disease and role of B-cell pathways

Several of the genes associated with breast cancer risk in our study have previously been associated with autoimmune diseases in GWAS studies, pointing to the possibility that dysregulated immune responses associated with autoimmune conditions, including chronic activation and proliferation of B cells, can alter breast cancer risk in AA women. Genes identified include *IL21*, *HLA-DQA1*, *HLA-DMA*, *TRAF1*, *ICOSLG*, and *SMAD3*. The *IL21* rs2390350 variant associated with increased risk of breast cancer among postmenopausal women is in high LD with rs907715 ($r^2 = 0.82$), an independent susceptibility locus for systemic lupus erythematosus in both AAs and EAs (34, 35), and was found to be borderline significant in our analyses (OR = 0.86; 95% CI: 0.79–0.95; $P_{\text{adj}} = 0.072$). The gene *PRKDC* involved in regulation of autoimmune responses (36), as well as T-cell tolerance, inflammatory disease, and DNA repair was the gene most strongly related to risk of ER⁻ breast cancer. The missense variant (rs8178033) identified in this gene, however, was considered to be benign by PolyPhen-2 (Score: 0.29). Given that AA women are more likely to be diagnosed with systemic autoimmune diseases compared with EAs (37), greater understanding is needed of how these pathways might impact breast cancer risk in these women.

Several variants in genes that play a role in B-cell activation and development were also noted in this study. This included SNPs in *POU2AF1*, a B-cell specific transcriptional coactivator required for B-cell maturation (38), and a polymorphism in *SLA2*, which encodes for a SRC-like adaptor protein (SLAP) required for maintaining normal levels of B-cell receptor expression and development that is associated with several oncogenic signaling pathways and rheumatoid arthritis (39). The variant rs17848049 in the *CXCR4* gene region was found to be associated with premenopausal breast cancer. The CXCR4 receptor binds to stromal cell-derived factor 1 and controls B-cell development, can activate inflammatory signaling pathways, and is dysregulated in autoimmune conditions (40). A recent finding from the Women's Health Initiative study of a specific prediagnostic autoimmune response signature related to humoral immunity in triple-negative breast cancers provide some support for B-cell pathways playing a role in breast cancer etiology (41), although our findings support an association for B-cell pathways in both ER⁺ and ER⁻ cancers among AAs.

Association with inflammation related to innate immunity

In this study, the associations found between *MAP3K1*, *IL1R1*, *IRAK2*, and *TRAF1* in the PPI network generated for ER⁺ cancers supports a role for inflammation related to innate immunity and the NF κ B pathway as a contributor to disease. MAP3K1 activates CHUK and IKK β , the central protein kinases of the NF κ B pathway, and IL1R1 mediates IL1-dependent activation of NF κ B and MAPK, and strongly induces IL8 expression, a major mediator of the inflammatory response (42). Signaling involves recruitment of adaptor proteins such as IRAK2, and *IRAK2*, *MAPK3*, and *IL8* were all found to be associated with increased risk of ER⁻ breast cancer. Associations with ER⁻ disease were also noted with other genes in innate immunity pathways including *HELLS* in premenopausal women. All three variants identified in *HELLS* (rs200175744, rs10882476, rs11188009) are associated with increased gene expression (31). Our findings are consistent with recent results indicating that inflammation is associated with breast cancer development (43).

Role for IL2, IL15, and IL21 signaling pathways

The closely related *IL2*, *IL15*, and *IL21* signaling pathways were found to play a prominent role in breast cancer risk in this study, and are involved in leukocyte development, and immune response activation and cessation. Signal transduction occurs via the Janus Kinase (JAK)–STAT pathway, the P13K–AKT pathway, and the MAPK pathway (reviewed in refs. 44, 45). IL21 triggers rapid activation of ERK1/2 required for IL21-induced cytokine production, which includes IL1, IL8, CCL3, CCL5, and CCL11 (46), and induces differentiation and production of cytotoxic T cells, macrophages, NK, and B cells. This is the first study to observe associations between breast cancer risk and *IL21* variants.

IL2RB, coding for one of 3 subunits in the IL2 receptor complex, was the most significant gene associated with overall breast cancer risk, and found to be relevant for both ER⁺ and ER⁻ cancers in postmenopausal women, with rs228952 being the top variant identified. Generally, few studies have examined the role of IL2 receptors in breast cancer. The high affinity IL2 receptor composed of a β , γ , and α subunit binds to IL2 and is highly expressed on Treg cells (47). Increased ratios of Treg cells to total T lymphocytes, an important determinant of immune suppression, was recently shown in a case-cohort study in EPIC-Heidelberg to be associated with increased risk of ER⁻ breast cancers (7). Dysregulation of IL15 expression, potentially mediated by altered IL-2R $\beta\gamma$ function, may alternatively contribute to autoimmune diseases by inhibiting Treg-mediated self-tolerance (44, 48). In this study, there were suggestions that the *IL15RA* gene was associated with ER⁻ breast cancer, with significant heterogeneity by ER status ($P = 0.02$), but no SNPs were found to be significant.

Study limitations

Despite this being the first large-scale study of AAs that comprehensively examines the role of immune-related pathways, there was still limited power to detect associations by ER and menopausal status, and to observe significant pathway and gene-level associations after correcting for multiple testing. Limited sample size may have also contributed to the observation that specific genes and variants showing the strongest associations differed by menopausal and ER status, similar to previous findings (49). These differences, however, may have been due in part to

age-related changes in immune function. Premenopausal women, for instance, are more prone to autoimmunity, while older postmenopausal women are at higher risk of chronic low-grade inflammation due to increased activation of the innate immune system (50). Most of the genetic variants of interest identified in this study were imputed, which is a potential study limitation, although we focused only on SNPs with high imputation INFO scores that achieved gene-wide significance. Regardless, this is the largest study to date in AAs that has examined the role of specific sub-components of immunity, such as the adaptive and innate immune response, inflammatory response, and cytokine-related pathways.

Summary

Results from this study represent an important extension of our understanding of how inherited genetic variation in immune pathways is relevant to breast cancer susceptibility and show the importance of pathways involved in innate immune response, immune activation, and immune suppression for both ER⁺ and ER⁻ cancers among AA women, including a role for Th2, B-cell, and PD-1-related pathways.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: C.-C. Hong, L.E. Sucheston-Campbell, S. Liu, K.P. Lee, A.F. Olshan, J.R. Palmer, C.B. Ambrosone

Development of methodology: C.-C. Hong, L.E. Sucheston-Campbell, S. Liu
Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): C.-C. Hong, S. Yao, E.V. Bandera, L.A. Rosenberg, C.A. Haiman, J.R. Palmer, C.B. Ambrosone

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): C.-C. Hong, L.E. Sucheston-Campbell, S. Liu, Q. Hu, S. Yao, K.L. Lunetta, J.T. Bensen, C.A. Haiman, S.I. Abrams

Writing, review, and/or revision of the manuscript: C.-C. Hong, L.E. Sucheston-Campbell, S. Liu, S. Yao, K.L. Lunetta, S.A. Haddad, E.A. Ruiz-Narváez, J.T. Bensen, T.-Y. David Cheng, E.V. Bandera, S.S. Evans, S.I. Abrams, E.A. Repasky, A.F. Olshan, J.R. Palmer, C.B. Ambrosone
Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): C.-C. Hong, A.F. Olshan
Study supervision: S. Liu, K.L. Lunetta, C.B. Ambrosone

Acknowledgments

The research conducted by the AMBER Consortium is funded by the NIH and Foundation grants P01 CA151135 (to C.B. Ambrosone, J.R. Palmer, A.F. Olshan), R01 CA058420 (to L. Rosenberg), UM1 CA164974 (to J.R. Palmer, L. Rosenberg), R01 CA098663 (to J.R. Palmer), R01 CA100598 (to C.B. Ambrosone), R01 CA185623 (to E.V. Bandera, C.C. Hong, K. Demissie), UM1 CA164973 (to L. Le Marchand, L.N. Kolonel, C.A. Haiman, L.R. Wilkens), R01 CA54281 (to L.N. Kolonel), R01 CA063464 (to B.E. Henderson), P50 CA58223 (to M.A. Troester, A.F. Olshan), U01 CA179715 (to M.A. Troester, A.F. Olshan), Department of Defense Breast Cancer Research Program, Era of Hope Scholar Award Program W81XWH-08-1-0383 (to C.A. Haiman), the Susan G. Komen for the Cure Foundation (to M.A. Troester, A.F. Olshan), the Breast Cancer Research Foundation (to C.B. Ambrosone), and the University Cancer Research Fund of North Carolina (to M.A. Troester, A.F. Olshan). This work was also supported by the National Cancer Institute's Cancer Center Support Grant to Roswell Park Cancer Institute (P30CA016056).

Data on breast cancer cases in the Black Women's Health Study were obtained from 24 state cancer registries (Arizona, California, Colorado, Connecticut, Delaware, District of Columbia, Florida, Georgia, Illinois, Indiana, Kentucky, Louisiana, Maryland, Massachusetts, Michigan, New Jersey, New York, North Carolina, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, and Virginia), and these results do not necessarily represent their views.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received May 18, 2017; revised August 23, 2017; accepted January 2, 2018; published OnlineFirst January 16, 2018.

References

- Cagliani R, Sironi M. Pathogen-driven selection in the human genome. *Int J Evol Biol* 2013;2013:204240.
- DeNardo DG, Coussens LM. Inflammation and breast cancer. *Balancing immune response: crosstalk between adaptive and innate immune cells during breast cancer progression.* *Breast Cancer Res* 2007;9:212.
- Quinnell RJ. Genetics of susceptibility to human helminth infection. *Int J Parasitol* 2003;33:1219-31.
- Palmer JR, Ambrosone CB, Olshan AF. A collaborative study of the etiology of breast cancer subtypes in African American women: the AMBER consortium. *Cancer Causes Control* 2014;25:309-19.
- Ambrosone CB, Zirpoli G, Hong CC, Yao S, Troester MA, Bandera EV, et al. Important role of menarche in development of estrogen receptor-negative breast cancer in African American Women. *J Natl Cancer Inst* 2015;107:pii: djv172.
- Palmer JR, Viscidi E, Troester MA, Hong CC, Schedin P, Bethea TN, et al. Parity, lactation, and breast cancer subtypes in African American women: results from the AMBER Consortium. *J Natl Cancer Inst* 2014;106:pii: dju237.
- Barth SD, Schulze JJ, Kühn T, Raschke E, Hüsing A, Johnson T, et al. Treg-Mediated immune tolerance and the risk of solid cancers: findings from EPIC-Heidelberg. *J Natl Cancer Inst* 2015;107:pii: djv224.
- Hong CC, Yao S, McCann SE, Dolnick RY, Wallace PK, Gong Z, et al. Pretreatment levels of circulating Th1 and Th2 cytokines, and their ratios, are associated with ER-negative and triple negative breast cancers. *Breast Cancer Res Treat* 2013;139:477-88.
- Ambrosone CB, Ciupak GL, Bandera EV, Jandorf L, Bovbjerg DH, Zirpoli G, et al. Conducting molecular epidemiological research in the age of HIPAA: a multi-institutional case-control study of breast cancer in African-American and European-American Women. *J Oncol* 2009;2009:871250.
- Millikan RC, Pittman GS, Newman B, Tse CK, Selmin O, Rockhill B, et al. Cigarette smoking, N-acetyltransferases 1 and 2, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 1998;7:371-8.
- Newman B, Moorman PG, Millikan R, Qaqish BF, Geradts J, Aldrich TE, et al. The Carolina Breast Cancer Study: integrating population-based epidemiology and molecular biology. *Breast Cancer Res Treat* 1995;35:51-60.
- Rosenberg L, Adams-Campbell L, Palmer JR. The Black Women's Health Study: a follow-up study for causes and preventions of illness. *J Am Med Women's Assoc* 1995;50:56-8.
- Kolonel LN, Henderson BE, Hankin JH, Nomura AM, Wilkens LR, Pike MC, et al. A multiethnic cohort in Hawaii and Los Angeles: baseline characteristics. *Am J Epidemiol* 2000;151:346-57.
- Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP. Molecular signatures database (MSigDB) 3.0. *Bioinformatics* 2011;27:1739-40.
- Haddad SA, Lunetta KL, Ruiz-Narvaez EA, Bensen JT, Hong CC, Sucheston-Campbell LE, et al. Hormone-related pathways and risk of breast cancer subtypes in African American women. *Breast Cancer Res Treat* 2015;154:145-54.
- Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* 2009;5:e1000529.
- Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al. STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 2015;43:D447-52.

Hong et al.

18. Carithers LJ, Ardlie K, Barcus M, Branton PA, Britton A, Buia SA, et al. A novel approach to high-quality postmortem tissue procurement: The GTEx Project. *Biopreserv Biobank* 2015;13:311–9.
19. Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, et al. Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res* 2012;22:1790–7.
20. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. *Nat Methods* 2010;7:248–9.
21. Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* 2012;40:D930–4.
22. Herrington FD, Nibbs RJ. Regulation of the adaptive immune response by the IkappaB family protein Bcl-3. *Cells* 2016;5:p11: E14.
23. Cavallo F, De Giovanni C, Nanni P, Forni G, Lollini PL. 2011: the immune hallmarks of cancer. *Cancer Immunol Immunother* 2011;60:319–26.
24. Ferrer-Admetlla A, Bosch E, Sikora M, Marques-Bonet T, Ramirez-Soriano A, Muntasell A, et al. Balancing selection is the main force shaping the evolution of innate immunity genes. *J Immunol* 2008;181:1315–22.
25. Holgate ST. Innate and adaptive immune responses in asthma. *Nat Med* 2012;18:673–83.
26. Kim HY, DeKruyff RH, Umetsu DT. The many paths to asthma: phenotype shaped by innate and adaptive immunity. *Nat Immunol* 2010;11:577–84.
27. Delisle JS, Giroux M, Boucher G, Landry JR, Hardy MP, Lemieux S, et al. The TGF-beta-Smad3 pathway inhibits CD28-dependent cell growth and proliferation of CD4 T cells. *Genes Immun* 2013;14:115–26.
28. Malhotra N, Kang J. SMAD regulatory networks construct a balanced immune system. *Immunology* 2013;139:1–10.
29. Sallusto F, Mackay CR, Lanzavecchia A. Selective expression of the eotaxin receptor CCR3 by human T helper 2 cells. *Science* 1997;277:2005–7.
30. Lei J, Rudolph A, Moysich KB, Behrens S, Goode EL, Bolla MK, et al. Genetic variation in the immunosuppression pathway genes and breast cancer susceptibility: a pooled analysis of 42,510 cases and 40,577 controls from the Breast Cancer Association Consortium. *Hum Genet* 2016;135:137–54.
31. GTEx Consortium. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* 2015;348:648–60.
32. Radstake TR, Gorlova O, Rueda B, Martin JE, Alizadeh BZ, Palomino-Morales R, et al. Genome-wide association study of systemic sclerosis identifies CD247 as a new susceptibility locus. *Nat Genet* 2010;42:426–9.
33. Zhemakova A, Stahl EA, Trynka G, Raychaudhuri S, Festen EA, Franke L, et al. Meta-analysis of genome-wide association studies in celiac disease and rheumatoid arthritis identifies fourteen non-HLA shared loci. *PLoS Genet* 2011;7:e1002004.
34. Qi JH, Qi J, Xiang LN, Nie G. Association between IL-21 polymorphism and systemic lupus erythematosus: a meta-analysis. *Genet Mol Res* 2015;14:9595–603.
35. Hughes T, Kim-Howard X, Kelly JA, Kaufman KM, Langefeld CD, Ziegler J, et al. Fine-mapping and transethnic genotyping establish IL2/IL21 genetic association with lupus and localize this genetic effect to IL21. *Arthritis Rheum* 2011;63:1689–97.
36. Mathieu AL, Verronese E, Rice GI, Fouyssac F, Bertrand Y, Picard C, et al. PRKDC mutations associated with immunodeficiency, granuloma, and autoimmune regulator-dependent autoimmunity. *J Allergy Clin Immunol* 2015;135:1578–88.
37. Ramos-Casals M, Brito-Zeron P, Kostov B, Siso-Almirall A, Bosch X, Buss D, et al. Google-driven search for big data in autoimmune geoepidemiology: analysis of 394,827 patients with systemic autoimmune diseases. *Autoimmun Rev* 2015;14:670–9.
38. Corcoran L, Emslie D, Kratina T, Shi W, Hirsch S, Taubenheim N, et al. Oct2 and Obf1 as Facilitators of B:T Cell Collaboration during a Humoral Immune Response. *Front Immunol* 2014;5:108.
39. Kazi JU, Kabir NN, Ronnstrand L. Role of SRC-like adaptor protein (SLAP) in immune and malignant cell signaling. *Cell Mol Life Sci* 2015;72:2535–44.
40. Wang A, Fairhurst AM, Tus K, Subramanian S, Liu Y, Lin F, et al. CXCR4/CXCL12 hyperexpression plays a pivotal role in the pathogenesis of lupus. *J Immunol* 2009;182:4448–58.
41. Katayama H, Boldt C, Ladd JJ, Johnson MM, Chao T, Capello M, et al. An autoimmune response signature associated with the development of triple-negative breast cancer reflects disease pathogenesis. *Cancer Res* 2015;75:3246–54.
42. Keating SE, Maloney GM, Moran EM, Bowie AG. IRAK-2 participates in multiple toll-like receptor signaling pathways to NFkappaB via activation of TRAF6 ubiquitination. *J Biol Chem* 2007;282:33435–43.
43. Hung RJ, Ulrich CM, Goode EL, Brhane Y, Muir K, Chan AT, et al. Cross cancer genomic investigation of inflammation pathway for five common cancers: lung, ovary, prostate, breast, and colorectal cancer. *J Natl Cancer Inst* 2015;107.
44. Boyman O, Sprent J. The role of interleukin-2 during homeostasis and activation of the immune system. *Nat Rev Immunol* 2012;12:180–90.
45. Zhang Y, Dong C. Regulatory mechanisms of mitogen-activated kinase signaling. *Cell Mol Life Sci* 2007;64:2771–89.
46. Fuqua CF, Akomeah R, Price JO, Adunyah SE. Involvement of ERK-1/2 in IL-21-induced cytokine production in leukemia cells and human monocytes. *Cytokine* 2008;44:101–7.
47. Setoguchi R, Hori S, Takahashi T, Sakaguchi S. Homeostatic maintenance of natural Foxp3(+) CD25(+) CD4(+) regulatory T cells by interleukin (IL)-2 and induction of autoimmune disease by IL-2 neutralization. *J Exp Med* 2005;201:723–35.
48. Waldmann TA. The biology of interleukin-2 and interleukin-15: implications for cancer therapy and vaccine design. *Nat Rev Immunol* 2006;6:595–601.
49. Quan L, Gong Z, Yao S, Bandera EV, Zirpoli G, Hwang H, et al. Cytokine and cytokine receptor genes of the adaptive immune response are differentially associated with breast cancer risk in American women of African and European ancestry. *Int J Cancer* 2013;134:1408–21.
50. Gubbels Bupp MR. Sex, the aging immune system, and chronic disease. *Cell Immunol* 2015;294:102–10.

Cancer Epidemiology, Biomarkers & Prevention

Genetic Variants in Immune-Related Pathways and Breast Cancer Risk in African American Women in the AMBER Consortium

Chi-Chen Hong, Lara E. Sucheston-Campbell, Song Liu, et al.

Cancer Epidemiol Biomarkers Prev 2018;27:321-330. Published OnlineFirst January 16, 2018.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-17-0434](https://doi.org/10.1158/1055-9965.EPI-17-0434)

Supplementary Material Access the most recent supplemental material at:
<http://cebp.aacrjournals.org/content/suppl/2018/01/13/1055-9965.EPI-17-0434.DC1>
<http://cebp.aacrjournals.org/content/suppl/2018/01/30/1055-9965.EPI-17-0434.DC2>

Cited articles This article cites 47 articles, 9 of which you can access for free at:
<http://cebp.aacrjournals.org/content/27/3/321.full#ref-list-1>

Citing articles This article has been cited by 1 HighWire-hosted articles. Access the articles at:
<http://cebp.aacrjournals.org/content/27/3/321.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/27/3/321>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.