

changing non-adherent Latinas' screening behaviors and promoting them to become agents of change in their communities. Limitations concern generalizability due to a non-probability based sample, and limited ability for causal inferences due to a lack of randomization.

Published online March 2, 2018.

doi: 10.1158/1055-9965.EPI-18-0048

©2018 American Association for Cancer Research.

Using Social Media to Reduce Multiple Risk Factors for CRC in Rural Appalachians: #CRCFREE

Voigts K, Adegboyega A, Bush H, Hatcher J

Purpose: To examine the efficacy, acceptability, feasibility, and applicability of a Facebook (FB) intervention designed to reduce multiple colorectal cancer (CRC) risk factors in older adults residing in rural Appalachian Kentucky. **Methods:** We piloted a 12 week FB intervention culturally tailored for older adults residing in rural Eastern Kentucky to impact CRC risk factors, including: nutrition, physical activity, and screening. Participants were aged 50+, had internet access, and were at risk for CRC. During the 12 week study, the participants received three daily posts via secret FB group regarding CRC risk factors. Demographics, dietary measures, body mass index (BMI), and CRC screening were assessed at baseline and post intervention. FB engagement and physical activity were tracked throughout the intervention. Dietary measures included the Healthy Eating Index (HEI) and Dietary Inflammatory Index (DII). Physical activity was tracked using Fitbits. Post-intervention focus group interviews were conducted to assess feasibility and acceptability. **Results:** Participants ($n = 57$) were Caucasian, aged 58 ± 6 years, predominately female (67%), and the majority reported at least a high school education (77%). Post intervention, participants experienced significant increase in HEI scores (49.94 ± 9.84 vs. 58.60 ± 12.06 , $P = < 0.01$). DII scores significantly decreased (2.44 ± 1.12 vs. 1.60 ± 1.63 , $P = 0.003$). There was no significant change in physical activity, BMI, or screening status. Participants, on average, viewed more than half of the posts. Focus group participants found FB posts to be useful and motivating. They reported that FB posts were educational and motivational. **Conclusion:** This pilot study shows promising preliminary data to support using a FB intervention in rural Appalachian older adults to decrease CRC risks. Participants were receptive to FB intervention, and FB provides a unique and accessible method for health promotion in hard to reach populations.

Published online March 2, 2018.

doi: 10.1158/1055-9965.EPI-18-0049

©2018 American Association for Cancer Research.

Prevalence and Characteristics of Liver Fibrosis Detected by Elastography: Results from the Cameron County Hispanic Cohort

Watt GP, Lee M, Pan JJ, Fallon MB, McCormick JP, Fisher-Hoch SP

Mexican Americans in south Texas have among highest rates of hepatocellular carcinoma (HCC) in the US. Non-invasive measures of liver fibrosis are needed to identify those at high risk of

HCC. The purpose of the study was to determine the prevalence of and factors associated with liver fibrosis using acoustic radiation force impulse (ARFI) elastography, an accurate and non-invasive modality, in the Cameron County Hispanic Cohort (CCHC). **Methods:** The CCHC is a well characterized population-based cohort in south Texas. Liver stiffness was measured by ARFI in 404 participants in the CCHC by two separate operators. Median liver stiffness > 1.34 m/s was considered significant (F2-F4) fibrosis. Ultrasound was used to determine presence of steatosis. Absence of HCV and HBsAg antigen, and absence of heavy drinking, was considered non-alcoholic fatty liver disease (NAFLD). We calculated design-based prevalence of fibrosis and determined clinical associations with (1) significant fibrosis and (2) simple steatosis using multinomial logistic regression. **Results:** Mean age was 51.1 years, 44.6% were male, and 29.1% had diabetes. The prevalence of significant fibrosis was 16.5%. In the fibrosis group, most (47/57, 84%) had NAFLD. We excluded participants with etiologies other than NAFLD for further analyses. After adjustment for age and sex, hypertension [Odds Ratio (OR) 3.0, 95% confidence interval (CI) 1.1–8.1], platelet count (OR 0.6, 95% CI, 0.5–1.0), and potassium level (OR 1.5, 95% CI, 1.1–2.0) were significantly associated with fibrosis, but not steatosis. HDL cholesterol (OR 0.5, 95% CI, 0.4–0.7), obesity (OR 2.2, 95% CI, 1.2–4.0) and insulin level (OR 2.1, 95% CI, 1.2–3.7) were significantly associated with steatosis, but not fibrosis. Elevated fasting glucose, diabetes, and elevated ALT levels were significantly associated with both steatosis and fibrosis. **Conclusion:** This is the first population-based application of liver elastography in the US. We find a high prevalence of fibrosis in Mexican Americans, dominated by NAFLD. Our results indicate a higher burden of fibrosis than population-based studies of liver fibrosis conducted elsewhere. We urge community intervention for the early detection of liver disease and prevention of NAFLD-related HCC.

Published online March 2, 2018.

doi: 10.1158/1055-9965.EPI-18-0050

©2018 American Association for Cancer Research.

Assessing the Feasibility of a Mentored Home-Based Vegetable Gardening Intervention among Breast Cancer Survivors in the Deep South

Bail JR, Frugé AD, Cases MG, Smith KP, Cantor AB, De Los Santos J, Locher JL, Cohen HJ, Demark-Wahnefried W

To assess the feasibility of a mentored home-based vegetable gardening intervention among Breast Cancer Survivors (BCS) residing in the Birmingham, Alabama metropolitan area. **Methods:** Using a wait-list control design, BCS were randomized to either a year-long vegetable gardening intervention or a wait-list control. Intervention participants were provided with necessary supplies and paired with a Master Gardener from the Cooperative Extension. Master Gardeners mentored participants in planning, planting, and maintaining 3 seasonal gardens over 12 months, conducted monthly home-visits, and checked in bi-weekly via telephone or email. Feasibility assessment criteria consisted of participant accrual, retention, and satisfaction rates of $\geq 80\%$. Target participant accrual was 100. Participant satisfaction data were collected after study completion via structured telephone debriefing. Descriptive statistics were conducted using SPSS V24.

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Using Social Media to Reduce Multiple Risk Factors for CRC in Rural Appalachians: #CRCFREE

K Voigts, A Adegboyega, H Bush, et al.

Cancer Epidemiol Biomarkers Prev 2018;27:354.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/27/3/354.1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link <http://cebp.aacrjournals.org/content/27/3/354.1>. Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.