

# Highlights

July 2018 • Volume 27 • Number 7

Selected Articles from This Issue

## Parents' Views on Best and Worst Reasons for HPV Vaccination

Gilkey et al. \_\_\_\_\_ Page 762

Healthcare providers use a wide variety of messages to encourage HPV vaccination, but few studies have empirically tested which messages are persuasive. Gilkey and colleagues conducted a best-worst scaling experiment to rank 11 reasons providers commonly give for guideline-consistent HPV vaccination. The data came from a national online survey of 1,177 U.S. parents of adolescents. The study found that parents ranked "cancer prevention" far higher as a reason for HPV vaccination than any of the other reasons tested. This finding held across subgroups of parents, including those with lower confidence in adolescent vaccines. Findings suggest an opportunity to simplify provider communication about HPV vaccination to focus on the issue of greatest importance to parents: cancer prevention.

## Circadian Misalignment and Hepatocellular Carcinoma

VoPham et al. \_\_\_\_\_ Page 719

Hepatocellular carcinoma (HCC) is an often-fatal cancer that is increasing in incidence across different parts of the world. Experimental evidence suggests that circadian misalignment impacts hepatocarcinogenesis. Geographic variation in light exposure, despite similar social and occupational schedules, may influence circadian systems and can be estimated by residential position in a time zone. Using the U.S. Surveillance, Epidemiology, and End Results Program, VoPham and colleagues found that increasing longitude moving east to west within a time zone, a proxy for circadian misalignment, was positively associated with HCC risk. Circadian misalignment may be a novel risk factor for HCC.

## NSAID Use and Colorectal Cancer

Kuo et al. \_\_\_\_\_ Page 737

Aspirin and nonsteroidal anti-inflammatory drugs (NSAIDs) are widely used clinically. Several studies have examined their effects on colorectal cancer (CRC) chemoprevention. However, the data in Asian populations are limited. Kuo and colleagues examined the primary preventive efficacy of aspirin and NSAIDs, separately, on CRC incidence in Taiwan by using the National Health Insurance Research Database with a large sample size. The results from this study have provided supportive evidence that aspirin or NSAID use is associated with a reduced risk of CRC incidence in an Asian population.

## Coffee, Tea, and the Oral Microbiome

Peters et al. \_\_\_\_\_ Page 814

Peters and colleagues examined the associations of coffee and tea intake with the oral bacterial ecosystem in a large cross-sectional study. The authors found that higher tea intake was associated with greater oral microbiota diversity, and shifts in overall community composition, but coffee was not associated with these microbiome parameters. As the oral microbiota may play a role in certain cancers, tea-driven changes to the oral microbiome may contribute to previously observed associations between tea and cancer.

# BLOOD CANCER DISCOVERY

## Highlights of This Issue

*Cancer Epidemiol Biomarkers Prev* 2018;27:717.

**Updated version** Access the most recent version of this article at:  
<http://cebp.aacrjournals.org/content/27/7/717>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://cebp.aacrjournals.org/content/27/7/717>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.