

Acrolein Is Involved in the Synergistic Potential of Cigarette Smoking- and Betel Quid Chewing-Related Human Oral Cancer



Han-Hsing Tsou¹, Chih-Hung Hu¹, Jin-Hui Liu², Chung-Ji Liu^{3,4}, Chien-Hung Lee^{5,6}, Tsung-Yun Liu^{1,2}, and Hsiang-Tsui Wang⁷

Abstract

Background: Cigarette smoking (CS) and betel quid (BQ) chewing are two known risk factors and have synergistic potential for the development of oral squamous cell carcinoma (OSCC) in Taiwan. The *p53* mutation characteristics in OSCC (G to A or G to T mutations) are similar to that of acrolein-induced DNA damage. Acrolein is a major cigarette-related carcinogen that preferentially causes *p53* mutations and inhibits DNA repair function in lung cancer. We hypothesize that acrolein is associated with OSCC carcinogenesis.

Methods: A total of 97 patients with OSCC and 230 healthy subjects with CS and/or BQ chewing histories were recruited. Slot blot analysis of Acr-dG adducts, an indicator of acrolein-induced DNA damage in buccal DNA, and LC/MS-MS analysis of 3-HPMA levels, urinary Acr metabolites, were performed.

Results: Our results showed that the level of Acr-dG adducts in buccal cells was 1.4-fold higher in patients with OSCC than in healthy subjects with CS and/or BQ chewing histories ($P < 0.001$). In addition, in healthy subjects, CS and BQ chewing were associated with significantly higher levels of 3-HPMA, indicating that CS and BQ chewing promotes acrolein absorption. However, 3-HPMA levels in patients with OSCC were significantly lower than those in healthy subjects, indicating impaired acrolein metabolism.

Conclusions: In this study, we provide a novel mechanism by which increased acrolein uptake and impaired metabolism may contribute to the synergistic potential of CS and BQ-induced OSCC.

Impact: Elevated acrolein-induced DNA damage (Acr-dG adducts) detected in buccal swabs may serve as an early indicator to identify patients at risk of developing OSCC.

Introduction

Oral cancer is a widely prevalent cancer worldwide and a leading cause of mortality in Southern Asia (1–3). Approximately 95% of oral cancer is oral squamous cell carcinoma (OSCC). The prevalence of the disease in different parts of the world reflects different forms and extents of exposure to etiologic agents. While cigarette smoking and alcohol consumption are the major risk factors in Western countries, betel quid (BQ) chewing and cigarette smoking (CS) are the primary causes of OSCC in Taiwan and both south and Southern Asia (4–7). Unfortunately, despite all of the advancements in the understanding of the disease process and recognition of the associated risk factors, the 5-year survival rate

remains at 50% (8). The key challenge to reduce the mortality and morbidity of this disease is to develop strategies to identify and detect OSCC when it is present at a very early stage, which will enable effective intervention and therapy. The detection of OSCC is currently based on expert clinical examinations and histologic analysis of suspicious areas, but it may be undetectable in hidden sites. Therefore, sensitive and specific biomarkers for OSCC may be helpful in screening high-risk patients (9).

Previous studies have shown that the incidence of oral cancer in Taiwan was 18-fold higher in patients who smoked than in abstainers and 89-fold higher in patients who smoked and BQ chewing than in abstainers (4). The tumor suppressor gene *p53* is frequently mutated in human cancers (10, 11), and its mutational patterns often bear the fingerprints of the etiologic carcinogens. Most notably, it has been found that >50% of aflatoxin B1-associated liver cancers have mutations in codon 249 of the *p53* gene and that *p53* mutations are concentrated at contiguous pyrimidines in sunlight-associated skin cancers (12, 13). Previous studies have shown that characteristics of *p53* mutations in OSCC are associated with CS and BQ chewing in Taiwan (14). *p53* mutations occur at exons 5–9 in nearly half of patients with OSCC, and G to A or G to T mutations are the predominant mutations that were observed and associated with CS and BQ chewing. Similar *p53* mutations in oral cavity tumors have been found (14–17). These findings suggest an important contributive role of tobacco carcinogens in *p53* mutations in a series of Taiwanese patients with OSCC.

Acrolein (Acr), an α,β -unsaturated aldehyde, is abundant in CS, cooking fumes, and automobile exhaust fumes (18). The main

¹Institute of Environmental and Occupational Health Sciences, National Yang-Ming University, Taipei, Taiwan. ²Institute of Food Safety and Health Risk Assessment, National Yang-Ming University, Taipei, Taiwan. ³Institute of Oral Biology, School of Dentistry, National Yang-Ming University, Taipei, Taiwan. ⁴Department of Oral and Maxillofacial Surgery, Mackay Memorial Hospital, Taipei, Taiwan. ⁵Department of Public Health, College of Health Sciences, Kaohsiung Medical University, Kaohsiung, Taiwan. ⁶Research Center for Environmental Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan. ⁷Department of Pharmacology, National Yang-Ming University, Taipei, Taiwan.

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Hsiang-Tsui Wang, National Yang-Ming University, Taipei 112, Taiwan. Phone: 886-2-2826-7097; Fax: 886-2-2826-4372; E-mail: htwang01@ym.edu.tw

doi: 10.1158/1055-9965.EPI-18-1033

©2019 American Association for Cancer Research.

route of Acr elimination involves the production of S-(3-hydroxypropyl)-N-acetylcysteine (3-HPMA) by conjunction with glutathione (GSH), the primary metabolite of Acr found in urine (19), which has been used as a biomarker for Acr exposure (20) and GSH has been identified as a protective scavenger of Acr. The free form of Acr induces α - and γ -hydroxy-1,N²-cyclic propano-2'-deoxyguanosine (α -OH-Acr-dG and γ -OH-Acr-dG) adducts in human cells (21, 22). It has been shown that both types of Acr-dG adducts are mutagenic and that they induce mainly G to T and G to A mutations (21–27). By mapping the Acr-dG adduct distribution at the nucleotide level in Acr-treated normal human bronchial epithelial (NHBE) cells, we found that the Acr-DNA-binding spectrum in the *p53* gene coincides with the *p53* mutational spectrum in CS-related lung cancer (28). A similar *p53* mutational spectrum between CS-related lung cancer and CS-related OSCC, such as mutational hotspots on codons 175, 248, and 273 of the *p53* gene, was observed (Supplementary Fig. S1). These findings raise the question of whether Acr-DNA adducts are responsible for *p53* mutations in CS- and BQ-induced OSCC. Previous studies clearly demonstrated the synergistic effects of CS and BQ in oral cancer (4, 6, 7). However, the underlying mechanism for the synergistic potential of BQ and CS remains elusive. To investigate the role of Acr in OSCC and the synergistic mechanisms of CS and BQ, this study was designed to compare Acr-dG adducts of buccal cells and urinary 3-HPMA levels in CS subjects and patients with OSCC either with or without BQ chewing histories.

Materials and Methods

Subjects and collection of buccal cells and urine

For patients with oral cancer, 97 patients treated at Mackay Memorial Hospital (Taipei, Taiwan) from February 2016 through August 2018 were recruited for participation in the study. All patients were interviewed uniformly by a well-trained interviewer before surgery. For each case, tumor tissue samples containing oral cavity and oropharynx neoplasms were surgically dissected and sent to the Department of Pathology for examination by the Association of Directors of Anatomic and Surgical Pathology (ADASP; ref. 29). Informed written consent was given by each participant or his (or her) relative, and our study protocol was approved by the Institutional Review Board of Mackay Memorial Hospital (Taipei, Taiwan). Experiments were conducted in accordance with the Declaration of Helsinki principles. For healthy participants, 230 healthy subjects, including 111 without a BQ chewing history, 107 with a BQ chewing history, and 12 non-smokers with a BQ chewing history, were recruited for participation in the study in cooperation with the Department of Public Health, Kaohsiung Medical University (Kaohsiung, Taiwan). All participants were interviewed by a well-trained interviewer. Buccal cells and urine samples were collected during the interviews. The questionnaire used in the interview sought detailed information on current and past cigarette smoking, alcohol consumption, and BQ chewing habits, occupational history, family disease, dietary history, and general demographic data. Informed written consent was given by each participant or his (or her) relative, and our study protocol was approved by the Institutional Review Board of Kaohsiung Medical University (Kaohsiung, Taiwan; IRB # KMH-IRB-20110270). Experiments were conducted in accordance with the Declaration of Helsinki principles. For the collection of buccal cells, subjects were asked to rinse their mouths with

water, followed by rinsing with antiseptic mouthwash and a subsequent water rinse. A trained technician placed one buccal brush against the inside of the cheek and scraped the brush against the center of a subject's cheek, applying firm pressure throughout the process. The brush was moved up and down as well as back and forth to ensure maximum cell collection, and the inside of the mouth was scraped at least 10–15 times. The procedure was repeated to obtain a second buccal swab sample on the other side of the cheek. A total of 2 buccal swabs were collected for each subject. Buccal cell and urine samples were stored at -30°C until use.

Slot blot assay for Acr-dG detection

Analysis of Acr-dG adducts in DNA samples was based on previously described methods (30, 31). Control genomic DNA was modified with Acr (0.5, 1, 2, and 5 mmol/L) at 37°C for 24 hours and purified with repeated phenol/ether extraction as Acr-dG adduct standards. After purification, modified DNA was precipitated with ethanol and dissolved in TE buffer (pH 8.0). Buccal DNA was extracted from buccal cells using Puregene buccal cell core kits (Qiagen) according to the manufacturer's instructions. Modified DNA or buccal DNA (0.25 μg) were loaded onto polyvinylidene difluoride membranes using a Bio-Dot SF micro-filtration apparatus (Bio-Rad). WesternDot 625 Western blotting kits (Invitrogen) were used for Western blot analysis according to the manufacturer's instructions. After blocking for 1 hour at room temperature in blocking buffer, the membrane was probed overnight at 4°C with anti-Acr-dG mouse mAbs (32). After washing with washing buffer to remove unbound primary antibodies, quantum dot-conjugated secondary antibodies (1:1,000 dilution) were added for 2 hours at room temperature. Ultimately, Acr-dG adducts were detected using a UVP BioDoc-It imaging system, and band density was quantified with UVP imaging software. After antibody detection, the same membrane was stained with methylene blue (Molecular Research Center) to indicate the amount of DNA.

Detection of 3-hydroxypropyl mercapturic acid and creatinine in urine

Analysis of 3-hydroxypropyl mercapturic acid (3-HPMA) and creatinine in urine was based on methods described previously (33). For analysis of 3-HPMA in urine, solid-phase extraction with Isolute ENV+ cartridges (Biotage) was used to prepare each sample before LC/MS-MS analysis. High-performance liquid chromatography (HPLC) was performed using an Agilent 1100 Series HPLC system with a quaternary pump (Agilent G1311A), a vacuum degasser (Agilent G1322A), and an autosampler (Agilent G1313A). The HPLC was directly connected to a triple quadrupole mass spectrometer (Finnigan TSQ Quantum Discovery MAX, Thermo Electron Corporation) equipped with an electrospray ionization (ESI) interface and a 10-port valve. A Hypersil GOLD aQ C18 2.1 mm \times 150 mm, 3 μm column (Thermo Fisher Scientific) was used for LC separation. For analysis of creatinine, HPLC was performed using an Agilent Series 1260 HPLC system with a binary pump (Agilent G1312B), a vacuum degasser (Agilent G1322A), and an autosampler (Agilent G1367E). The HPLC was directly connected to a variable wavelength detector (Agilent G1314F). A NUCLEODUR C18 HTec 250 mm \times 4.6 mm \times 4.6 μm column was used for LC separation. The samples were analyzed at a wavelength of 240 nm via a variable wavelength detector.

Detection of NNAL in urine

Analysis of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) in urine was based on modified methods previously described by Shah and colleagues (34). Urine samples were frozen and then stored at -30°C until analysis. Initially, urine samples were thawed, equilibrated to room temperature, and vortexed. The urine samples were centrifuged at $3,000 \times g$ for 10 minutes at 4°C , and 3 mL of the supernatant was collected. Solid-phase extraction with mixed cation exchange (MCX) cartridges (Waters) was used to prepare each sample before LC/MS-MS analysis. Each cartridge was conditioned with 2 mL of methanol, followed by 2 mL of water. The total NNAL volume, a volume of 1-mL aliquots of urine or water blanks, was added to 15-mL centrifuge tubes. Five milliliters of phosphate buffer solution (pH = 7.4) and 1,000 units of β -glucuronidase (G-0876; Sigma Chemical Co.) were added to each sample. The tubes were incubated in a shaking water bath at 37°C overnight. Spiked [13C6] NNAL (0.39 nmol) was added to each sample. The free NNAL volume, a volume of 2-mL aliquots of urine or water blanks, was added to centrifuge tubes and spiked [13C6] NNAL (0.13 nmol) was added to each sample. This mixture was loaded onto the prepared MCX cartridges. Each cartridge was washed with 1 mL of 1 mol/L HCl, followed by 1 mL of methanol and 1 mL of methanol with 25% ammonium hydroxide in water (5/5/90, v/v/v). Each cartridge was eluted with 3 mL of 25% ammonium hydroxide in methanol (5/95, v/v). The eluates were combined and dried with a rotary evaporator. Each sample was then reconstituted in 50 μL of the mobile phase. LC/MS-MS analysis was conducted with an Agilent 1200 series vacuum degasser, binary pump, well plate autosampler, and thermostatted column compartment and paired with an Agilent 6430 Triple Quad LC-MS operated in electrospray ionization (ESI) mode (Agilent Technologies, Inc.). All data were acquired and analyzed using Mass Hunter software, version B01.03. A Hypersil GOLD aQ C18 2.1 mm \times 150 mm \times 3 μm column (Thermo Fisher Scientific) was used for LC separation. The buffers used were (A) water + 0.1% formic acid and (B) acetonitrile + 0.1% formic acid. The linear LC gradient was as follows: time = 0 minutes, 0% B; time = 4 minutes, 95% B; time = 4.5 minutes, 0% B; time = 8 minutes, 0% B. Multiple reaction monitoring was used for MS analysis. The NNAL data were acquired in positive electrospray ionization (ESI) mode by monitoring the following transitions: 210/93.1 with a collision energy of 17 V. The [13C6] NNAL data were acquired in ESI mode by monitoring the following transitions: 216/98.1 with a collision energy of 21 V. The jet stream ESI interface had a gas temperature of 350°C , gas flow rate of 10 L/minute, nebulizer pressure of 30 psi, and capillary voltage of 4,000 V.

Statistical analyses

Descriptive statistics were presented as the mean \pm SD or as the number (percentage). Shapiro-Wilk test was used to test for normality. Student *t* tests were used to determine statistical significance. Pearson correlation analysis was used to analyze the correlation between Acr-related metabolites and clinical parameters. A minimum of three independent replicate experiments of slot blot analysis or isotope dilution HPLC mass spectrometry were performed to justify the use of the statistical tests. All calculated *P* values were two-tailed. Statistical significance was defined as a *P* < 0.05. All analyses were performed with the IBM SPSS Statistics software package, version 23.0.

Cell culture and Acr treatment

Human oral squamous carcinoma cells (SAS cells), were provided by Dr. Kuo-Wei Chang at the Institute of Oral Biology, School of Dentistry, National Yang-Ming University (Taipei, Taiwan), and were authenticated by short tandem repeat analysis (35). SAS cells were grown in DMEM supplemented with 10% FBS. Acr stock solution (Sigma-Aldrich) was freshly prepared before use. Cells at 70% confluency were washed with PBS buffer (137 mmol/L NaCl, 2.7 mmol/L KCl, 8 mmol/L Na_2HPO_4 , and 1.46 mmol/L KH_2PO_4 , pH 7.0) and treated with Acr (50 $\mu\text{mol/L}$) in complete culture medium for 3 hours, as indicated, at 37°C in the dark.

Measurement of total glutathione (GSH) and GSH/GSSG

For total GSH levels and GSH/GSSG, a glutathione fluorometric assay kit (BioVision) was used according to the manufacturer's instructions. Briefly, Acr-treated cells (2×10^6) were lysed in 100 μL of assay buffer, and 60 μL of each homogenate was mixed with perchloric acid and then centrifuged at $13,000 \times g$ for 2 minutes to pellet the insoluble materials. The supernatant was collected, neutralized with 20 μL of ice-cold 6N KOH, and analyzed for GSH assay. Finally, the total amount of GSH was measured using a fluorescence plate reader at excitation/emission = 340/420 nm.

Results

Subjects and collection of buccal cells and urine

Baseline sociodemographic variables are shown in Tables 1 and 2. Of the 327 participants recruited, 97 were patients with OSCC and 230 were healthy subjects. All patients with OSCC have been clinically followed for 2–3 years. The demographic data of

Table 1. Baseline sociodemographic variables of patients with OSCC (*n* = 97)

Characteristics	
Age, y	
Mean \pm SD	59.5 \pm 11.0
Range	29–89
Sex [n (%)]	
Male	82 (84.5)
Female	15 (15.5)
Site of primary tumor [n (%)] ^a	
Lip	4 (4.35)
Tongue	22 (23.9)
Buccal mucosa	27 (29.3)
Gingiva	23 (25.0)
Hard palate	6 (6.52)
Retromolar trigone	4 (4.35)
Oropharynx	5 (5.43)
Hypopharynx	1 (1.09)
Clinical stage [n (%)] ^a	
Stage I	18 (19.1)
Stage II	16 (17.0)
Stage III	6 (6.38)
Stage IV	54 (57.4)
Cigarette smoker [n (%)] ^b	75 (77.3)
Alcohol drinker [n (%)] ^b	40 (41.2)
Betel quid chewer [n (%)] ^b	74 (76.3)

^aDiagnosis case number for site of primary tumor is 92, and the case number of clinical staging is 94.

^bStudy participants were asked if they had ever smoked cigarettes, chewed BQ, and had alcohol on a regular basis (at least once a week). Those who responded "yes" to these questions were classified as tobacco, BQ, and alcohol users.

Table 2. Baseline sociodemographic variables of healthy subjects with CS or BQ chewing histories

N	CS		CS + BQ		BQ	
	111	P ^a	107	P ^a	12	P ^a
Male (%)	88.3	—	88.8	0.47	83.3	0.31
(M/F)	99/12	—	95/12	—	10/2	—
Age, y	41.5	—	42.7	0.23	44.8	0.22
Range	19–70	—	21–64	—	28–67	—
Smoking (%)	100	—	100	—	—	—
(±)	111/0	—	107/0	—	0/12	—
Smoking years	23.1	—	27.1	<0.01	—	—
Range	1–53	—	5–50	—	—	—
Drinking alcohol (%)	29.7	—	70.1	—	58.3	—
(±)	33/78	—	75/32	—	7/5	—
Chewing betel quid (%)	—	—	100	—	100	—
(±)	—	—	107/0	—	12/0	—
Chewing betel days	—	—	34.9	—	23.4	—

^aThe $P < 0.05$ versus smoking (CS) by Student t test.

the patients are shown in Table 1. Advanced stage III and IV cancer was diagnosed in 63.8% (60/94) of patients. The most common primary tumor site was the buccal mucosa. A total of 77.3% (75/97) of the patients smoked, 41.2% (40/97) were users of alcohol and 76.3% (74/97) chewed BQ. The healthy subjects, including smokers without a BQ chewing history ($N = 107$), smokers with a BQ chewing history ($N = 111$), and nonsmokers with a BQ chewing history ($N = 12$), were matched for age and gender for the buccal cell and urine sample studies (Table 2). The number of years as a smoker with a BQ chewing history (27.1 years) was significantly greater than that of smokers without a BQ chewing history (23.1 years; $P < 0.01$).

Increased Acr-dG adducts of buccal DNA in patients with OSCC compared with healthy subjects with CS and/or BQ chewing histories

We examined buccal DNA samples collected from healthy subjects and OSCC subjects by slot blot analysis using anti-Acr-dG antibodies to analyze Acr-induced DNA adducts [α - and γ -hydroxy-1, N^2 -cyclic propano-2'-deoxyguanosine (α -OH-Acr-dG and γ -OH-Acr-dG) adducts] that have been described previously (refs. 30, 31; Supplementary Fig. S2; Fig. 1A–C). We found that the Acr-dG levels in the buccal DNA of patients with OSCC were 1.4-fold higher than those of healthy subjects with CS or BQ chewing histories ($P < 0.001$; Fig. 1D; Table 3); however, there was no significant difference in healthy subjects with different CS or BQ chewing histories (Supplementary Fig. S3). To analyze the CS exposure spectrum between OSCC and healthy subjects, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL), a urinary biomarker for assessing tobacco exposure, was used (refs. 36–38; Supplementary Fig. S4). We found that Acr-dG levels in buccal DNA of patients with OSCC with CS and BQ chewing histories were 1.3 higher than those of healthy subjects with CS and BQ chewing histories ($P < 0.05$; Fig. 1E), while there was no significant difference in NNAL/Cre between OSCC and healthy subjects (Supplementary Table S1). These results partially excluded the possibility of different CS exposure spectra between healthy subjects and patients with OSCC. Interestingly, we further found that Acr-dG levels in tumor tissue DNA were 1.8-fold higher than those from buccal DNA in the same group of patients with OSCC ($P < 0.01$; Supplementary Fig. S5A and S5B). Overall, we conclude that the levels of Acr-dG adducts in buccal DNA or tumor DNA of patients with OSCC were higher than those in healthy

subjects, suggesting a higher degree of Acr-induced DNA damage in patients with OSCC compared with normal subjects.

Increased urinary 3-HPMA/Cre in healthy subjects with CS and BQ chewing histories compared with healthy subjects with only CS histories

We analyzed the levels of urinary 3-HPMA, the major metabolite of Acr, using isotope dilution HPLC mass spectrometry that has been previously established (33) in urine samples of patients with OSCC and healthy subjects. The results show that 3-HPMA levels in healthy subjects with CS and BQ chewing histories (8.9 ± 9.3 $\mu\text{mole/g}$ creatinine) were significantly higher compared with those in healthy subjects with only CS histories (5.8 ± 4.7 $\mu\text{mole/g}$ creatinine; $P < 0.01$) or only BQ chewing histories (3.6 ± 6.1 $\mu\text{mole/g}$ creatinine; $P < 0.01$; Fig. 2A; Table 3). Furthermore, we found that in healthy subjects with CS and BQ chewing histories, 3-HPMA levels (9.2 ± 9.7 $\mu\text{mole/g}$ creatinine) were significantly higher than those in healthy subjects with only CS histories (6.0 ± 5.0 $\mu\text{mole/g}$ creatinine; $P < 0.01$) in matched smoking years (smoking years >15 years; Fig. 2B). The correlation between urinary 3-HPMA and smoking years of healthy subjects was studied. We found that an increase in urinary 3-HPMA/Cre was strongly correlated with an increase in the number of smoking years in healthy subjects with smoking histories without BQ chewing histories ($P = 0.0027$) or with a BQ chewing histories ($P = 0.0389$; Fig. 2C).


Decreased urinary 3-HPMA/Cre in patients with OSCC compared with healthy subjects

Interestingly, we found that urinary 3-HPMA levels in patients with OSCC (0.7 ± 0.7 $\mu\text{mole/g}$ creatinine) were significantly lower than those in healthy subjects with CS or BQ chewing histories (7.1 ± 7.4 $\mu\text{mole/g}$ creatinine; $P < 0.001$; Fig. 3A). A similar phenomenon was observed in patients with OSCC (0.6 ± 0.6 $\mu\text{mole/g}$ creatinine) with CS and BQ chewing histories and healthy subjects with CS and BQ chewing histories (9.0 ± 9.5 $\mu\text{mole/g}$ creatinine; $P < 0.001$; Fig. 3B), while there were no significant differences between NNAL/Cre levels in patients with OSCC and healthy subjects (Supplementary Table S1). 3-HPMA is a detoxifying metabolite of Acr when conjugated with GSH, and it has been shown that intracellular GSH levels are inversely correlated with Acr-induced cytotoxicity (39). Furthermore, we found that Acr decreased total intracellular GSH levels or GSH/GSSG levels in SAS cells (Fig. 3C). These results indicate that cellular GSH levels determine the detoxifying activity of Acr, and chronic exposure of Acr through CS may reduce cellular protection for Acr-induced DNA damage.

Discussion

The importance of the $p53$ tumor suppressor gene in the process of carcinogenesis is well established, and a high incidence of $p53$ mutations has been demonstrated in tobacco-related cancers (10, 40). Previous studies have shown that a high incidence (48.66%, 91/187) of $p53$ mutations in OSCC exhibited mutations in the conserved exons of the $p53$ gene, which is similar to reports from Western countries (16, 41). The most prevalent types of $p53$ mutations found in these studies were G to A transitions and G to T transversions. Acr induces mutagenic Acr-dG adducts and mainly G to T and G to A mutations in human cells (21–27). By mapping the Acr-dG adduct distribution at the nucleotide level in

Tsou et al.

**Figure 1.**

Slot blot analysis of acrolein-induced DNA (Acr-dG) adducts in buccal DNA of healthy subjects and patients with OSCC. **A**, The chemical reaction of Acr with deoxyguanosine (dG) residues of DNA to form α - and γ -hydroxy-1, N²-cyclic propano-2'-deoxyguanosine (α -OH-Acr-dG and γ -OH-Acr-dG) adducts. **B**, A typical slot blot result is shown. DNA was spotted on the membrane, hybridized with the Acr-dG antibody, then with the quantum dot-conjugated second antibody, and the binding of Acr-dG antibody was analyzed using fluorescence development. First, four bands of the first lane, human genomic DNA isolated from cultured OSCC cells (SAS), were modified with different concentrations of Acr (0–5 mmol/L) at 37°C for 24 hours as Acr-dG standards (Acr-dG STD); the second and third lanes include DNA from healthy subjects; and the fourth, fifth and sixth lanes include DNA from patients with OSCC. Left, equivalent amounts of DNA loaded in the membrane that was stained with methylene blue; right, fluorescence development and the band intensity was quantified with a UVP image analyzer. **C**, Relative Acr-dG levels were calculated by the fluorescence intensity of Acr-dG stained with an anti-Acr-dG antibody normalized to the amount of loaded DNA stained with methylene blue. The standard curve was determined by the fluorescence intensity of the relative Acr-dG DNA adduct level in genomic DNA modified with different concentrations of Acr (0, 0.5, 2, and 5 mmol/L) as Acr-dG adduct standards. Relative Acr-dG adduct levels in buccal cells of healthy subjects ($n = 222$) and OSCC patient buccal cells ($n = 80$; **D**) and relative Acr-dG adduct in healthy subjects with CS and BQ chewing histories ($n = 101$) and OSCC patients with comparable CS and BQ chewing histories ($N = 51$) as detected by the slot blot analysis described above (**E**). Bar graphs of data were collected from three independent slot blot experiments. Data were presented as the mean \pm SD. Student *t* tests were used to determine statistical significance, and two-tailed *P* values are shown (*, $P < 0.05$; ***, $P < 0.001$).

Acr-treated normal human bronchial epithelial cells, we found that the Acr–DNA binding spectrum in the *p53* gene coincides with the *p53* mutational spectrum in lung cancer (28). Interest-


ingly, there are some similar *p53* mutation hotspots and mutational spectra between CS-induced lung cancer and OSCC (Supplementary Fig. S1). On the basis of these findings, we hypothesized that Acr is a major etiologic agent in OSCC induced by CS and BQ chewing. In this study, we found that the levels Acr-dG adducts were significantly higher in patients with OSCC compared with those in healthy subjects with comparable CS and BQ chewing histories (Fig. 1D and E). These results suggest that increased Acr-dG levels are due to differences in Acr processing other than different CS exposure spectra between healthy subjects and patients with OSCC. In addition, we found substantially increased Acr-dG adduct levels in tumor tissue DNA compared with those in buccal DNA in the same population of patients with

Table 3. Urinary 3-HPMA levels in healthy subjects with CS or BQ chewing histories and patients with OSCC

	Creatinine Mean \pm SD (mg/dL)	3-HPMA Mean \pm SD (μ mol/L)	3-HPMA/Cr Mean \pm SD (μ mole/g)
CS ($n = 111$)	71.9 \pm 65.7	4.3 \pm 6.5	5.8 \pm 4.7
CS+BQ ($n = 107$)	167.4 \pm 117.9	16.1 \pm 23.2	8.9 \pm 9.3
BQ ($n = 12$)	196.3 \pm 154.8	9.0 \pm 15.7	3.6 \pm 6.1
Oral cancer (OSCC; $n = 97$)	96.8 \pm 67.8	0.6 \pm 0.7	0.7 \pm 0.7

Figure 2.

Levels of 3-HPMA/Cre in the urine of healthy subjects with CS or BQ chewing histories. 3-HPMA and Cre levels of healthy subjects were measured using isotope dilution HPLC mass spectrometry, as described in the Materials and Methods section. **A**, 3-HPMA and creatinine (Cre) levels of 230 healthy subjects with CS or BQ chewing histories included smokers without BQ chewing histories (S, $n = 111$), smokers with BQ chewing histories (S+B, $n = 107$), and nonsmokers with BQ chewing histories (B, $n = 12$). **B**, 3-HPMA and Cre levels of healthy subjects with matched CS histories (>15 smoking years) included smokers without BQ chewing histories (S, $N = 84$) and smokers with BQ chewing histories (S+B, $N = 95$). Bar graphs of data were presented as the mean \pm SD. Student *t* tests were used to determine statistical significance, and two-tailed *P* values are shown. **, $P < 0.01$. S indicates smokers without BQ chewing histories; S+B indicates smokers with BQ chewing histories; and B indicates nonsmokers with BQ chewing histories. **C**, The correlation between urinary 3-HPMA/Cre levels and smoking years of healthy subjects. Urinary 3-HPMA/Cre levels in 111 smokers without BQ chewing histories (top) and 107 smokers with BQ chewing histories (bottom) were evaluated by Pearson correlation analysis.




OSCC (Supplementary Fig. S5A and S5B). Furthermore, we also did sequencing analysis on exon 5–9 of *p53* gene. Results showed that in 12 of 26 samples, mutations were found on *p53* gene and

the major types of *p53* mutation were G to T and G to A mutation (66.7%, 8/12 samples; Supplementary Fig. S5C). Taken together, these results indicate that the accumulation of Acr-dG adducts

Figure 3.

Levels of 3-HPMA/Cre in the urine of healthy subjects and patients with OSCC. 3-HPMA/creatinine (Cre) levels of 230 healthy subjects with CS or BQ chewing histories and 97 patients with OSCC (**A**) and 3-HPMA/Cre levels in healthy subjects with CS and BQ chewing histories ($n = 101$) and OSCC patients with comparable CS and BQ chewing histories ($n = 51$) were measured (**B**) as described in Fig. 2. Bar graphs of data were presented as the mean \pm SD. Student *t* tests were used to determine statistical significance, and two-tailed *P* values are shown. ***, $P < 0.001$. **C**, Total GSH and GSH/GSSG levels in SAS cells treated with Acr (50 $\mu\text{mol/L}$, 3 hours) were analyzed using a glutathione fluorometric assay kit (Biovision Inc.). Bar graphs of data were collected from three independent experiments. Data were presented as the mean \pm SD. Student *t* tests were used to determine statistical significance, and two-tailed *P* values are shown (***, $P < 0.001$; *, $P < 0.05$).



Tsou et al.

may induce *p53* mutations and contribute to oral carcinogenesis. To the best of our knowledge, this is the first study to hypothesize Acr etiology in OSCC.

Urinary 3-HPMA, the primary metabolite of Acr found in urine (19), has been used as a biomarker for Acr exposure through cigarette smoking or cooking fumes (20). Our results showed that 3-HPMA levels were closely correlated with the number of smoking years of healthy subjects with smoking histories (Fig. 2C). However, the correlation is stronger in smokers without BQ chewing histories than smokers with BQ chewing histories. This result suggested that BQ chewing probably influences the metabolism of Acr inside the human body. In the matched smoking years (>15 smoking years), we found that 3-HPMA levels are much higher in smokers with BQ chewing histories than those in smokers without BQ chewing histories (Fig. 2B). BQ is mixed with the betel nut and its additives. Commonly mixed additives include the areca nut, unripened fruit of *piper betel* linn, betel leaf of *piper betel* linn, *piper betel* linn, slaked lime, and other spices. In 1987, the International Agency for Research on Cancer (IARC) stated that the use of BQ with tobacco is a carcinogen in humans (Group 1; ref. 42). The addition of slaked lime to the betel nut block neutralizes the astringency of a large amount of tannic acid and polyphenolic compounds in the betel nut, but it causes the oral cavity to be alkaline. Studies have shown that the pH of the oral cavity after chewing the betel nut can reach 8.2–11 (43, 44). The alkaline oral environment (pH 8–11) following the chewing of BQ-containing slake lime might facilitate the absorption of Acr (*pKa* 9.6) during CS (45). This finding provides a possible explanation for the synergistic effects of CS and BQ chewing in oral cancer.

In addition, we found that urinary 3-HPMA levels were significantly lower in patients with OSCC than in healthy subjects with comparable CS or BQ chewing histories (Fig. 3A and B; Table 3; Supplementary Table S1). Previous studies have shown decreased GSH levels in the blood of patients with oral cancer (0.69 ± 0.11 mmol/L) compared with those in healthy subjects (2.54 ± 0.33 mmol/L; ref. 46). The reduced level of GSH in these patients may have been the cause of reduced urine 3-HPMA levels in patients with oral cancer, reflecting their reduced capacity to systemically neutralize Acr toxicity. However, GSH levels in oral tissues directly affect Acr-induced DNA damage in oral tissues. Acr has been shown to reduce intracellular GSH rapidly and further induce oxidative stress (47, 48), which is similar to our results shown in Fig. 3C. In addition, Acr has been shown to reduce DNA repair capacity (21, 22, 39). These results suggest that chronic exposure to Acr may reduce cellular detoxifying activity and repair capacity against Acr. Furthermore, oral tissue GSH levels may change due to pathologic changes. Previous studies have shown significantly increased levels of GSH in tumor tissue compared with those in tumor-free tissue of patients with oral cancer ($P < 0.001$; refs. 46, 49). These results indicate that enhanced GSH levels or antioxidant capacities in tumor tissue can make them less susceptible to oxidative stress, conferring a selective growth advantage on tumor cells. Taken together, these results suggest that increased Acr-dG levels in buccal cells of patients with OSCC, coupled with the reduced repair and detoxifying capacity of Acr-induced damage, may contribute to oral carcinogenesis.

One limitation of our study is that we could not exclude other exposure possibilities to Acr other than CS. Acr is an important pollutant that is widely distributed in the environment because it is a prominent byproduct of the incomplete combustion of

organic matter and is also formed from carbohydrates, vegetable oils, animal fats, and amino acids during the heating of foods (18, 50). Therefore, humans are exposed to Acr by the inhalation of incompletely combusted organic matter and overheated cooking oil fumes or by the ingestion of many foods (51). However, smoking of tobacco products is equivalent to or exceeds the total human exposure to Acr from all other sources (52). Previous studies have suggested that a high level of oral exposure to Acr through these sources induces detrimental effects on the oral cavity, including salivary quality and contents, oral resistance to oxidative stress, and stress mechanism activation in a variety of oral cells (50). Therefore, protective pathways against Acr-induced cytotoxicity in the oral cavity need to be further investigated.

OSCC is one of the most common epithelial malignancies with significant morbidity and mortality (1–3). The key challenge to reduce the mortality and morbidity of this disease is to develop strategies to identify and detect OSCC when it is present at a very early stage, which will enable effective intervention and therapy. The detection of OSCC is currently based on expert clinical examinations and histologic analysis of suspicious areas (53, 54), but it may be undetectable in hidden sites. Therefore, sensitive and specific biomarkers for OSCC may be helpful in screening high-risk patients (9). These results provide a novel mechanism by which increased Acr uptake and impaired metabolism may contribute to OSCC carcinogenesis. Elevated Acr-dG levels in buccal swabs may serve as an early indicator to identify patients at risk.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: H.-T. Wang, H.-H. Tsou, C.-J. Liu, T.-Y. Liu
Development of methodology: H.-T. Wang, H.-H. Tsou, C.-H. Hu, C.-H. Lee
Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): H.-H. Tsou, C.-H. Hu, J.-H. Liu, C.-J. Liu, C.-H. Lee, T.-Y. Liu
Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): H.-T. Wang, H.-H. Tsou, C.-H. Hu, C.-H. Lee
Writing, review, and/or revision of the manuscript: H.-T. Wang, H.-H. Tsou, C.-J. Liu, C.-H. Lee
Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): H.-T. Wang, T.-Y. Liu
Study supervision: H.-T. Wang, T.-Y. Liu

Acknowledgments

We thank Dr. Moon-shong Tang, New York University, for providing monoclonal anti-Acr-dG antibody and Dr. Kuo-Wei Chang, Institute of Oral Biology, School of Dentistry, National Yang-Ming University (Taipei, Taiwan), for providing SAS cells. We thank Ms. Tse-Wen Wang and Ms. Pei-Ying Che, two graduate students of Institute of Food Safety and Health Risk Assessment, National Yang-Ming University (Taipei, Taiwan), for their great help with slot blot analysis and *p53* sequencing analysis. We also thank Ms. Chia-Lin Wang, Assistant Researcher of Taipei Veterans General Hospital Department of Medical Research, for her great help with LC/MS-MS analysis. This work was supported by Ministry of Science and Technology, Taiwan (grant no. 104-2320-B-010-040-MY3, to H.-T. Wang; 106-3114-B-010001, to H.-T. Wang; 107-2320-B-010-018, to H.-T. Wang).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received September 18, 2018; revised November 9, 2018; accepted February 26, 2019; published first March 6, 2019.

References

- Warnakulasuriya S. Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol* 2009;45:309–16.
- Petersen PE. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Community Dent Oral Epidemiol* 2003;31(Suppl 1):3–23.
- Gupta N, Gupta R, Acharya AK, Patthi B, Goud V, Reddy S, et al. Changing trends in oral cancer - a global scenario. *Nepal J Epidemiol* 2016;6:613–9.
- Ko YC, Huang YL, Lee CH, Chen MJ, Lin LM, Tsai CC. Betel quid chewing, cigarette smoking and alcohol consumption related to oral cancer in Taiwan. *J Oral Pathol Med* 1995;24:450–3.
- Kumar M, Nanavati R, Modi TG, Dobariya C. Oral cancer: etiology and risk factors: a review. *J Cancer Res Ther* 2016;12:458–63.
- Tsai KY, Su CC, Lin YY, Chung JA, Lian Ie B. Quantification of betel quid chewing and cigarette smoking in oral cancer patients. *Community Dent Oral Epidemiol* 2009;37:555–61.
- Wen CP, Tsai SP, Cheng TY, Chen CJ, Levy DT, Yang HJ, et al. Uncovering the relation between betel quid chewing and cigarette smoking in Taiwan. *Tob Control* 2005;14(Suppl 1):16–22.
- Ries LMD, Krapcho M. SEER cancer statistics review, 1975–2005. Bethesda, MD: NIH/NCI; 2008. Available from: https://seer.cancer.gov/archive/csr/1975_2005/.
- Saxena S, Sankhla B, Sundaragiri KS, Bhargava A. A review of salivary biomarker: a tool for early oral cancer diagnosis. *Adv Biomed Res* 2017; 6:90.
- Olivier M, Eeles R, Hollstein M, Khan MA, Harris CC, Hainaut P. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum Mutat* 2002;19:607–14.
- Rivlin N, Brosh R, Oren M, Rotter V. Mutations in the p53 tumor suppressor gene: important milestones at the various steps of tumorigenesis. *Genes Cancer* 2011;2:466–74.
- Staib F, Hussain SP, Hofseth LJ, Wang XW, Harris CC. TP53 and liver carcinogenesis. *Hum Mutat* 2003;21:201–16.
- Giglia-Mari G, Sarasin A. TP53 mutations in human skin cancers. *Hum Mutat* 2003;21:217–28.
- Hsieh LL, Wang PF, Chen IH, Liao CT, Wang HM, Chen MC, et al. Characteristics of mutations in the p53 gene in oral squamous cell carcinoma associated with betel quid chewing and cigarette smoking in Taiwanese. *Carcinogenesis* 2001;22:1497–503.
- Wong YK, Liu TY, Chang KW, Lin SC, Chao TW, Li PL, et al. p53 alterations in betel quid- and tobacco-associated oral squamous cell carcinomas from Taiwan. *J Oral Pathol Med* 1998;27:243–8.
- Blons H, Laurent-Puig P. TP53 and head and neck neoplasms. *Hum Mutat* 2003;21:252–7.
- Singh RD, Patel KR, Patel PS. p53 mutation spectrum and its role in prognosis of oral cancer patients: a study from Gujarat, West India. *Mutat Res* 2016;783:15–26.
- Moghe A, Ghare S, Lamoreau B, Mohammad M, Barve S, McClain C, et al. Molecular mechanisms of acrolein toxicity: relevance to human disease. *Toxicol Sci* 2015;143:242–55.
- Stevens JF, Maier CS. Acrolein: sources, metabolism, and biomolecular interactions relevant to human health and disease. *Mol Nutr Food Res* 2008;52:7–25.
- Carmella SG, Chen M, Zhang Y, Zhang S, Hatsukami DK, Hecht SS. Quantitation of acrolein-derived (3-hydroxypropyl)mercapturic acid in human urine by liquid chromatography-atmospheric pressure chemical ionization tandem mass spectrometry: effects of cigarette smoking. *Chem Res Toxicol* 2007;20:986–90.
- Wang HT. Effect of acrolein in DNA damage, DNA repair, and lung carcinogenesis. New York: New York University; 2012. p.1–190.
- Tang MS, Wang HT, Hu Y, Chen WS, Akao M, Feng Z, et al. Acrolein induced DNA damage, mutagenicity and effect on DNA repair. *Mol Nutr Food Res* 2011;55:1291–300.
- Wang HT, Zhang S, Hu Y, Tang MS. Mutagenicity and sequence specificity of acrolein-DNA adducts. *Chem Res Toxicol* 2009;22:511–7.
- Yang IY, Miller H, Wang Z, Frank EG, Ohmori H, Hanaoka F, et al. Mammalian translesion DNA synthesis across an acrolein-derived deoxyguanosine adduct. Participation of DNA polymerase ϵ in error-prone synthesis in human cells. *J Biol Chem* 2003;278:13989–94.
- Sanchez AM, Minko IG, Kurtz AJ, Kanuri M, Moriya M, Lloyd RS. Comparative evaluation of the bioreactivity and mutagenic spectra of acrolein-derived alpha-HOPdG and gamma-HOPdG regioisomeric deoxyguanosine adducts. *Chem Res Toxicol* 2003;16:1019–28.
- Minko IG, Washington MT, Kanuri M, Prakash L, Prakash S, Lloyd RS. Translesion synthesis past acrolein-derived DNA adduct, gamma-hydroxypropanodeoxyguanosine, by yeast and human DNA polymerase ϵ . *J Biol Chem* 2003;278:784–90.
- Kanuri M, Minko IG, Nechev LV, Harris TM, Harris CM, Lloyd RS. Error prone translesion synthesis past gamma-hydroxypropano deoxyguanosine, the primary acrolein-derived adduct in mammalian cells. *J Biol Chem* 2002;277:18257–65.
- Feng Z, Hu W, Hu Y, Tang MS. Acrolein is a major cigarette-related lung cancer agent: preferential binding at p53 mutational hotspots and inhibition of DNA repair. *Proc Natl Acad Sci U S A* 2006;103: 15404–9.
- Association of Directors of Anatomical and Surgical Pathology. Recommendations for the reporting of specimens containing oral cavity and oropharynx neoplasms. *Mod Pathol* 2000;13:1038–41.
- Wang HT, Weng MW, Chen WC, Yobin M, Pan J, Chung FL, et al. Effect of CpG methylation at different sequence context on acrolein- and BPDE-DNA binding and mutagenesis. *Carcinogenesis* 2013;34:220–7.
- Lee HW, Wang HT, Weng MW, Hu Y, Chen WS, Chou D, et al. Acrolein- and 4-Aminobiphenyl-DNA adducts in human bladder mucosa and tumor tissue and their mutagenicity in human urothelial cells. *Oncotarget* 2014;5: 3526–40.
- Pan J, Awoyemi B, Xuan Z, Vohra P, Wang HT, Dyba M, et al. Detection of acrolein-derived cyclic DNA adducts in human cells by monoclonal antibodies. *Chem Res Toxicol* 2012;25:2788–95.
- Tsou HH, Hsu WC, Fuh JL, Chen SP, Liu TY, Wang HT. Alterations in acrolein metabolism contribute to Alzheimer's disease. *J Alzheimers Dis* 2018;61:571–80.
- Shah KA, Halquist MS, Karnes HT. A modified method for the determination of tobacco specific nitrosamine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in human urine by solid phase extraction using a molecularly imprinted polymer and liquid chromatography tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 2009;877: 1575–82.
- Peng SY, Tu HF, Yang CC, Wu CH, Liu CJ, Chang KW, et al. miR-134 targets PDCD7 to reduce E-cadherin expression and enhance oral cancer progression. *Int J Cancer* 2018;143:2892–904.
- Hecht SS. Tobacco carcinogens, their biomarkers and tobacco-induced cancer. *Nat Rev Cancer* 2003;3:733–44.
- Goniewicz ML, Eisner MD, Lazcano-Ponce E, Zielinska-Danch W, Koszowski B, Sobczak A, et al. Comparison of urine cotinine and the tobacco-specific nitrosamine metabolite 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) and their ratio to discriminate active from passive smoking. *Nicotine Tob Res* 2011;13:202–8.
- Hecht SS. Carcinogen derived biomarkers: applications in studies of human exposure to secondhand tobacco smoke. *Tob Control* 2004; 13(Suppl 1):i48–56.
- Wang HT, Hu Y, Tong D, Huang J, Gu L, Wu XR, et al. Effect of carcinogenic acrolein on DNA repair and mutagenic susceptibility. *J Biol Chem* 2012; 287:12379–86.
- Greenblatt MS, Bennett WP, Hollstein M, Harris CC. Mutations in the p53 tumor suppressor gene: clues to cancer etiology and molecular pathogenesis. *Cancer Res* 1994;54:4855–78.
- Ostwald C, Gogacz P, Hillmann T, Schweder J, Gundlach K, Kundt G, et al. p53 mutational spectra are different between squamous-cell carcinomas of the lip and the oral cavity. *Int J Cancer* 2000;88:82–6.
- Overall evaluations of carcinogenicity: an updating of IARC monographs volumes 1 to 42. *IARC Monogr Eval Carcinog Risks Hum Suppl* 1987;7: 1–440.
- Nair U, Bartsch H, Nair J. Alert for an epidemic of oral cancer due to use of the betel quid substitutes gutkha and pan masala: a review of agents and causative mechanisms. *Mutagenesis* 2004;19:251–62.
- Donoghue M, Basandi PS, Adarsh H, Madhushankari GS, Selvamani M, Nayak P. Habit-associated salivary pH changes in oral submucous fibrosis—a controlled cross-sectional study. *J Oral Maxillofac Pathol* 2015;19: 175–81.

Tsou et al.

45. Hayon E, Simic M. Acid-base properties of free radicals in solution. *Acc Chem Res* 1974;7:114–21.
46. Fiaschi AI, Cozzolino A, Ruggiero G, Giorgi G. Glutathione, ascorbic acid and antioxidant enzymes in the tumor tissue and blood of patients with oral squamous cell carcinoma. *Eur Rev Med Pharmacol Sci* 2005;9:361–7.
47. Colombo G, Dalle-Donne I, Orioli M, Giustarini D, Rossi R, Clerici M, et al. Oxidative damage in human gingival fibroblasts exposed to cigarette smoke. *Free Radic Biol Med* 2012;52:1584–96.
48. Avezov K, Reznick AZ, Aizenbud D. Oxidative damage in keratinocytes exposed to cigarette smoke and aldehydes. *Toxicol In Vitro* 2014;28:485–91.
49. Srivastava KC, Austin RD, Shrivastava D. Evaluation of oxidant-antioxidant status in tissue samples in oral cancer: a case control study. *Dent Res J* 2016;13:181–7.
50. Aizenbud D, Aizenbud I, Reznick AZ, Avezov K. Acrolein-an alpha,beta-unsaturated aldehyde: a review of oral cavity exposure and oral pathology effects. *Rambam Maimonides Med J* 2016;7:e0024.
51. Abraham K, Andres S, Palavinskas R, Berg K, Appel KE, Lampen A. Toxicology and risk assessment of acrolein in food. *Mol Nutr Food Res* 2011;55:1277–90.
52. Gerberding JL. Toxicological profile for Acrolein. Atlanta, GA: Agency for Toxic Substances and Disease Registry; 2007. Available from: <https://www.atsdr.cdc.gov/ToxProfiles/tp.asp?id=557&tid=102>.
53. Rethman MP, Carpenter W, Cohen EE, Epstein J, Evans CA, Flaitz CM, et al. Evidence-based clinical recommendations regarding screening for oral squamous cell carcinomas. *Tex Dent J* 2012;129:491–507.
54. Rethman MP, Carpenter W, Cohen EE, Epstein J, Evans CA, Flaitz CM, et al. Evidence-based clinical recommendations regarding screening for oral squamous cell carcinomas. *J Am Dent Assoc* 2010;141:509–20.

Cancer Epidemiology, Biomarkers & Prevention

Acrolein Is Involved in the Synergistic Potential of Cigarette Smoking— and Betel Quid Chewing-Related Human Oral Cancer

Han-Hsing Tsou, Chih-Hung Hu, Jin-Hui Liu, et al.

Cancer Epidemiol Biomarkers Prev 2019;28:954-962. Published OnlineFirst March 6, 2019.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-18-1033](https://doi.org/10.1158/1055-9965.EPI-18-1033)

Supplementary Material Access the most recent supplemental material at:
<http://cebp.aacrjournals.org/content/suppl/2019/03/06/1055-9965.EPI-18-1033.DC1>

Cited articles This article cites 51 articles, 9 of which you can access for free at:
<http://cebp.aacrjournals.org/content/28/5/954.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/28/5/954>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.