

Iron Intake and the Risk of Colorectal Cancer^{1,2}

John I. Wurzelmann,³ Andrew Silver,
Dina M. Schreinemachers, Robert S. Sandler, and
Richard B. Everson

Division of Digestive Diseases, Center for Gastrointestinal Biology and Disease, General Clinical Research Center, University of North Carolina, Chapel Hill, North Carolina 27599-7038 [J. I. W., A. S., R. S. S.], and U.S. Environmental Protection Agency, National Health and Environmental Effects Research Laboratory, Human Studies Division (MD-58C), Research Triangle Park, North Carolina 27711 [D. M. S., R. B. E.]

Abstract

Prospectively gathered data from the National Health and Nutrition Examination Survey I and the National Health Evaluation Follow-Up Study were analyzed to evaluate the risk of colorectal cancer due to consumption of iron. Morbidity and mortality data due to colorectal cancer were available on 14,407 persons first interviewed in 1971 and followed through 1986. A total of 194 possible colorectal cancers occurred in this group over the 15-year period. Subsite analysis showed that the risk of colon cancer due to iron intake was elevated throughout the colon for both men and women, with the highest adjusted risks for the interquartile range seen in the proximal colon for females (relative risk, 1.51; 95% confidence interval, 1.41-1.60). The risk of rectal cancer was not significantly elevated for men or women. Elevated serum iron was also associated with increased risk; however, this effect was strongest in the distal (rather than proximal) colon and was significant only among females (adjusted relative risk, 1.73; 95% confidence interval, 1.03-2.92). The mean transferrin saturation was higher among cases than controls (30.7 versus 28.7%), but total iron-binding capacity did not seem to predict the occurrence of colorectal cancer. Proportional hazards models confirmed that the effects of iron and serum iron were not confounded by age, gender, energy consumption, fat intake, or other known risk factors for colorectal cancer. These data suggest that iron may confer an increased risk for colorectal cancer, and that the localization of risk may be attributable to the mode of epithelial exposure. It seems that luminal exposure to

iron increases risk proximally, whereas humoral exposure increases risk distally. These differences may be due to such factors as oxidation state, binding proteins and the presence of other cofactors such as bile acids, products of bacterial metabolism.

Introduction

Iron is a necessary nutrient that has been associated with increased cancer risk in humans. *In vitro* and *in vivo* data suggest that iron may be capable of mutagenic effects mediated through free-radical generation or tumor promotion through nutritional mechanisms (1-3). As a transition metal, iron is possessed of loosely bound electrons that are capable of participating in lipid peroxidation reactions. Such reactions are thought to lead to DNA damage and, in some cases, neoplasia (4-7).

An earlier interval analysis based upon NHANES⁴ I (8) suggested that increased iron stores were associated with increased colorectal cancer risk, but this finding was based on a total of only 39 cases. The magnitude of risk was not specified, and there were too few cases to conduct a site- and gender-specific analysis. We sought to extend these observations by analyses of additional cases and years of follow-up data collected through 1987.

Materials and Methods

NHANES I aimed to determine the baseline health and nutritional status of the noninstitutionalized U.S. population. A total of 14,407 individuals, 25-74 years of age, were assessed between 1972 and 1974. Follow-up was obtained from 1982-1984, at which time 93% of the original cohort was traced, and again in 1986, when subjects older than 55 at intake were traced. The 1987 NHEFS was conducted for all members of the original cohort known not to be deceased as of 1986 ($n = 11,750$). Ultimately, 11,018 subjects (93.8%) were traced, and 9,998 subjects were interviewed (9). Measurements included a 24-h dietary intake and a food frequency questionnaire. Iron and other nutrient intakes were calculated by conversion of reported intakes using nutrient information from the U.S. Department of Agriculture Handbook No. 8. Supplemental iron was not considered. Biochemical measurements, including serum iron and total iron-binding capacity, were measured spectrophotometrically with the use of a modification of the automated Technicon AAI method (10). Detailed information on the design and conduct of the NHEFS has been published elsewhere (11-13).

A total of 196 possible colorectal cancers were identified in the cohort of 14,407. Two cases were excluded because they were subsequently identified as anal cancers. The diagnoses

Received 10/19/95; revised 4/4/96; accepted 4/4/96.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported in part by grants from the NIH (P30 DK34987, T32 DK07364, and RR00046).

² Although the research described in this article has been supported by the United States Environmental Protection Agency, it has not been subjected to Agency review and therefore does not necessarily reflect the views of the Agency and no official endorsement should be inferred. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

³ To whom requests for reprints should be addressed, at Division of Digestive Diseases, CB # 7038, Room 156, Glaxo Building, University of North Carolina, Chapel Hill, NC 27599-7038. Phone: (919) 966-2516.

⁴ The abbreviations used are: NHANES, National Health and Nutrition Examination Survey; NHEFS, NHANES Epidemiologic Follow-Up Study; CI, confidence interval; RR, relative risk.

were confirmed by examination of hospital and nursing home records, including pathology reports. When a subject was found to be deceased, a death certificate was obtained from the appropriate state vital statistics office (14). To eliminate confounding by occult disease or the inclusion of individuals not at risk for colon cancer, statistical analysis was limited to individuals not censored within the first 3 years of follow-up, and to individuals more than or equal to 31 years old, the age of the youngest case at intake. Nutritive intake was adjusted for caloric content by the regression method of Willett *et al.* (15). The crude risk was calculated by comparing cancer incidence in the second, third, and fourth quartiles of exposure to cancer incidence in the first quartile. This was done to demonstrate the presence and magnitude of the effect in the raw data. Ninety-five percent CIs were calculated according to the method described in Kleinbaum *et al.* (16). Cutoffs for quartiles were taken from univariate analyses of the initial data set. The significance of trends was calculated using the Mantel-Haenszel test (17). The means were compared by two-tailed *t* test (18).

The adjusted RR was calculated for the interquartile range of exposure, using proportional hazards models adjusting for age and gender (19). The location was coded by the rules of the International Classification of Diseases, Ninth Revision (20). The following codes were included in the analysis as colon cancers: 153.0, 153.1, 153.2, 153.4, 153.6, 153.7, 153.8, and 153.9; whereas 154.0 and 154.1 were classified as rectal cancers. The location of cancers was classified as: (a) either rectal or colon; and (b) either proximal or distal. Proximal cancers were considered to be those up to and including the splenic flexure. Distal colon cancers were distal to the splenic flexure (including the rectum). This was defined as such because it is presently understood that proximal colon cancers may have a different biology than distal colon or rectal cancers. It is not clear whether the risk changes gradually as one moves distally (suggesting the difference is due to an exposure gradient) or the difference starts abruptly at the rectosigmoid junction (suggesting that biological susceptibility is the reason for these differences). To simply split cancers into colon and rectal cancers is to assume the latter model, and possibly miss effects that may be present if the first model holds. Rather than assume one model, we have elected to show the effects based upon either assumption. Of note, a subset of colon cancers that did not have their sites specified (International Classification of Diseases codes 153.8 and 153.9; *n* = 39) could not be included in the analysis of proximal *versus* distal cancers. Risk was calculated for iron intake, serum iron, total iron-binding capacity, and transferrin saturation. Transferrin saturation was obtained by dividing serum iron by the total iron-binding capacity. The independence of the main effects was assayed by the inclusion of these variables in proportional hazards models with known risk factors for colorectal cancer (including age, body-mass index, fat and energy consumption, aspirin intake, vegetable consumption, physical activity, and education). Potential interactions were assayed by evaluating the RR due to iron at different levels of these same factors.

Results

The data set of 14,407 included all persons ages 25–74 who completed a medical examination at the time of the original NHANES I sample (1971–1975). After eliminating persons less than 31 years old and those censored within the first 3 years of follow-up, the final data set used as a basis for regression and stratified analyses was comprised of 11,317 individuals. A total

Table 1 Demographic and clinical comparisons of the initial and final data sets, NHANES I and NHEFS, 1982–1987

	Initial data set	Final data set
White	83.5%	84.1%
Black	15.3%	14.8%
Female	59.7%	59.1%
Age	50.0%	52.8%
College education	20.6%	20.7%
Colon cancers	<i>n</i> = 140	<i>n</i> = 118
Rectal cancers	<i>n</i> = 54	<i>n</i> = 38

Table 2 Mean iron indices of cases compared with unaffected subjects NHANES I and NHEFS, 1982–1987

	Females		Males	
	Controls	Cases	Controls	Cases
Iron intake (mg) (SE)	9.6 (.07)	9.4 (.58)	13.5 (.11)	14.4 (.87)
Serum iron (mcg/dl) (SE)	96.7 (.51)	104.5 (4.9)	106.1 (.64)	111.0 (4.8)
TIBC (mcg/dl) (SE)	365 (.86)	372 (6.5)	348 (.87)	343 (6.51)
Transferrin Sat. (%) (SE)	27.1 (0.2)	28.5 (1.3)	31.0 (0.2)	33.0 (1.6)

of 156 individuals who developed colorectal cancer were included. There were 118 colon cancers and 38 rectal cancers. Within this data set, iron intake was available for 8876 persons (including 136 cases), and serum iron was available for 8345 persons (including 126 cases). Whites and women comprised 81.8% and 60.1% of the total sample, respectively. The initial and final data sets were similar in composition, as shown in Table 1. In the final data set, cases were significantly older than unaffected subjects: 62.2 years compared to 53.3 years, respectively (*P* < 0.0001). Cases also had significantly higher serum iron (107.7 *versus* 100.4 mcg/dl; *P* = 0.03) and transferrin saturation (30.7 *versus* 28.7%; *P* = 0.05) by two-tailed *t* test. Breakdown of a comparison by gender is presented in Table 2.

The crude risk of cancer due to iron intake by site and gender for the entire study population is shown in Table 3. Risk was highest for women in the proximal colon (RR, 7.76; 95% CI, 1.36–44.3) and was also elevated in the colon in general. A similar (but less pronounced) pattern is seen in males. The crude risk attributable to serum iron is shown in Table 4. Unlike iron intake, the risk attributable to serum iron increases as one progresses distally, with the peak risk seen in the rectum, again in women (RR, 9.77; 95% CI, 1.81–52.8). Risk is not significantly elevated for males distally. Significant effects were not seen for either total iron-binding capacity or transferrin saturation.

Proportional hazards models were run to adjust for possible confounders and interactions. After adjustment for age and gender, iron intake is associated with increased risk in the proximal colon (RR, 1.44; 95% CI, 1.23–1.69), but not in the distal colon (RR, 1.03; 95% CI, 0.80–1.32). Serum iron predicts rectal cancer (RR, 1.57; 95% CI, 1.08–2.30) and, to a lesser extent, colon cancer (RR, 1.24; 95% CI, 0.96–1.59). When these analyses were stratified by gender, the risk for rectal cancer was again higher for women for serum iron (RR, 1.73; 95% CI, 1.03–2.92) and for iron intake in the proximal colon (RR, 1.51, 95% CI, 1.41–1.60). Models including variables for age, sex, fat, calories, body mass index, physical activity, vegetable and meat intake, and recent aspirin consumption did not show these variables to be confounders, insofar as they did not lead to appreciable changes in risk due to iron.

Table 3 Crude risk due to iron intake by location and gender (RR, 95% CI), NHANES I and NHEFS, 1982-1987

	Proximal	Cases	Distal	Cases	Colon	Cases	Rectum	Cases
All								
1st Q		5		11		10		8
2nd Q	2.62 (0.95-7.32)	14	1.45 (0.69-3.08)	17	2.24 (1.10-4.59)	24	1.29 (0.63-3.15)	11
3rd Q	2.26 (0.82-6.23)	12	1.12 (0.50-2.49)	13	2.63 (1.32-5.25)	28	1.18 (0.47-2.98)	10
4th Q	3.15 (1.41-7.04)	21	1.47 (0.73-2.97)	16	3.35 (1.74-6.46)	36	1.01 (0.42-2.42)	9
Trend	$P = 0.007$		$P = 0.61$		$P < 0.001$		$P = 0.98$	
Males								
1st Q		4		5		6		3
2nd Q	1.97 (0.57-6.80)	6	2.10 (0.71-6.24)	8	1.75 (0.62-11.0)	8	2.63 (0.70-9.95)	6
3rd Q	1.39 (0.35-5.51)	4	1.67 (0.52-5.37)	6	2.30 (0.87-6.13)	10	2.31 (0.58-9.26)	5
4th Q	3.08 (1.06-8.97)	12	2.06 (0.72-5.88)	10	3.73 (1.62-8.59)	22	1.72 (0.42-7.17)	5
Trend	$P = 0.06$		$P = 0.27$		$P = 0.001$		$P = 0.58$	
Females								
1st Q		1		6		4		5
2nd Q	5.89 (0.95-36.6)	8	1.11 (0.40-3.10)	9	2.94 (1.04-8.31)	16	0.74 (0.22-2.54)	5
3rd Q	5.78 (0.93-36.1)	8	0.85 (0.29-4.76)	7	3.24 (1.17-8.97)	18	0.73 (0.21-2.51)	5
4th Q	7.76 (1.36-44.3)	9	0.87 (0.28-2.68)	6	3.01 (1.05-8.64)	14	0.70 (0.21-2.86)	5
Trend	$P = 0.05$		$P = 0.67$		$P = 0.08$		$P = 0.22$	

Table 4 Crude risk due to serum iron by location and gender (RR, 95% CI), NHANES I and NHEFS, 1982-1987

	Proximal	Cases	Distal	Cases	Colon	Cases	Rectum	Cases
All								
1st Q		13		8		23		3
2nd Q	0.82 (0.37-1.80)	11	1.93 (0.84-4.43)	16	0.84 (0.47-1.58)	20	3.86 (1.20-12.5)	12
3rd Q	1.23 (0.60-2.55)	16	2.24 (1.00-5.04)	18	1.05 (0.59-1.85)	24	4.32 (1.37-13.6)	13
4th Q	0.63 (0.24-1.65)	6	2.04 (0.85-4.89)	12	1.36 (0.77-2.41)	23	3.63 (1.05-12.5)	8
Trend	$P = 0.70$		$P = 0.10$		$P = 0.24$		$P = 0.06$	
Males								
1st Q		6		4		8		7
2nd Q	0.76 (0.25-2.37)	6	1.52 (0.46-4.98)	8	1.05 (0.42-2.59)	11	2.28 (0.48-10.8)	6
3rd Q	1.06 (0.38-2.96)	9	1.59 (0.50-5.07)	9	1.32 (0.57-3.09)	15	1.76 (0.35-8.87)	5
4th Q	0.44 (0.11-1.67)	3	1.20 (0.37-4.56)	6	1.08 (0.43-2.72)	10	1.73 (0.32-9.2)	4
Trend	$P = 0.40$		$P = 0.73$		$P = 0.73$		$P = 0.73$	
Females								
1st Q		7		4		15		1
2nd Q	0.79 (0.25-2.47)	5	2.19 (0.68-7.05)	8	0.66 (0.29-1.51)	9	6.58 (1.05-41.1)	6
3rd Q	1.23 (0.43-3.49)	7	2.75 (0.89-8.49)	9	0.74 (0.33-1.68)	9	9.77 (1.81-52.8)	8
4th Q	0.79 (0.21-3.03)	3	2.74 (0.82-9.20)	6	1.58 (0.76-3.29)	13	7.31 (1.13-47.2)	4
Trend	$P = 0.99$		$P = 0.09$		$P = 0.32$		$P = 0.04$	

Discussion

Stevens *et al.* (10) first suggested an association between iron stores and cancer risk in 1988, based upon NHANES I data. This report was based on a total of 242 cancers, including 22 colon and rectal cancers (21). An in-depth analysis of individual cancer types was not possible, given the small numbers involved. A subsequent report was based on a total of 39 colorectal cancers (8). We have undertaken a detailed analysis of risk factors for colon and rectal cancer, expecting that the additional data gained since the last report would bring more statistical power to the analysis.

Meat intake has generally been associated with increased risk of colorectal cancer (15, 22-32), but the specific mechanisms of carcinogenesis have been debated. Risk has been attributed to fat intake, but fat itself is not mutagenic and is better characterized as a tumor promoter. It has been postulated that mutagens are produced through the effects of bacterial enzymes on fat or bile acids (33, 34), or by the production of polycyclic aromatic hydrocarbons in the course of protein pyrolysis (35). On the other hand, iron is also consumed when meat is ingested and may induce damage and/or increased

proliferation through lipid peroxidation (4, 5, 36). An effect may result merely from iron's status as a nutrient necessary for proliferation. In this regard, it has been argued that the anemia often seen with malignancy is an adaptation intended to slow tumor growth (37). Several studies have shown augmentation of tumor growth in rodent models with iron supplementation (2, 3, 37-39). Siegers *et al.* (40) have shown this effect to be dose dependent. In one study (38), supplementation with phytate (a chelator of iron found in high-fiber diets) abolished this effect, thus creating a theoretical basis for the protective effect of dietary fiber. Conversely, iron deficiency seems to inhibit the growth of tumor cells (1). Recently, Fukuchi *et al.* (41) have shown that iron deprivation increases p53 expression in two cell lines, suggesting that the carcinogenic effects of iron may be mediated by alteration of normal mechanisms for the maintenance of genomic stability. In this regard, the iron chelator has been shown to be a potent inhibitor of DNA-synthesis and cell proliferation (42, 43).

There is also evidence that iron may cause cancer in humans. Individuals with hereditary hemochromatosis are at increased risk for hepatocellular carcinoma (44, 45). Bradbear

et al. (46) note a nonsignificant excess of colorectal cancers in genetic hemochromatosis patients. Colorectal adenomas and gastric cancers have been associated with increased serum ferritin (47, 48).

The data we present supports the hypothesis that iron intake increases the risk of proximal cancers. This effect seems to be more pronounced in women. A variety of influences are thought to determine the location of cancers. McMichael and Potter (49) first noted an increased propensity for proximal cancer in women and attributed this to the effect of estrogen on the delivery of bile acids to the colon. Other factors that have been associated with proximal cancers include genotype and advanced age (50). Cancers due to hereditary mismatch repair defects are thought to favor the right side of the colon (51). These cancers would typically occur in younger individuals. However, there was no difference between the ages of cases reporting proximal cancers and those reporting distal cancers.

There are numerous physiological differences between the right and left colon that may explain this site-specific variation in risk. The left colon is more densely colonized with bacteria, and the proportion of anaerobic bacteria increases as one moves distally. The number and species of bacteria probably influence the production and metabolism of potential carcinogens.

Serum iron effects, on the other hand, are significant only in the distal bowel and rectum. This exposure differs from that of ingested iron in that it is relatively constant, controlled by endogenous binding proteins, not exposed to the actions or debris of bacterial metabolism, and not subjected to the high reducing potential of the colon's luminal milieu. In this data set, there is a weak correlation ($r = 0.04$; $P = 0.0001$) between serum iron and iron intake. This is because serum levels are also influenced by gastrointestinal losses, extraintestinal bleeding, efficiency of absorption, and an equilibrium with extravascular stores.

This study has the advantages of prospective design and large sample size, including a representative cross-section of the U.S. population. On the other hand, the measurement of dietary intake was relatively crude, and it may be argued that the effects attributed to iron were due to some other constituent of meat. However, meat intake was not a risk factor or a confounder, and the adjustment for fat intake did not alter the risk due to iron. In summary, these data suggest that iron may be an important determinant of colorectal cancer risk. This effect may be modified by several genetic, hormonal, and physiological factors, which in turn may impact the location of disease.

References

- Hann, H. W., Stahlhut, M. W., and Blumberg, B. S. Iron nutrition and tumor growth: decreased tumor growth in iron-deficient mice. *Cancer Res.*, 48: 4168–4170, 1988.
- Singh, M., Lu, J., Briggs, S. P., McGinley, J. N., Haegele, A. D., and Thompson, H. J. Effect of excess dietary iron on the promotion stage of 1-methyl-1-nitrosourea-induced mammary carcinogenesis: pathogenetic characteristics and distribution of iron. *Carcinogenesis (Lond.)*, 15: 1567–1570, 1994.
- Thompson, H. J., Kennedy, K., Witt, M., and Juzefyk, J. Effect of dietary deficiency or excess on the induction of mammary carcinogenesis by 1-methyl-1-nitrosourea. *Carcinogenesis (Lond.)*, 12: 111–114, 1991.
- Minotti, G., and Aust, S. D. The role of iron in the initiation of lipid peroxidation. *Chem. Phys. Lipids*, 44: 191–208, 1987.
- Sevanian, A., and Hochstein, P. Mechanisms and consequences of lipid peroxidation in biological systems. *Annu. Rev. Nutr.*, 5: 365–390, 1985.
- de Kok, T. M. C. M., ten Vaarwek, F., Zwingman, I., van Maanen, J. M. S., and Kleinjans, J. C. S. Peroxidation of linoleic, arachidonic, and oleic acid in relation to the induction of oxidative DNA damage and cytogenetic effects. *Carcinogenesis (Lond.)*, 15: 1399–1404, 1994.
- Baader, S. L., Bruchelt, G., Carmine, T. C., Lode, H. N., Rieth, A. G., and Niethammer, D. Ascorbic acid-mediated iron release from cellular ferritin and its relation to the formation of DNA strand breaks in neuroblastoma cells. *J. Cancer Res. Clin. Oncol.*, 120: 415–421, 1994.
- Stevens, R. G., Graubard, B. I., Micozzi, M. S., Neriishi, K., and Blumberg, B. S. Moderate elevation of body iron level and increased risk of cancer occurrence and death. *Int. J. Cancer*, 56: 364–369, 1994.
- U. S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Center for Health Statistics. Public Use Data Tape Documentation: NHANES I Epidemiologic Follow-Up Study, 1987, pp. 3–6. Hyattsville, MD: U.S. Government Printing Office, 1991.
- Stevens, R. G., Jones, D. Y., Micozzi, M. S., and Taylor, P. R. Body iron stores and the risk of cancer. *N. Engl. J. Med.*, 319: 1047–1052, 1988.
- Plan and operation of the NHANES I Epidemiologic Follow-Up Study, 1987. *Vital Health Stat. [1]*, 27: , 1991.
- National Center for Health Statistics. Plan and operation of the National Health and Nutrition Examination Survey. *Vital Health Stat. [1]*, 10: , 1977.
- Cohen, B. B., Barbano, H. E., Cox, C. S., *et al.* Plan and operation of the NHANES I Epidemiologic Follow-Up Study, 1982–1984. *Vital Health Stat. [1]*, 22: 1–4, 1987.
- Finucane, F. F. Public Use Data Tape Documentation: NHANES I Epidemiologic Follow-Up Study, 1986: Vital and Tracing Status. Hyattsville, MD: U. S. Dept. of Health and Human Services, Public Health Services, Centers for Disease Control, and National Center for Health Statistics, 1990.
- Willett, W. C., Stampfer, M. J., Colditz, G. A., Rosner, B. A., and Speizer, F. E. Relation of meat, fat, and fiber intake to the risk of colon cancer in a prospective study among women. *N. Engl. J. Med.*, 323: 1664–1672, 1990.
- Kleinbaum, D. G., Kupper, L. L., and Morgenstern, H. *Epidemiologic Research, Principles, and Quantitative Methods*, pp. 298–299. New York: Van Nostrand Reinhold, 1982.
- Mantel, N. [Chi]-square tests with one degree of freedom: extensions of the Mantel-Haenszel procedure. *J. Am. Stat. Assoc.*, 58: 690–700, 1959.
- Gossett, W. S. The probable error of the mean. *Biometrika*, 6: 1–25, 1908.
- Cox, D. R., and Oakes, D. *Analysis of Survival Data*. London: Chapman and Hall, 1984.
- Manual of the International Statistical Classification of Diseases, Injuries, and Causes of Death: Based on the Recommendations of the Ninth Revision Conference, 1975, and Adopted by the Twenty-ninth Health Assembly, Vol. 1. Geneva: World Health Organization, 1977.
- Molbak, K., Aaby, P., Hojlyng, N., and da Silva, A. P. J. Risk factors for cryptosporidium diarrhea in early childhood: a case-control study from Guinea-Bissau. *Am. J. Epidemiol.*, 139: 734–740, 1994.
- Armstrong, B., and Doll, R. Environmental factors and cancer incidence and mortality in different countries with special reference to dietary practices. *Int. J. Cancer*, 15: 617–631, 1975.
- Jensen, O. M., and MacLennan, R. Dietary factor and colorectal cancer in Scandinavia. *Isr. J. Med. Sci.*, 15: 329–334, 1979.
- McKeown-Eyssen, G. E., and Bright-See, E. Dietary factors in colon cancer: international relationships. *Nutrition and Cancer*, 1992.
- Rose, D. P., Boyar, A. P., and Wynder, E. L. International comparisons of mortality rates for cancer of the breast, ovary, prostate, and colon, and per capita food consumption. *Cancer (Phila.)*, 58: 2363–2371, 1986.
- Kato, I., Tominaga, S., and Kuroishi, T. Per capita foods/nutrients intake and mortality from gastrointestinal cancers in Japan. *Jpn. J. Cancer Res.*, 78: 453–459, 1987.
- Haenszel, W., Berg, J. W., Segi, M., Kurihara, M., and Locke, F. B. Large-bowel cancer in Hawaiian Japanese. *J. Natl. Cancer Inst.*, 51: 1765–1778, 1973.
- Phillips, R. L. Role of life-style and dietary habits in risk of cancer among Seventh-Day Adventists. *Cancer Res.*, 35: 3513–3522, 1975.
- Gerhardsson de Verdier, M., Hagman, U., Peters, R. K., Steineck, G., and Overvik, E. Meat, cooking methods, and colorectal cancer: a case-referent study in Stockholm. *Int. J. Cancer*, 49: 520–525, 1991.
- Miller, A. B., Howe, G. R., Jain, M., Craib, K. J. P., and Harrison, L. Food items and food groups as risk factors in a case-control study of diet and colorectal cancer. *Int. J. Cancer*, 32: 155–161, 1983.
- Tajima, K., and Tominaga, S. Dietary habits and gastrointestinal cancers: a comparative case-control study of stomach and large intestinal cancers in Nagoya, Japan. *Jpn. J. Cancer Res.*, 76: 705–716, 1985.
- Lee, H. P., Gourley, L., Duffy, S. W., Estève, J., Lee, J., and Day, N. E. Colorectal cancer and diet in an Asian population: a case-control study among Singapore Chinese. *Int. J. Cancer*, 43: 1007–1016, 1989.
- Wakabayashi, K., Nagao, M., Esumi, H., and Sugimura, T. Food-derived mutagens and carcinogens. *Cancer Res. (Suppl.)*, 52: 2092s–2098s, 1992.

34. Hill, M. J., Drasar, B. S., Aries, V., Crowther, J. S., Hawksworth, G., and Williams, R. E. O. Bacteria and the aetiology of cancer of the large bowel. *Lancet*, *1*: 95–100, 1971.
35. Prival, M. J. Carcinogens and mutagens present as natural components of food or induced by cooking. *Nutr. Cancer*, *6*: 236–253, 1985.
36. Halliwell, B. Oxidants and human disease: some new concepts. *FASEB J.*, *1*: 358–364, 1987.
37. Bergeron, R. J., Streiff, R. R., and Elliott, G. T. Influence of iron on *in vivo* proliferation and lethality of L1210 cells. *J. Nutr.*, *115*: 369–374, 1985.
38. Nelson, R. L., Yoo, S. J., Tanure, J. C., Andrianopoulos, G., and Misumi, A. The effect of iron on experimental colorectal carcinogenesis. *Anticancer Res.*, *9*: 1477–1482, 1989.
39. Siegers, C. P., Bumann, D., Baretton, G., and Younes, M. Dietary iron enhances the tumor rate in dimethylhydrazine-induced colon carcinogenesis in mice. *Cancer Lett.*, *41*: 251–256, 1988.
40. Siegers, C. P., Bumann, D., Trepkau, H. D., Schadwinkel, B., and Baretton, G. Influence of dietary iron overload on cell proliferation and intestinal tumorigenesis in mice. *Cancer Lett.*, *65*: 245–249, 1992.
41. Fukuchi, K., Tomoyasu, S., Watanabe, H., Kaetsu, S., Tsuruoka, N., and Gomi, K. Iron deprivation results in an increase in p53 expression. *Biol. Chem. Hoppe-Seyler*, *376*: 627–630, 1995.
42. Robbins, E., and Pederson, T. Iron, its intracellular localization and possible role in cell division. *Proc. Natl. Acad. Sci. USA*, *66*: 1244–1251, 1970.
43. Blatt, J., and Stitley, S. Antineuroblastoma activity of desferrioxamine in human cell lines. *Cancer Res.*, *47*: 1749–1750, 1987.
44. Niederau, C., Strohmeyer, G., and Stremmel, W. Epidemiology, clinical spectrum and prognosis of hemochromatosis. *Adv. Exp. Med. Biol.*, *356*: 293–302, 1994.
45. Niederau, C., Fischer, R., Sonnenberg, A., Stremmel, W., Trampisch, H. J., and Strohmeyer, G. Survival and causes of death in cirrhotic and in noncirrhotic patients with primary hemochromatosis. *N. Engl. J. Med.*, *313*: 1256–1262, 1985.
46. Bradbear, R. A., Bain, C., Siskind, V., *et al.* Cohort study of internal malignancy in genetic hemochromatosis and other chronic nonalcoholic liver diseases. *J. Nat. Cancer Inst.*, *75*: 81–84, 1985.
47. Nelson, R. L., Davis, F. G., Sutter, E., Sobin, L. H., Kikendall, J. W., and Bowen, P. Body iron stores and the risk of colonic neoplasia. *J. Nat. Cancer Inst.*, *86*: 455–460, 1994.
48. Nomura, A., Chyou, P. H., and Stemmermann, G. N. Association of serum ferritin levels with the risk of stomach cancer. *Cancer Epidemiol., Biomarkers & Prev.*, *1*: 547–550, 1992.
49. McMichael, A. J., and Potter, J. D. Do intrinsic differences in lower alimentary tract physiology influence the sex-specific risks of bowel cancer and other biliary and intestinal diseases. *Am. J. Epidemiol.*, *118*: 620–627, 1983.
50. Schub, R., and Steinheber, F. U. Rightward shift of colon cancer: a feature of the aging gut. *J. Clin. Gastroenterol.*, *8*: 630–634, 1986.
51. Thibodeau, S. N., Bren, G., and Schaid, D. Microsatellite instability in cancer of the proximal colon. *Science (Washington DC)*, *260*: 816–889, 1993.

Cancer Epidemiology, Biomarkers & Prevention

AACR American Association
for Cancer Research

Iron intake and the risk of colorectal cancer.

J I Wurzelmann, A Silver, D M Schreinemachers, et al.

Cancer Epidemiol Biomarkers Prev 1996;5:503-507.

Updated version Access the most recent version of this article at:
<http://cebp.aacrjournals.org/content/5/7/503>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link <http://cebp.aacrjournals.org/content/5/7/503>. Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.