

Missing the Target for Routine Human Papillomavirus Vaccination: Consistent and Strong Physician Recommendations Are Lacking for 11 to 12-Year-Old Males

Susan T. Vadaparampil^{1,2,3}, Teri L. Malo⁴, Steven K. Sutton^{1,3}, Karla N. Ali¹, Jessica A. Kahn⁵, Alix Casler⁶, Daniel Salmon⁷, Barbara Walkosz⁸, Richard G. Roetzheim^{1,9}, Gregory D. Zimet¹⁰, and Anna R. Giuliano^{1,2,3}

Abstract

Background: Rates of routine human papillomavirus (HPV) vaccination of adolescent males in the United States are low. Leading health organizations advocate consistent and strong physician recommendations to improve HPV vaccine dissemination. This study describes the prevalence and correlates of consistent and strong physician recommendations for HPV vaccination of adolescent males.

Methods: We surveyed pediatric and family medicine physicians in Florida about their HPV vaccine recommendations for male vaccine-eligible age groups (11–12, 13–17, 18–21 years). Descriptive statistics compared consistency and strength of HPV recommendations across age groups. Multivariable logistic regression examined factors associated with consistent and strong recommendations for 11- to 12-year-olds.

Results: We received 367 completed surveys (51% response rate). Physicians most often consistently and strongly recommended HPV vaccine to males ages 13 to 17 (39%) compared

with ages 11 to 12 (31%) and 18 to 21 (31%). Consistent and strong recommendation for 11- to 12-year-old males was more likely to be delivered by Vaccine for Children providers and less likely among physicians who reported more personal barriers to vaccination, particularly concerns about vaccine safety, concerns about adding vaccines to the vaccine schedule, and difficulty in remembering to discuss HPV vaccination.

Conclusion: Physicians' current consistency and strength of HPV vaccine recommendations do not align with national recommendations. Interventions to improve HPV vaccine recommendations must also consider the influence of physicians' personal barriers to HPV vaccine delivery.

Impact: As one of the first studies to examine both consistency and strength of physicians' HPV vaccine recommendations for males, our findings can inform future interventions focused on facilitating physicians' recommendations. *Cancer Epidemiol Biomarkers Prev*; 25(10); 1–12. ©2016 AACR.

Introduction

In 2011, the Advisory Committee on Immunization Practices (ACIP) recommended routine human papillomavirus (HPV) vaccination of all males ages 11 to 12 years, catch-up vaccination

for males ages 13 to 21 years, and vaccination for men who have sex with men ages 22 to 26 (1). The ACIP updated this recommendation in 2015 to include the recently licensed 9-valent HPV vaccine (2). Yet HPV vaccination uptake among adolescent males nationally and in Florida is modest. In 2014, HPV vaccine initiation rate among males both in Florida and nationally was approximately 41%; coverage with ≥ 3 dose among males ages 13 to 17 was 22% in the United States and 17.5% in Florida (range: 9% in Alabama to 43% in Rhode Island; ref. 3). Available data suggest rates are even lower for 11- to 12-year-olds, who constitute the target age group for vaccination (4). Florida has among the nation's highest rates of HPV-related diseases in males, including the fifth highest rate of anal cancer (5). Given the established efficacy of preventing HPV infections that cause anal and penile cancer (6, 7), HPV vaccination has tremendous potential for primary prevention (8, 9), particularly for males from racial/ethnic and sexual minority groups disproportionately affected by HPV-related diseases (10–12).

The Centers for Disease Control and Prevention (CDC; ref. 13) and the President's Cancer Panel (14) advocate consistent and strong physician recommendations, particularly for the target age group of 11- to 12-year-old adolescents, as a primary approach to improving HPV vaccine dissemination. Yet relatively little

¹Division of Population Science, Moffitt Cancer Center, Tampa, Florida. ²Center for Infection Research in Cancer, Moffitt Cancer Center, Tampa, Florida. ³Department of Oncologic Sciences, College of Medicine, University of South Florida, Tampa, Florida. ⁴Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, North Carolina. ⁵Department of Pediatrics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio. ⁶Department of Pediatrics, Orlando Health Physician Associates, Orlando, Florida. ⁷Department of International Health and Health Behavior Society, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland. ⁸Klein Buechel, Boulder, Colorado. ⁹Department of Family Medicine, College of Medicine, University of South Florida, Tampa, Florida. ¹⁰Department of Pediatrics, Indiana University School of Medicine, Indianapolis, Indiana.

Corresponding Author: Susan T. Vadaparampil, Moffitt Cancer Center, 12902 Magnolia Drive MRC-CANCONT, Tampa, FL 33612. Phone: 813-745-1997; Fax: (813) 449-8020; E-mail: Susan.Vadaparampil@moffitt.org

doi: 10.1158/1055-9965.EPI-15-1294

©2016 American Association for Cancer Research.

research has examined the consistency and strength of physicians' HPV vaccine recommendations specific to males following the ACIP's recommendation for routine vaccination. Regarding consistency, results from a statewide survey of pediatricians, family physicians, and nurse practitioners in Minnesota showed that about 46% of providers routinely recommended HPV vaccine for boys ages 11 to 12 years (15). A national survey found that 61% of pediatricians and family physicians started routinely recommending HPV vaccine for males at or before age 12. For strength, a national survey of pediatricians and family physicians found that 52% of pediatricians and 41% of family physicians strongly recommend HPV vaccine for 11- to 12-year-old males (16), compared with 64% of pediatricians and 50% of family physicians in Hawaii (17). To our knowledge, only one study has examined both consistency and strength of physicians' HPV vaccine recommendations as part of a larger recommendation quality measure but did not focus solely on male patients (16). Because available literature demonstrates a disparity in recommendation practices such that fewer physicians report consistent and strong HPV vaccine recommendations for males compared with females (15–18), it is important to further explore factors associated with physicians' HPV vaccine recommendations for males.

Understanding recommendation practices and factors associated with those practices in a state with low vaccine uptake is essential to designing and implementing interventions to improve physicians' HPV vaccine recommendations. The medical encounter presents competing demands that pull physicians in many directions and influence how they recommend/provide preventive services during patient encounters. These ideas serve as the basis for the Competing Demands Model (CDM; ref. 19), originally developed to understand delivery of clinical preventive services in the primary care setting. The CDM proposes that physician (e.g., specialty, attitudes), patient (e.g., knowledge, attitudes), and practice environment (e.g., setting, location) factors influence physicians' delivery of preventive health services. Although several of these CDM factors have been explored by our group (20–22) and others (23), few have delved into vaccine-specific factors within the practice environment, such as the presence of a vaccine coordinator and use of specific types of reminder systems for series initiation and completion.

This study examines the prevalence of physician recommendation of HPV vaccination to vaccine-eligible males ages 11 to 12, 13 to 17, and 18 to 21 in Florida. Given ACIP recommendations emphasize routine vaccination targeting 11- to 12-year-olds and an emphasis on multilevel approaches to improving HPV vaccination rates (24), we assessed examined physician-reported factors relevant to the physician and practice (both general and vaccine specific) domains of the CDM as correlates of HPV vaccine recommendation for this age group.

Materials and Methods

Recruitment

Physicians were recruited from the American Medical Association (AMA) Physician Masterfile, a database of all licensed U.S. physicians (25). Our initial sampling frame excluded those who (i) were trainees; (ii) locum tenens; (iii) reported their major professional activity as nonpatient care; (iv) were ≥ 65 years of age, as the AMA Masterfile has been shown to have a significant lag in updating retired physicians (26); and (v) listed a post office box

for their address (precluding use of FedEx mailing). Florida pediatric and family medicine (FM) physicians were sampled on the basis of their proportional representation in the Florida physician primary care workforce and randomly selected from the AMA Masterfile ($n = 770$). We selected only 1 physician per group practice. Informed by Dillman tailored design method (27), prenotice postcards were mailed in May 2014, followed by the first survey mailing in June 2014. Although the survey itself was anonymous, we limited subsequent mailings by using a postage paid postcard that was included with every survey mailing where physicians were asked to provide their name, address, and select from options including: (i) I have completed and sent the survey back in the prepaid envelope; or (ii) I do not wish to participate and have returned the blank survey and the \$25 incentive. Follow-up mailings were sent to all those from whom we did not receive a completed postcard through August 2014. Participating physicians received a \$25 cash incentive.

Of the 770 surveys mailed, 367 were received. After accounting for undeliverable surveys ($n = 36$) and ineligible respondents ($n = 10$), the overall response rate was 51% (367/724). We excluded 12 participants who reported not seeing male patients ages 9 to 26, for an analytic sample size of 355.

Instrument

Where possible, we used previous survey items to assess HPV vaccination recommendation (20, 28–33); new items that were created to measure constructs not assessed in previous studies are noted below. The final 49-item survey assessed three domains relevant to the CDM: physician characteristics, physician-reported general and vaccine-specific practice characteristics.

Physician-reported characteristics were perceived personal and parental barriers related to HPV vaccination, HPV-related knowledge, and demographic characteristics. Sixteen items assessed physician-reported barriers to immunizing male patients against HPV. Perceived parental HPV vaccination barriers for 9- to 17-year-old male patients were measured using 14 items. Response options for physician-reported and perceived parental barriers were on a 4-point Likert scale (1 = not a barrier at all to 4 = a major barrier). Items were summed to create scores for perceived personal (range: 16–64; Cronbach $\alpha = 0.88$) and parental barriers (range: 14–56; Cronbach $\alpha = 0.90$). Lower scores indicated lower perceived barriers. Knowledge was measured using 9 items regarding HPV infection, disease, and vaccine guidelines for males. We built on previous surveys by creating new items to assess knowledge specific to guidelines and financial coverage for vaccinating males. One point was awarded for each correct response, and correct responses were summed to create a knowledge score (range: 0–9).

Physician-reported general practice characteristics included number of physicians in the practice, practice situation (single specialty, multispecialty, other), practice type (private, other), race/ethnic category of the majority of patients seen, whether the practice serves Medicaid patients, typical daily patient load, and practice location (urban, suburban, rural, other). Vaccine-specific practice characteristics were administration of HPV vaccine, Vaccines for Children (VFC) provider, strategies for remembering to discuss HPV vaccine with male patients, specific strategies to get patients into the office for the first and subsequent dose(s) of vaccine, total number of strategies used to get patients into the office for the first and subsequent dose(s) of vaccine, and presence of an office vaccine coordinator. We created new survey items to

assess whether other health care professionals (i.e., medical assistant, nurse, nurse practitioner, physician assistant) in the practice discuss and recommend HPV vaccine. The final survey can be obtained by emailing the corresponding author.

In addition to items evaluating domains relevant to CDM, our survey included new and previous items that assessed the context and content of vaccination recommendation (34). New items also assessed provider's acceptance and use of the CDC messages and materials to support HPV vaccine recommendation for adolescent males (35) and intervention preferences.

The primary outcome variables were HPV vaccine recommendations to vaccine-eligible males ages 11 to 12, 13 to 17, and 18 to 21. Given that national guidelines call for consistent and strong recommendations, we assessed recommendation consistency and strength using two questions from prior studies (33). The first question asked providers to indicate how often they recommended HPV vaccination by age group: never/almost never (<10%), occasionally (10%–39%), about half the time (40%–59%), usually (60%–90%), and always/almost always (>90%). Providers were also asked how strongly they recommended vaccination by age group: I recommend against; I make no recommendation for or against; I recommend, but not strongly; and I strongly recommend. In addition to reviewing these items individually, we created a composite variable to reflect that national guidelines call for both consistent and strong recommendations. We combined these two questions into one variable to compare those who reported recommending vaccination consistently (always/almost always) and strongly to physicians who reported any other combination of responses.

Statistical analysis

Frequencies and percentages were calculated for independent and outcome variables. To reflect national guidelines for consistent and strong HPV vaccine recommendations targeted to adolescent males ages 11 to 12, we focused analyses on assessing correlates of recommendation for this age group. This was done in three steps. First, simple logistic regression models examined each correlate. Second, within each domain, significant univariate correlates were entered into a multivariable model. A backward elimination approach (significance level of stay = 0.05) was used to determine those correlates, making relatively independent contributions to consistent and strong HPV vaccine recommendations for each domain. Significant correlates in the final model for each domain were then entered into a multivariable model using a backward elimination approach to generate a model of the relatively independent correlates across domains. ORs and their 95% confidence intervals (CI) were estimated in each model. All analyses used two-tailed tests with the significance level set at $P < 0.05$ and were performed using the SAS 9.3 statistical software package (SAS Institute Inc.).

Results

The sample was almost equally comprised of female (51.0%) and male (49.0%) physicians (Table 1), with an average age of 48.7 years (SD = 9.0). Most were white (67.7%) and non-Hispanic (75.1%). About half specialized in FM (49.9%) and had been practicing for ≥ 16 years (51.9%). HPV vaccine knowledge scores averaged 5.7 (SD = 2.1; range: 0–9).

Physicians most often reported their practice had two physicians (49.9%), was single specialty (66.8%), private practice

(67.2%), and in a suburban location (52.6%). More than one third (36.6%) reported seeing mostly patients from minority groups and about one fifth (20.2%) reported no definable majority. The majority reported seeing Medicaid patients either along with other types of insurance (67.6%) or solely (1.9%). Almost half (44.5%) saw 20 to 29 patients daily. The majority reported administering HPV vaccine in their practice (68.8%) and nearly half were VFC providers (45.9%). More than half did not use a strategy to get patients in for the first dose of HPV vaccine (52.5%), but most used ≥ 1 strategies to get patients vaccinated with subsequent doses (84.3%). Physicians reported using flagging charts (16.6%), automatic prompts (22.2%), and electronic queries (13.6%) to remind themselves to discuss HPV vaccine. About three fourths reported having a vaccine coordinator (72.2%).

The anonymous nature of the survey precludes examining the degree to which survey responders and nonresponders were similar on demographic and practice characteristics. However, we were able to compare responding physicians to the population of physicians in Florida, meeting our study eligibility criteria on characteristics, including age, sex, and clinical specialty. We found no statistically significant difference between responding physicians and the larger population of Florida physicians for age, sex, and clinical specialty (all $P > 0.05$).

Compared with older adolescent groups, a lower proportion of physicians reported consistently (34.8%), strongly (42.9%), and both consistently and strongly (30.6%) recommending HPV vaccine to the 11- to 12-year-old group (Fig. 1). As shown in Figs. 2 and 3, physicians' mean perceived personal barriers score was 30.3 (SD = 9.6; range: 16–61), and the mean perceived parental barriers score was 37.0 (SD = 8.9; range: 14–56). As can be seen in Table 2, consistent and strong HPV vaccine recommendation for 11 to 12 year olds was associated with the following variables in univariate analyses: physician gender, specialty, personal barriers, HPV knowledge, patient race, Medicaid patients seen, practice location, and all vaccine-specific practice characteristics.

Within each domain (physician characteristics, general practice characteristics, and vaccine-specific practice characteristics), a multivariable analysis using a backward stepwise approach was applied, starting with the significant univariate correlates in each group (Table 2). Two physician characteristics were significantly associated with consistent and strong recommendations: pediatric specialty [adjusted OR (AOR) = 2.55; 95% CI, 1.38–4.71] and fewer (or lower) physician-reported barriers (AOR = 0.94; 95% CI, 0.90–0.97). The lone general practice characteristic was serving Medicaid patients (OR = 3.11; 95% CI, 1.48–6.53). The lone vaccine-specific characteristic was being a VFC provider (OR = 6.48; 95% CI, 3.47–12.1).

A multivariable analysis was performed using backward stepwise regression, starting with the significant correlates in the domain-specific multivariable models (Table 2). In the final multivariable model, fewer (or lower) physician barriers (AOR = 0.91; 95% CI, 0.88–0.94) and being a VFC provider (AOR = 5.43; 95% CI, 2.80–10.55) were statistically significant correlates of HPV vaccine recommendation in 11- to 12-year-olds.

Given that the total score for physician barriers was inversely associated with strong and consistent HPV vaccine recommendations, we performed a *post hoc* analysis to explore each of the 16 physician barriers as a correlate. The results of these

Vadaparampil et al.

Table 1. Sample characteristics (N = 355)

Physician characteristics	n (%)	Mean (SD)
Gender		
Female	178 (51.0)	
Male	171 (49.0)	
Age		48.7 (9.0)
30–39	64 (18.3)	
40–49	124 (35.5)	
≥50	161 (46.1)	
Race		
White/Caucasian	233 (67.7)	
Black/African American	22 (6.4)	
Asian	38 (11.1)	
Other	51 (14.8)	
Ethnicity		
Hispanic	86 (24.9)	
Non-Hispanic	259 (75.1)	
Years practicing		
10 or fewer	100 (29.3)	
11–15	64 (18.8)	
16 or more	177 (51.9)	
Clinical specialty		
Pediatrics	155 (44.4)	
Family Medicine	174 (49.9)	
Other ^a	20 (5.7)	
HPV knowledge (range 0–9)		5.7 (2.1)
Physician barriers score (range 16–61) ^b		30.3 (9.6)
Parental barriers score (range 14–56) ^b		37.0 (8.9)
General practice characteristics	n (%)	Mean (SD)
Number of physicians in practice		
1	100 (28.7)	
2	174 (49.9)	
6–15	46 (13.2)	
16 or more	29 (8.3)	
Practice situation		
Single specialty	233 (66.8)	
Multispecialty	91 (26.1)	
Other	25 (7.2)	
Practice type		
Private practice office	229 (67.2)	
Other	112 (32.8)	
Practice location		
Urban	129 (37.7)	
Suburban	180 (52.6)	
Rural/other	33 (9.7)	
Race of majority of patients seen		
White, non-Hispanic	143 (43.2)	
Minority group(s)	121 (36.6)	
No definable majority ^c	67 (20.2)	
Medicaid patients served		
Medicaid, only	6 (1.9)	
Medicaid and others	215 (67.6)	
No Medicaid	97 (30.5)	
Typical daily patient load		
Less than 15	41 (11.8)	
15–19	96 (27.6)	
20–29	155 (44.5)	
30 or more	56 (16.1)	
Vaccine-specific characteristics	n (%)	Mean (SD)
Administer HPV vaccine		
Yes	242 (68.8)	
No	110 (31.2)	
VFC provider		
Yes	162 (45.9)	
No	156 (44.2)	
Do not know	35 (9.9)	
Strategies used for remembering to discuss HPV vaccine with eligible patients		
Flag charts	55 (15.5)	
Use automated electronic medical record prompts	74 (20.9)	
Perform periodic electronic queries to identify vaccine-eligible patients	45 (12.7)	
Used any of the 3 strategies	111 (31.3)	

(Continued on the following page)

Table 1. Sample characteristics (*N* = 355) (Cont'd)

Strategies to get patients into office for first dose of HPV vaccine	
Send reminder regarding preventive visit	75 (21.3)
Send letter or call patients specifically for HPV vaccine	37 (10.4)
Place reminder flag/tag in patients' medical record	31 (8.7)
Use a computerized immunization database/registry to track when first dose is due	47 (13.2)
Use some other strategy	18 (5.1)
Do not use any strategy	156 (43.9)
Not applicable: Do not administer vaccine	60 (16.9)
Strategies to get patients into office for second and third dose of HPV vaccine	
Record dose due date on card kept by patient	75 (21.3)
Send letter or call patients specifically for HPV vaccine	62 (17.5)
Place reminder flag/tag in patients' medical record	43 (12.1)
Schedule patient for next dose during office visit	189 (53.2)
Use a computerized immunization database/registry to track when first dose is due	49 (13.8)
Use some other strategy	20 (5.6)
Do not use any strategy	43 (12.1)
Not applicable: Do not administer vaccine	62 (17.5)
Number of strategies used for first dose of vaccine ^d	
None	140 (50.0)
1	89 (31.8)
2 or more	51 (18.2)
Number of strategies used for second and third doses of vaccine ^e	
None	38 (13.3)
1	134 (47.0)
2 or more	113 (39.7)
Other HCP in practice recommends HPV vaccine	
Yes	153 (43.7)
No	197 (56.3)
Other HCP in practice discusses HPV vaccine	
Yes	148 (42.4)
No	201 (57.6)
Vaccination coordinator in practice ^f	
Yes	252 (72.2)
No	88 (25.2)
Do not know	9 (2.6)

Abbreviation: HCP, health care provider.

^aOther clinical specialty includes urgent care, acute care, internal medicine, hospice, geriatrics, general physician, primary care, emergency room.

^bResponse options for physician-reported and perceived parental barriers were on a 4-point Likert scale (1 = not a barrier at all to 4 = a major barrier). Items were summed to create scores for perceived personal (range: 16–64) and parental barriers (range: 14–56). Lower scores indicated lower perceived barriers.

^cThere was no racial/ethnic group that comprised the majority of their patients.

^dStrategies used to get patients into the office for the first dose of HPV vaccine for those who administer vaccine.

^eStrategies used to get patients into the office for the second and third doses of HPV vaccine for those who administer vaccine.

^fQuestion was asked as follows: "Is there a vaccine coordinator in your office (i.e., someone responsible for purchasing, receiving, and storing vaccine shipments, maintaining vaccine inventory, training staff members on vaccine administration, etc.)?"

analyses are presented in Table 3. Alpha was adjusted to 0.0031 (0.05/16) for univariate analyses and $\alpha = 0.05$ for multivariable analyses. Ten of the 16 individual barriers were significant correlates (left column). Only one of these 10 barriers was significant in a multivariable model (middle column), highlighting the positive correlation among the individual barriers which is also exhibited in the relative high Cronbach α (0.88). The barriers that remained statistically significant following a backward stepwise approach (right column) were physician concern about vaccine safety, concern about adding another vaccine to the schedule, and remembering to discuss the vaccine.

Discussion

Recent efforts to improve physician recommendation for HPV vaccination have focused on two critical components. First, the recommendation should be consistent for all adolescent males and females, particularly the 11- to 12-year-old age group, for which routine vaccination is recommended. This is clearly demonstrated by the 2011 (1) and 2015 (2) ACIP

guidelines endorsed by all professional medical societies that provide preventive care to adolescent males. Second, as noted in the President's Cancer Panel Report focused on HPV vaccination, the recommendation must be strong (24). Our study demonstrates that, despite available guidelines, a minority of physicians indicated that they consistently (35%), strongly (43%), and both consistently and strongly (31%) recommend HPV vaccination to the target 11- to 12-year-old adolescent male age group. Although our study focused on recommendation for males, studies that have examined differences in recommendation by gender highlight a marked disparity in vaccine recommendations between boys and girls (15–18). Without immediate and targeted intervention, the nation is unlikely to achieve Healthy People 2020 goal of 80% of 13- to 15-year-old adolescents receiving the entire 3-dose HPV vaccine series (36).

Relative to physicians reporting high HPV vaccination barriers, those reporting low barriers were more likely to consistently and strongly recommend vaccination. This finding aligns with previous research reporting that barriers to vaccination were associated




Figure 1.

Percentage of physicians reporting consistent and strong recommendation of HPV vaccination ($n = 355$). The number of patients seen differs by patient age group and category: 11 to 12 years, $n = 284$ to 305; 13 to 17 years, $n = 303$ to 318; 18 to 21 years, $n = 312$ to 321.

with physicians not strongly recommending HPV vaccination to females ages 11 to 12 years (23) or offering the vaccine at all (37). Thus, one approach to improving vaccination recommendation may be to reduce provider barriers.

Barriers significantly associated with recommendation in the current study included concerns about vaccine safety, adding another vaccine to the schedule, and remembering to discuss vaccination. Interestingly, a systematic review of provider communication about HPV vaccination identified no studies demonstrating an association between providers' perceptions of HPV vaccine safety and their recommendation behaviors (38). However, our survey took place more recently than those included in the systematic review and may reflect the cumulative impact of media coverage that presents HPV vaccination in a controversial political context and as a vaccine for females (39). Thus, providers responding to our survey may be less aware of the safety of HPV vaccine for males. The ACIP recommendations for universal male HPV vaccination also coincided with political debates surrounding the vaccine during the 2012 presidential campaign (40). In addition, physician concerns about safety may be influenced by events such as the Japanese government's decision in June 2013 to stop proactive recommendation of HPV vaccine due to safety concerns (41, 42).

Physicians in our study also expressed concerns about adding another vaccine to the schedule. This concern has been documented in the context of adding a new vaccine to the early childhood immunization schedule (43) and HPV vaccination (20, 23). This finding is particularly concerning and should be addressed when

suggesting that providers deliver a bundled recommendation that includes HPV alongside tetanus, diphtheria, and acellular pertussis (Tdap) and meningococcal vaccines, which are also recommended at ages 11 to 12 for boys and girls (44).

Difficulty in remembering to discuss HPV vaccine with males was the third barrier associated with recommendation for males. Given the strong correlation between provider recommendation and HPV vaccine uptake in males (45), providers forgetting to discuss the vaccine offers one possible explanation for lower series initiation rates in males compared with females, despite the 2011 ACIP recommendation for routine vaccination in age-eligible males (1). The President's Cancer Panel recommends use of provider reminders to reduce missed opportunities for HPV vaccination (24). Almost two thirds of respondents (67%) did not use any reminders to discuss HPV vaccination with patients and approximately half (44%) used no strategy to get patients into the office for the first dose. In our study, we specifically examined the manual and electronic strategies for remembering to discuss HPV vaccine with male patients and specific strategies to get patients into the office for the first and subsequent dose(s) of vaccine. Although our study found no association between use of reminders and recommendation, it is possible that the relatively small number of those using these approaches precluded detection of a statistically significant association.

One study found that clinics using an electronic health record-based point-of-care reminder system targeting both clinicians and patients resulted in significantly more young women initiating and completing the HPV vaccine series in a timely fashion. The





Figure 2.

Physicians' barriers when immunizing male patients against HPV. STIs, sexually transmitted infections. *N* values differ for each barrier due to missing data. Missing data for each item ranged from 8 to 23 participants.

authors suggested that this combined strategy may have supported the creation of a common agenda that facilitates HPV vaccine series initiation (46). A recent literature review of interventions to increase HPV vaccination rates suggested that there was sufficient evidence to support the use of Community Preventive Task Force recommendations related to client reminder and recall systems as well as provider assessment and feedback

(47). Thus, future interventions may benefit from considering point-of-care reminders that simultaneously prompt physicians and patients to discuss HPV vaccination (48).

In our study, physicians who participated in the VFC program were more likely to recommend vaccination than those not participating. VFC-enrolled providers administer to Medicaid-eligible, uninsured, or American Indian or Alaska Native children

**Figure 3.**

Physicians' perceptions of parental barriers when immunizing male patients against HPV. STIs, sexually transmitted infections. Number of respondents ranged from 261 to 263. Many nonresponders see only patients 18 years or older.

younger than age 19 (49, 50). VFC also serves underinsured children in limited public settings (e.g., federally qualified health centers). Davis and colleagues (51) found that physician recommendation for pneumococcal conjugate vaccine (PCV7) was associated with the child's insurance coverage for the vaccine and

state VFC policy. Another study demonstrated that patients of VFC providers were 59% more likely to be up-to-date with early-childhood vaccines than patients of non-VFC providers (52). However, it is unclear why these providers are more likely to recommend vaccination. It is possible that participating in the

Table 2. Predictors of consistent and strong recommendation to 11- to 12-year-olds for physicians at practices that administer HPV vaccine (*N* = 208)

	OR (95% CI) ^a	Multivariable AOR (95% CI) ^b	Multivariable AOR (95% CI) ^c
Physician characteristics			
Female gender	1.45 (0.82–2.57)	—	—
Age	0.99 (0.96–1.02)	—	—
White, non-Hispanic race/ethnicity	1.00 (0.80–1.27)	—	—
Years practicing	0.99 (0.96–1.03)	—	—
Pediatric clinical specialty	2.54 (1.36–4.75)	0.74 (0.21–2.67)	—
Physician barriers score	0.92 (0.89–0.96)	0.94 (0.89–1.00)	0.93 (0.89–0.97)
Perceived parental barriers score	0.96 (0.93–1.00)	0.98 (0.93–1.03)	—
HPV knowledge	1.24 (1.06–1.45)	1.08 (0.88–1.34)	—
Practice characteristics: General			
Number of physicians in practice	0.89 (0.64–1.24)	—	—
Single-specialty practice	0.54 (0.29–0.99)	0.45 (0.18–1.13)	—
Private practice office	0.49 (0.26–0.90)	0.69 (0.25–1.89)	0.45 (0.22–0.92)
Majority of patients non-Hispanic White	0.42 (0.23–0.78)	0.69 (0.31–1.57)	—
Medicaid patients seen	2.89 (1.32–6.28)	0.58 (0.17–2.34)	—
Typical number of patients/day	1.01 (0.73–1.40)	—	—
Urban practice location	2.08 (1.16–3.71)	2.57 (1.17–5.64)	2.08 (1.08–4.02)
Practice characteristics: Vaccination			
VFC provider	3.74 (1.87–7.47)	5.73 (1.31–25.1)	3.80 (1.70–8.54)
Flag charts of vaccine-eligible patients	1.94 (0.99–3.81)	—	—
Use automated electronic medical record prompts ^d	1.59 (0.84–3.01)	—	—
Perform periodic electronic queries to identify vaccine-eligible patients ^d	1.78 (0.86–3.70)	—	—
Other HCP recommends vaccination	1.24 (0.69–2.25)	—	—
Other HCP discusses vaccination	1.28 (0.69–2.35)	—	—
Strategies for first vaccination: None	1.30 (0.92–1.87)	—	—
Strategies for second vaccination: None	1.19 (0.77–1.84)	—	—
Vaccination coordinator	3.06 (1.10–8.52)	1.09 (0.28–4.33)	—

NOTE: Bold ORs and CIs are statistically significant. For vaccination subset, the 5 variables not applied to the multivariable analysis were excluded due to high covariation with one or more variables that were included.

Abbreviation: HCP, health care provider.

^aSimple logistic regression model examined association between individual predictor and outcome variable.

^bLogistic regression model with all significant individual predictors. Sample size is 162.

^cFinal model following backward stepwise regression starting with all significant individual predictors. Sample size is 193.

^dMore than 10% of the data were missing for this predictor.

VFC program reduces physicians' up-front costs by offering free vaccines for VFC-eligible children (53). Providers also do not have to send patients elsewhere to receive vaccines. It is also possible that VFC providers also represent those who are more supportive of vaccines overall and willing to offer in-office vaccination in

their practice setting. Future qualitative interviews with VFC and non-VFC providers may help to further explore these possible explanations.

To our knowledge, our study is among the first to focus specifically on physicians' HPV vaccine recommendations for

Table 3. Physician barriers as predictors of consistent and strong recommendation of HPV vaccine to 11- to 12-year-olds

Physician barrier	Univariate OR (95% CI) ^a	Multivariable AOR (95% CI) ^b	Multivariable AOR (95% CI) ^c
Concerns about vaccine safety	0.32 (0.18–0.56)	0.55 (0.27–1.12)	0.47 (0.25–0.86)
Concerns about vaccine efficacy	0.37 (0.22–0.61)	0.90 (0.45–1.80)	—
Discussing sex	0.41 (0.22–0.75)	—	—
Riskier sex behaviors	0.65 (0.42–1.00)	—	—
Adding another vaccine to schedule	0.35 (0.22–0.56)	0.65 (0.38–1.13)	0.57 (0.34–0.95)
Up-front cost	0.52 (0.40–0.69)	0.84 (0.50–1.41)	—
Insufficient reimbursement	0.55 (0.43–0.71)	0.80 (0.44–1.46)	—
Insufficient insurance coverage	0.63 (0.50–0.80)	1.22 (0.71–2.09)	—
VFC program does not cover males of age >18	0.86 (0.70–1.07)	—	—
Determining insurance coverage	0.62 (0.49–0.79)	1.11 (0.71–1.74)	—
Vaccine expiring before use	0.44 (0.32–0.62)	0.78 (0.49–1.25)	—
Insufficient time to discuss	0.37 (0.24–0.55)	0.86 (0.50–1.45)	—
Remember to discuss	0.28 (0.18–0.45)	0.49 (0.29–0.84)	0.40 (0.25–0.64)
Will not complete the series	0.64 (0.48–0.86)	—	—
Not required by schools	1.01 (0.80–1.28)	—	—
Rarely see adolescent males	0.69 (0.53–0.90)	—	—

NOTE: Bold ORs and CIs represents statistical significance with $\alpha = 0.05/16 = 0.0031$ for univariate analyses and $\alpha = 0.05$ for multivariable analyses.

^aOR with CI from simple logistic regression model for each physician barrier item.

^bMultivariable model with all statistically significant variables from univariate analysis.

^cMultivariable model following backward stepwise regression starting with all statistically significant variables from univariate analysis.

males after the ACIP guidelines for routine vaccination of adolescent males. In addition, this is among the first observational studies to describe the current use of electronic and manual reminders for both physicians and patients related to HPV vaccination. This study has several notable strengths, including a statewide sample of primary care providers and an examination of physicians' recommendation with respect to both consistency and strength. There are also limitations. Our cross-sectional survey design precluded our ability to make causal inferences about variables significantly associated with recommendation of HPV vaccination. In addition, we surveyed physicians from a single state; although this design limits our ability to generalize findings to physicians practicing in our states, it allowed us to focus on HPV vaccine recommendations in a state with relatively high rates of HPV-related disease in males. We provided quantitative "anchors" for our questions regarding consistency but not strength of vaccine recommendation. Physicians may vary in their interpretation of a strong recommendation. Physicians may have reported socially desirable responses regarding practice behaviors; however, the anonymity of the survey likely reduced this bias. Physicians most in favor of HPV vaccination may have completed the survey, possibly providing an overestimate of the proportion that consistently and strongly recommend HPV vaccination. In our sample, 31% of provider-reported HPV vaccine currently is not administered to males in their clinical setting. However, we did not include any follow-up questions as to why they currently do not administer. It is possible that they simply do not vaccinate in their office. Although less likely, they may specifically not offer HPV vaccination for males. Finally, our study was limited to physicians, although other health care providers may recommend HPV vaccination. Thus, study of groups delivering care to adolescent males, such as nurses, medical assistants, and physician assistants, is warranted.

Physician recommendation is key to increasing HPV vaccine coverage; yet physicians' current HPV vaccine recommendations do not align with national guidelines and U.S. health organization recommendations. Interventions are needed to support HPV vaccine recommendation consistency and strength. Our research and others' suggest these interventions also should include education about and strategies as well as policy level interventions to address financial barriers to HPV vaccination. Current efforts to bolster physician recommendation have largely been focused on communication skills and office-based strategies to increase HPV vaccination rates. For example, the CDC's You Are the Key Campaign (54) largely provides patient and provider educational materials and provider communication strategies. More recently, the American Academy of Pediatrics HPV Champion Toolkit (55) has extended resources provided in the You Are the Key Campaign to include tools to facilitate change at the practice level (e.g., electronic health record-based reminder recalls). Our survey

found that Florida physicians' barriers to consistently and strongly recommending HPV vaccine were related to concerns about vaccine safety, concerns about adding vaccines to the vaccine schedule, and difficulty in remembering to discuss HPV vaccination. These findings suggest that future interventions should include components to address these issues at the physician and practice level. By addressing physicians' challenges and supporting their HPV vaccine recommendations to their male patients, we can increase HPV vaccine coverage in Florida and reduce HPV-related disease.

Disclosure of Potential Conflicts of Interest

J.A. Kahn reports receiving a commercial research grant from Merck, Inc. A. Casler has received speakers bureau honoraria from and is a consultant/advisory board member for Merck (Vaccine Division). D. Salmon reports receiving commercial research grants from Pfizer and Crucell and is a consultant/advisory board member for Merck. G.D. Zimet reports receiving commercial research grants from Merck and Roche and is a consultant/advisory board member for Merck. A.R. Giuliano reports receiving a commercial research grant from and is a consultant/advisory board member for Merck & Co. No potential conflicts of interest were disclosed by the other authors.

Authors' Contributions

Conception and design: S.T. Vadaparampil, J.A. Kahn, D. Salmon, R.G. Roetzheim, G.D. Zimet, A.R. Giuliano

Development of methodology: T.L. Malo, J.A. Kahn, R.G. Roetzheim, G.D. Zimet

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): S.T. Vadaparampil, T.L. Malo

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): S.T. Vadaparampil, S.K. Sutton, J.A. Kahn, D. Salmon, G.D. Zimet

Writing, review, and/or revision of the manuscript: S.T. Vadaparampil, T.L. Malo, S.K. Sutton, K.N. Ali, J.A. Kahn, A. Casler, D. Salmon, B. Walkosz, R.G. Roetzheim, G.D. Zimet, A.R. Giuliano

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): S.K. Sutton, K.N. Ali

Study supervision: S.T. Vadaparampil, T.L. Malo

Acknowledgments

We thank Janine Cory and Allison Kennedy Fisher for their survey feedback and Dr. Ji-Hyun Lee for assistance with sample selection.

Grant Support

This research was supported by funding from the Bankhead-Coley Cancer Research Program (4BB10). T.L. Malo is supported by the UNC Lineberger Cancer Control Education Program (R25CA057726). This work was also supported in part by the Biostatistics Core at the H. Lee Moffitt Cancer Center & Research Institute, an NCI-designated Comprehensive Cancer Center (P30-CA076292).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received January 7, 2016; revised July 21, 2016; accepted July 25, 2016; published OnlineFirst August 2, 2016.

References

- Centers for Disease Control and Prevention. Recommendations on the use of quadrivalent human papillomavirus vaccine in males—Advisory Committee on Immunization Practices (ACIP), 2011. *MMWR Morb Mortal Wkly Rep* 2011;60:1705–8.
- Petrosky E, Bocchini JA Jr, Hariri S, Chesson H, Curtis CR, Saraiya M, et al. Use of 9-valent human papillomavirus (HPV) vaccine: updated HPV vaccination recommendations of the advisory committee on immunization practices. *MMWR Morb Mortal Wkly Rep* 2015;64:300–4.
- Reagan-Steiner S, Yankey D, Jeyarajah J, Elam-Evans LD, Singleton JA, Curtis CR, et al. National, regional, state, and selected local area vaccination coverage among adolescents aged 13–17 years - United States, 2014. *MMWR Morb Mortal Wkly Rep* 2015;64:784–92.

4. Cullen KA, Stokley S, Markowitz LE. Uptake of human papillomavirus vaccine among adolescent males and females: Immunization Information System sentinel sites, 2009–2012. *Acad Pediatr* 2014;14:497–504.
5. Centers for Disease Control and Prevention. Human papillomavirus (HPV)-associated cancers, by anatomic site and state – United States, 2004–2008. Available from: <http://www.cdc.gov/cancer/hpv/pdf/HPV-state-rates.pdf>.
6. Giuliano AR, Palefsky JM, Goldstone S, Moreira ED Jr, Penny ME, Aranda C, et al. Efficacy of quadrivalent HPV vaccine against HPV infection and disease in males. *N Engl J Med* 2011;364:401–11.
7. Palefsky JM, Giuliano AR, Goldstone S, Moreira ED Jr, Aranda C, Jessen H, et al. HPV vaccine against anal HPV infection and anal intraepithelial neoplasia. *N Engl J Med* 2011;365:1576–85.
8. Gillison ML, Chaturvedi AK, Lowy DR. HPV prophylactic vaccines and the potential prevention of noncervical cancers in both men and women. *Cancer* 2008;113:3036–46.
9. Marty R, Roze S, Bresse X, Largeron N, Smith-Palmer J. Estimating the clinical benefits of vaccinating boys and girls against HPV-related diseases in Europe. *BMC Cancer* 2013;13:10.
10. Centers for Disease Control and Prevention (CDC). Human papillomavirus-associated cancers - United States, 2004–2008. *MMWR Morb Mortal Wkly Rep* 2012;61:258–61.
11. Machalek DA, Poynten N, Jin F, Fairley CK, Farnsworth A, Garland SM, et al. Anal human papillomavirus infection and associated neoplastic lesions in men who have sex with men: a systematic review and meta-analysis. *Lancet Oncol* 2012;13:487–500.
12. Daling JR, Madeleine MM, Johnson LG, Schwartz SM, Shera KA, Wurscher MA, et al. Human papillomavirus, smoking, and sexual practices in the etiology of anal cancer. *Cancer* 2004;101:270–80.
13. Centers for Disease Control and Prevention. National and state vaccination coverage among adolescents aged 13–17 years – United States, 2011. *MMWR Morb Mortal Wkly Rep* 2012;61:671–7.
14. Brewer NT, Croyle R. President's Cancer Panel: achieving widespread HPV vaccine uptake. Bethesda, MD: National Cancer Institute; 2012.
15. McRee AL, Gilkey MB, Dempsey AF. HPV vaccine hesitancy: findings from a statewide survey of health care providers. *J Pediatr Health Care* 2014;28:541–9.
16. Gilkey MB, Malo TL, Shah PD, Hall ME, Brewer NT. Quality of physician communication about human papillomavirus vaccine: findings from a national survey. *Cancer Epidemiol Biomarkers Prev* 2015;24:1673–9.
17. Soon R, Dela Cruz MR, Tsark JU, Chen JJ, Braun KL. A survey of physicians' attitudes and practices about the human papillomavirus (HPV) vaccine in Hawaii. *Hawaii J Med Public Health* 2015;74:234–41.
18. Allison MA, Hurley LP, Markowitz L, Crane LA, Brtnikova M, Beaty BL, et al. Primary care physicians' perspectives about HPV vaccine. *Pediatrics* 2016;137:1–9.
19. Jaen CR, Stange KC, Nutting PA. Competing demands of primary care: a model for the delivery of clinical preventive services. *J Fam Pract* 1994;38:166–71.
20. Vadaparampil ST, Kahn JA, Salmon D, Lee JH, Quinn GP, Roetzheim R, et al. Missed clinical opportunities: provider recommendations for HPV vaccination for 11–12 year old girls are limited. *Vaccine* 2011;29:8634–41.
21. Vadaparampil ST, Malo TL, Kahn JA, Salmon DA, Lee JH, Quinn GP, et al. Physicians' human papillomavirus vaccine recommendations, 2009 and 2011. *Am J Prev Med* 2014;46:80–4.
22. Vadaparampil ST, Staras SAS, Malo TL, Eddleton KZ, Christie J, Rodriguez M, et al. Provider factors associated with disparities in human papillomavirus vaccination among low-income 9- to 17-year-old girls. *Cancer* 2013;119:621–8.
23. Daley MF, Crane LA, Markowitz LE, Black SR, Beaty BL, Barrow J, et al. Human papillomavirus vaccination practices: a survey of US physicians 18 months after licensure. *Pediatrics* 2010;126:425–33.
24. Rimer B, Harper H, Witte ON. Accelerating HPV vaccine u: urgency for action to prevent cancer. A report to the President of the United States from the President's Cancer Panel. Bethesda, MD: National Cancer Institute; 2014.
25. Freed GL, Nahra TA, Wheeler JR. Counting physicians: inconsistencies in a commonly used source for workforce analysis. *Acad Med* 2006;81:847–52.
26. Kletke PR. Physician workforce data: when the best is not good enough. *Health Serv Res* 2004;39:1251–5.
27. Dillman D. Mail and internet surveys: the tailored design method. New York, NY: Wiley; 2000.
28. Kahn JA, Zimet GD, Bernstein DI, Riedesel JM, Lan D, Huang B, et al. Pediatricians' intention to administer human papillomavirus vaccine: the role of practice characteristics, knowledge, and attitudes. *J Adolesc Health* 2005;37:502–10.
29. Riedesel JM, Rosenthal SL, Zimet GD, Bernstein DI, Huang B, Lan D, et al. Attitudes about human papillomavirus vaccine among family physicians. *J Pediatr Adolesc Gynecol* 2005;18:391–8.
30. Kahn JA, Rosenthal SL, Tissot AM, Bernstein DI, Wetzel C, Zimet GD. Factors influencing pediatricians' intention to recommend human papillomavirus vaccines. *Ambul Pediatr* 2007;7:367–73.
31. Daley MF, Liddon N, Crane LA, Beaty BL, Barrow J, Babbel C, et al. A national survey of pediatrician knowledge and attitudes regarding human papillomavirus vaccination. *Pediatrics* 2006;118:2280–9.
32. Malo TL, Giuliano AR, Kahn JA, Zimet GD, Lee JH, Zhao X, et al. Physicians' human papillomavirus vaccine recommendations in the context of permissive guidelines for male patients: a national study. *Cancer Epidemiol Biomarkers Prev* 2014;23:2126–35.
33. Allison MA, Dunne EF, Markowitz LE, O'Leary ST, Crane LA, Hurley LP, et al. HPV vaccination of boys in primary care practices. *Acad Pediatr* 2013;13:466–74.
34. Malo TL, Ali KN, Sutton SK, Perkins RB, Giuliano AR, Vadaparampil ST. The content and context of physicians' communication with males about human papillomavirus vaccination. *Hum Vaccin Immunother* 2016;12:1511–8.
35. Scherr CL, Augusto B, Ali K, Malo TL, Vadaparampil ST. Provider-reported acceptance and use of the Centers for Disease Control and Prevention messages and materials to support HPV vaccine recommendation for adolescent males. *Vaccine* 2016;34:4229–34.
36. U.S. Department of Health and Human Services. Healthy People 2020. 2014[cited 2014 Jul 1]. Available from: <http://healthypeople.gov/2020/topicsobjectives2020/pdfs/HP2020objectives.pdf>.
37. Ko EM, Missmer S, Johnson NR. Physician attitudes and practice toward human papillomavirus vaccination. *J Low Genit Tract Dis* 2010;14:339–45.
38. Gilkey MB, McRee AL. Provider communication about HPV vaccination: a systematic review. *Hum Vaccin Immunother* 2016;12:1454–68.
39. Gollust SE, LoRusso SM, Nagler RH, Fowler EF. Understanding the role of the news media in HPV vaccine uptake in the United States: synthesis and commentary. *Hum Vaccin Immunother* 2016;12:1430–4.
40. Krakow M, Rogers B. Collateral damage and critical turning points: public health implications of HPV vaccine news coverage for boys and men in 2011. *Health Commun* 2016;31:1081–8.
41. Dornbusch HJ, Stiris T, Del Torso S, Ross-Russell R, Zavrsnik J, Wettergren B, et al. Human papillomavirus vaccination crisis in Japan. *J Paediatr Child Health* 2015;51:1146–7.
42. Hanley SJ, Yoshioka E, Ito Y, Kishi R. HPV vaccination crisis in Japan. *Lancet* 2015;385:2571.
43. Kempe A, Patel MM, Daley MF, Crane LA, Beaty B, Stokley S, et al. Adoption of rotavirus vaccination by pediatricians and family medicine physicians in the United States. *Pediatrics* 2009;124:e809–16.
44. Bratic JS, Seyferth ER, Bocchini JA Jr. Update on barriers to human papillomavirus vaccination and effective strategies to promote vaccine acceptance. *Curr Opin Pediatr* 2016;28:407–12.
45. Reiter PL, Gilkey MB, Brewer NT. HPV vaccination among adolescent males: results from the National Immunization Survey-Teen. *Vaccine* 2013;31:2816–21.
46. Ruffin MITV, Plegue MA, Rockwell PG, Young AP, Patel DA, Yeazel MW. Impact of an electronic health record (EHR) reminder on human papillomavirus (HPV) vaccine initiation and timely completion. *J Am Board Fam Med* 2015;28:324–33.
47. Oliver K, Frawley A, Garland E. HPV vaccination: population approaches for improving rates. *Hum Vaccin Immunother* 2016;12:1589–93.

Vadaparampil et al.

48. Jacobson RM, Agunwamba AA, St Sauver JL, Finney Rutten LJ. The most effective and promising population health strategies to advance human papillomavirus vaccination. *Expert Rev Vaccines* 2016;15:257–69.
49. Centers for Disease Control and Prevention. VFC: for parents. 2012. Available from: <http://www.cdc.gov/vaccines/programs/vfc/parents/default.html>.
50. Lee GM, Santoli JM, Hannan C, Messonnier ML, Sabin JE, Rusinak D, et al. Gaps in vaccine financing for underinsured children in the United States. *J Am Med Assoc* 2007;298:638–43.
51. Davis MM, Ndiaye SM, Freed GL, Clark SJ. One-year uptake of pneumococcal conjugate vaccine: a national survey of family physicians and pediatricians. *J Am Board Fam Pract* 2003;16:363–71.
52. Mennito SH, Darden PM. Impact of practice policies on pediatric immunization rates. *J Pediatr* 2010;156:618–22.
53. Centers for Disease Control and Prevention. VFC healthcare providers information flyer. 2014; Available from: <http://www.cdc.gov/vaccines/programs/vfc/providers/questions/qa-flyer-hcp.html>.
54. Centers for Disease Control and Prevention. HPV vaccine resources for healthcare Professionals. 2015; Available from: <http://www.cdc.gov/vaccines/who/teens/for-hcp/hpv-resources.html>.
55. American Academy of Pediatrics. HPV Champion Toolkit. 2015. Available from: <https://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Pages/HPV-Champion-Toolkit.aspx>.

Cancer Epidemiology, Biomarkers & Prevention

Missing the Target for Routine Human Papillomavirus Vaccination: Consistent and Strong Physician Recommendations Are Lacking for 11 to 12-Year-Old Males

Susan T. Vadaparampil, Teri L. Malo, Steven K. Sutton, et al.

Cancer Epidemiol Biomarkers Prev Published OnlineFirst August 2, 2016.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-15-1294](https://doi.org/10.1158/1055-9965.EPI-15-1294)

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link <http://cebp.aacrjournals.org/content/early/2016/09/12/1055-9965.EPI-15-1294>. Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.