

Supplemental table 4. Polymorphisms in the MBL2 gene among Caucasians exposed to secondhand smoke in the Mayo clinic study

	Cases n	Controls n	<u>OR</u> (95% CI)	P-Value
MBL2-01				
<i>G/G</i>	67	103	1.00 (reference)	
<i>G/A</i>	29	30	1.62 (0.88-2.96)	0.12
<i>A/A</i>	2	2	1.43 (0.19-10.83)	0.73
MBL2-02				
<i>C/C</i>	97	131	1.00 (reference)	
<i>C/T</i>	1	4	0.36 (0.04-3.38)	0.37
<i>T/T</i>	0	0	-	-
MBL2-03				
<i>G/G</i>	90	118	1.00 (reference)	
<i>G/A</i>	7	17	0.57 (0.22-1.45)	0.24
<i>A/A</i>	1	0	-	-
MBL2-11				
<i>C/C</i>	49	51	1.00 (reference)	
<i>C/G</i>	40	57	2.11 (0.89-4.95)	0.09
<i>G/G</i>	9	27	2.88 (1.23-6.74)	0.02
MBL2-12				
<i>G/G</i>	50	87	1.00 (reference)	
<i>G/C</i>	44	43	1.91 (1.08-3.37)	0.03
<i>C/C</i>	4	5	1.37 (0.34-5.42)	0.66
MBL2-25				
<i>C/C</i>	59	84	1.00 (reference)	
<i>C/T</i>	34	46	1.04 (0.59-1.84)	0.89
<i>T/T</i>	5	5	1.26 (0.34-4.64)	0.73

*Odds ratios are adjusted for age, gender, exposure to adult secondhand smoke, first degree relative with lung cancer, and education level